



## SHORT COMMUNICATION

# Effect of Starch Noodle (*Dangmyeon*) and Pork Intestines on the Rehydration Stability of Korean Blood Sausage (*Sundae*)

Youngmin Kim, Hyejin Jang, Sangdong Lim, and Sangpil Hong\*

Korea Food Research Institute, Wanju 55365, Korea

## OPEN ACCESS

**Received** July 16, 2020  
**Revised** October 26, 2020  
**Accepted** October 28, 2020

\*Corresponding author : Sangpil Hong  
Korea Food Research Institute,  
Wanju 55365, Korea  
Tel: +82-63-219-9098  
Fax: +82-63-219-9876  
E-mail: [spHong@kfri.re.kr](mailto:spHong@kfri.re.kr)

\*ORCID  
Youngmin Kim  
<https://orcid.org/0000-0002-7570-7284>  
Hyejin Jang  
<https://orcid.org/0000-0002-1485-9196>  
Sangdong Lim  
<https://orcid.org/0000-0002-1500-4413>  
Sangpil Hong  
<https://orcid.org/0000-0002-4060-0129>

**Abstract** This study was conducted to examine the effects of starch noodles (*dangmyeon*; SNs) with different starch sources and porcine intestines (PIs) with different pH on the rehydration stability of Korean blood sausage (*sundae*). Mungbean SN3 and PI3 (pH 9.18) showed significantly higher values of 80.69%–91.67% and 79.66%–80.98%, respectively, regardless of the drying methods (hot air, vacuum and freeze drying) ( $p < 0.05$ ). A number of larger pores were observed only in the cross-section of the freeze dried SN and PI through SEM. SN2 (potato starch) and PI3 (pH 9.18) showed lower expansion ( $^{\circ}\Delta L$  6.90 mm) and higher expansion ratio ( $^{\circ}\Delta L$  26.29 mm), respectively, after rehydration of freeze dried sample ( $p < 0.05$ ). From the application of SN2 (potato starch) and PI (0.5%–2.0% Na-pyrophosphate) to freeze dried *sundae* manufacturing, higher rehydration stability of more than 91.5% was obtained. These results suggested that potato SN and treatment of PI with Na-pyrophosphate is useful for desirable rehydration stability of freeze dried *sundae*.

**Keywords** *sundae*, Korean blood sausage, starch noodle, porcine intestines, rehydration stability

## Introduction

Sausages are a meat product that is usually made from ground meat (usually pork, but sometimes beef or poultry), salt, spices and other flavorings (Caldironi and Ockerman, 1982; Oh, 2012). Many countries produce similar types of sausages even though the materials and methods used are different (Caldironi and Ockerman, 1982; Oh, 2012). Among the many different sausage products, blood sausage (*sundae* in Korean) is a popular ready-to-eat food in Korea (Choi et al., 2015; Son et al., 1999), which is similar to western blood sausage, although the manufacturing methods differ (Silva et al., 2013).

*Sundae* is generally prepared by steaming porcine intestines (PIs) stuffed with various ingredients such as minced pork, starch noodles (SNs), pork blood, and vegetables (Park, 2017). The many different *sundae* products tend to be named after

their city of origin or main ingredient, i.e. Pyongando (SNs), Hamgyeongdo (bigger), Gaesung (pork), Byungcheon (rich ingredients), Baekam (ground ingredients), Ojingeo *sundae* (squid) and Chapssal *sundae* (glutinous rice), with each type having its own special characteristics and manufacturing method (Oh et al., 2012). However, when cooking *sundae* or rehydrating dried *sundae* as a convenient home meal replacement (HMR) product, swelling of the inner part and contraction of the outer part often occur simultaneously, resulting in an undesirable shape (Kim et al., 2019). These problems are attributable to the physicochemical properties of the inner SNs and outer PI with which *sundae* is made (Kim et al., 2019).

SNs, also called cellophane or glass noodles, are routinely used as one of the main ingredients of *sundae*, and are derived from sweet potatoes, called *dangmyeon* in Korea (Chen et al., 2002).

However, many kinds of SNs have also been made from corn, sweet potato, potato, and tapioca, which can have different physicochemical properties that influence the swelling or spreadability of *sundae* (Chen, 2003; Lee et al., 2002; Yook and Lee, 2001). In this study, SN products from sweet potato, potato and mungbean were chosen as one of main experimental materials because they have different starch composition (Chen, 2003; Yook and Lee, 2001), which can show different expansion effect on the *sundae*.

PI, a natural casing material, is usually used to pack the inner material of *sundae*. Kim et al. (2019) tried to change the contraction properties of PI by controlling its pH in order to reduce the contraction of PI that occurs when hot water is added to rehydrate dried *sundae*. Choe and Kim (2016) showed that the addition of natural vinegar to swelled pig skin increased moisture content and decreased hardness in *sundae*, though acid was commonly known to lower the quality of meat products (Kim and Lim, 1994). On the other hand, lower or higher pH from the pI (isoelectric point) of beef improved its water-retention capacity, followed by an increase in the yields and tenderness of marinated beef after cooking (Hong et al., 2013). Based on the above studies, it is expected that water retention capacity or tenderness of PI can be increased like pig skin or beef through adjustment of pH to acidic or alkaline.

The purpose of this study is to examine the rehydration stability of Korean blood sausage (*sundae*) according to the swelling and contraction characteristics of SNs and PI, in the cooking, drying and rehydration.

## Materials and Methods

### Materials

The main ingredients of *sundae*, including pork, pork blood, PI, and SNs, were purchased from a local market in the Jeonju area of Korea. SN products were classified into sweet potato (Ottogi, Anyang, Korea), potato (Qingdao richsun trading, Qingdao, China), and mung bean (Qingdao richsun trading) ones. Samples of SN products were cut into strips measuring 3–5 cm and boiled for 6 min, and then left to cool immediately.

The raw ingredients such as pork, PI and porcine blood were stored at  $-80^{\circ}\text{C}$  (RT50H6120SL, Samsung, Suwon, Korea) and defrosted before use. The PI was used after removing the fat layer and washing it with distilled water. Citric acid (Edentown FNB, Incheon, Korea) and sodium pyrophosphate (ESfood, Gunpo, Korea) were used for the pH control.

### Manufacturing *Sundae*

Table 1 shows the recipe for *sundae*. The inner part ingredients were mixed and stuffed into the PI using a stuffing machine (SV-5, Newtech, Daegu, Korea). The diameter of the stuffing machine used to make the casing was 25 mm. The error range of the casing diameter of manufactured *sundae* was within 5%. The samples were heated at  $80^{\circ}\text{C}$  for 50 min in a

**Table 1. Formulation of *Sundae***

Ingredients	Contents (%)
Pork	44.5
Porcine intestine	16.0
Starch noodles	7.5
Pork blood	2.5
Starch	5.0
Isolated soybean protein	4.0
Vegetable <sup>1)</sup>	17.3
Seasoning <sup>2)</sup>	3.2
Total	100

<sup>1)</sup> Vegetable included the following: onion, 6.0%; green onion, 5.0%; carrot, 5.0%; garlic, 1%; and ginger, 0.25%.

<sup>2)</sup> Seasoning included the following: salt, 1.0%; sugar, 1.0%; condiment, 1.0%; and pepper, 0.25%.

water bath (SB-9, Eyela, Tokyo, Japan) and then left to cool for 30 min at room temperature. The manufactured *sundae* was cut into the proper size and stored at  $-80^{\circ}\text{C}$  or freeze-dried under 20 Pa for 24 h.

### Dry yields

Samples were freeze-dried (FDU-1200, Eyela, Tokyo, Japan), hot air-dried (HK-DO1000F, Hankuk S&I, Hwaseong, Korea) and vacuum-dried (DRV622DA, Advantec, Tokyo, Japan) for 24 h. Hot air drying and vacuum drying were performed at a temperature of  $60^{\circ}\text{C}$  and  $40^{\circ}\text{C}$ , respectively, with the degree of the vacuum set at 80 kPa. Each sample was weighed before and after drying, and the dry yields were calculated as a percentage of the pre-drying weight.

### SEM (Scanning Electron Microscopy)

SEM (Scanning Electron Microscopy, Hitachi S4700, Tokyo, Japan) was used to analyze the swelling effect of the samples treated by freeze, hot air and vacuum drying for 24 h. The samples were fixed to SEM stubs with double-sided tape, then coated with a layer of gold with a thickness of 10 nm. All of the samples were observed at an accelerating voltage of 20.0 kV with a magnification level of  $\times 1,500$ .

### Length and thickness

Vernier calipers (CD-15CPX, Mitutoyo, Japan) was used to measure changes in the length ( $\Delta L$ ) and thickness ( $\Delta T$ ) of the dried samples before and after heating them, and in the length ( $^*\Delta L$ ) and thickness ( $^*\Delta T$ ) of the freeze-dried samples before and after heating them.

### pH

The PI samples were soaked in a pretreatment solution (distilled water, 2% citric acid and 0.5%–2% sodium pyrophosphate) at  $4^{\circ}\text{C}$  for 2 h at a ratio of 1:10. The pH of each solution was measured using a pH meter (S210, Mettler Toledo, Worthington, OH, USA). The pretreated samples were washed 3 times with distilled water and then stocked at  $4^{\circ}\text{C}$  until use in experiments. The pH of the PI was measured after homogenizing 5 g samples with 20 mL of distilled water at 12,000 rpm for 1 min in a

homogenizer (PT1200E, Kinematica, Luzern, Switzerland).

### Rehydration stability

The freeze dried *sundae* cut into a proper size was rehydrated in a beaker with 100 mL of boiling distilled water, and width of the PI part was measured every min for 3 min. The rehydration stability was expressed in relative ratio (%) of width of PI part before and after rehydration.

### Statistical analysis

The experiment was replicated 3 times, and SPSS Statistics (ver. 23, IBM, Armonk, NY, USA) were used for the statistical analysis. One-way ANOVA was performed for the significant test between each sample, and Duncan's multiple range test was performed to determine whether there were any significant differences between the mean values ( $p < 0.05$ ).

## Results and Discussion

### Dry yields

The freeze, hot air and vacuum drying methods are well known as common and conventional techniques. Hot air drying is a technique by heated air (Celma et al., 2009) and vacuum drying is a method by reducing pressure and drying at low temperature (Baslar et al., 2014; Wu et al., 2007) while freeze drying (FD) is a technique for drying foods at low temperatures and under high vacuum (Karam et al., 2016). These drying methods have their own characteristics, and may influence drying yields or quality according to drying conditions such as kinds of sample, temperature and time. Table 2 shows the dry yields of the SN and PI samples after freeze, hot air and vacuum drying. The dry yields of the SN ranged from 60.02% to 91.67%. Jung et al. (1991) reported a water-binding capacity of 77% for mung bean starch, 79% for potato starch, and 83% for sweet potato starch. In addition, the water content of starch when isolated from SNs was as follows: 11.6% for mung bean starch, 8.9% for potato starch, and 8.7% for sweet potato starch (Chen, 2003). In our results, however, SN3 (mung bean starch) showed a significant difference in dry yield, ranging from 80.69% to 91.67%, when compared to SN1 (sweet potato starch, 60.96%–64.62%) and SN2 (potato starch, 60.13%–60.96%), regardless of the drying method. This indicates that the different properties of the SN products used in this experiment resulted in significant differences between the dry yields.

On the other hand, the dry yields of the PI samples ranged from 65.23% to 93.95%. There was a significant difference in PI2 (citric acid), PI3 (sodium pyrophosphate) and PI1 (control, distilled water), regardless of the drying method. These results indicate that the water-holding capacity and degree of expansion of the PI2 and PI3 samples are relatively higher than the control (PI1) due to the pH shifting from the isoelectric point (Kim et al., 2000). Hong et al. (2013) showed that shifting of pH from pI (isoelectric point) of beef improved the water-holding capacity by as much as 9%–15%, followed by an increase in the yield and tenderness of marinated beef after cooking.

Therefore, the above results correlate with the high water-holding capacity resulting from the pH shift from the isoelectric points and the humectant effect achieved by reducing water loss during drying.

### pH

pH is an important factor that affects the quality of meat, including its freshness, water-holding capacity, tenderness, binding ability, color and texture (Hong et al., 2013; Yim et al., 2016). The pH value of PI2 with 2% citric acid was 2.37, and

**Table 2.** Dry yield of SN and PI according to various dry methods

Samples		Dry yield (%)
Freeze drying	SN1 <sup>1)</sup>	62.02±5.75 <sup>b</sup>
	SN2	60.35±7.14 <sup>b</sup>
	SN3	80.69±2.29 <sup>a</sup>
	PI1 <sup>2)</sup>	67.12±0.45 <sup>c</sup>
	PI2	91.98±0.23 <sup>a</sup>
	PI3	79.66±2.66 <sup>b</sup>
Hot air drying	SN1	64.62±6.00 <sup>b</sup>
	SN2	60.96±5.12 <sup>b</sup>
	SN3	91.67±8.33 <sup>a</sup>
	PI1	67.62±0.19 <sup>c</sup>
	PI2	92.18±0.41 <sup>a</sup>
	PI3	80.98±1.40 <sup>b</sup>
Vacuum drying	SN1	60.96±5.69 <sup>b</sup>
	SN2	60.13±4.51 <sup>b</sup>
	SN3	90.00±8.82 <sup>a</sup>
	PI1	65.23±1.40 <sup>c</sup>
	PI2	93.95±5.27 <sup>a</sup>
	PI3	79.68±2.73 <sup>b</sup>

<sup>1)</sup> SN1, sweet potato starch; SN2, potato starch; SN3, mung bean starch.

<sup>2)</sup> PI1, distilled water; PI2, 2% citric acid; PI3, 2% sodium pyrophosphate.

<sup>a-c</sup> Means on the same column with different letters are significantly different ( $p < 0.05$ ).

SN, starch noodle; PI, porcine intestine.

that of PI3 with 2% sodium pyrophosphate was 9.18, while that of PI1 with distilled water was 6.13 (data not shown), suggesting different levels of water-holding capacity or tenderness. The accumulation of met-myoglobin is faster in meat with a low pH than in meat with high pH condition (Ledward, 1985; Owen and Lawrie, 1975).

In general, the pH value of raw meat used to make ham and sausages is 5.8–6.2, so it is very difficult to produce satisfactory products in terms of quality when meat with a pH of 5.7 or less is used (Kim and Lim, 1994). Hong et al. (2013) indicated that lower or higher pH than the pI (isoelectric point) of beef improved the water-holding capacity, yield and tenderness of marinated beef after cooking.

Therefore, PI3 (pH 9.18) was expected to be the most advantageous in terms of improving the rehydration stability of *sundae*.

### Length and thickness

The degree of expansion of the SN and the degree of contraction of the PI, which differently used of SN products and the respective PI treatments, were compared by observing changes in the length and thickness of boiled and rehydrated SN and PI samples after freeze-drying. As shown in Table 3, SN3 ( $\Delta L$  7.72 mm) showed a significant difference from SN1 ( $\Delta L$  5.03 mm) and SN2 ( $\Delta L$  5.39 mm) after boiling, and this pattern was also similar to the changes of length observed when SN samples were rehydrated after freeze-drying where SN3 ( $\Delta L^*$  6.90 mm) was significantly longest, compared to SN1 ( $\Delta L^*$

**Table 3. Relative changes in the length and thickness of SN**

Samples	Relative change (mm)			
	$\Delta L$	* $\Delta L$	$\Delta T$	* $\Delta T$
SN1 <sup>1)</sup>	5.03±0.41 <sup>b</sup>	1.81±2.01 <sup>b</sup>	0.03±0.02 <sup>a</sup>	0.22±0.03 <sup>a</sup>
SN2	5.39±1.53 <sup>b</sup>	1.47±0.41 <sup>b</sup>	0.03±0.02 <sup>a</sup>	0.20±0.16 <sup>a</sup>
SN3	7.72±1.48 <sup>a</sup>	6.90±0.05 <sup>a</sup>	0.05±0.01 <sup>a</sup>	0.14±0.01 <sup>a</sup>

<sup>1)</sup> SN1, sweet potato starch; SN2, potato starch; SN3, mung bean starch.

<sup>a,b</sup> Means on the same column with different letters are significantly different ( $p < 0.05$ ).

SN, starch noodle.

1.81 mm) and SN2 ( $\Delta L^*$  1.47 mm), respectively. This means that SN3 has a stronger expansion property than SN1 and SN2. On the other hand, a significant difference was not observed in the changes of thickness among the SN samples after boiling or after rehydration of the freeze-dried samples. These results may be due to the relatively lower changes.

In the case of the PI samples (Table 4), after soaking treatment, the length of PI3 was observed to expand significantly more (42.15 mm) than PI1 and PI 2 at 24.77 mm and 25.90 mm, respectively. These expansion levels were also maintained in the freeze-drying step. On the other hand, when the freeze-dried PI samples were rehydrated with hot water, PI3 contracted to 26.29 mm in length, while PI1 and PI2 contracted to 15.63 mm and 21.02 mm, respectively. Even though the PI3 samples showed a relatively higher degree of contraction, the overall length of PI3 (16.29 mm) is relatively longer than that of PI1 (5.63 mm) and PI2 (11.02 mm). These results are interesting and may seem controversial, but if *sundae* is manufactured under the same conditions, the PI3 sample with a higher expansion capability is expected to be advantageous in terms of maintaining the shape of freeze-dried *sundae* during rehydration.

## SEM

SEM images were analyzed to observe a cross section of SN composed of different starch and PI samples were treated by various drying methods. As shown in Fig. 1, freeze-dried SN (A-a, B-a, C-a) retained a number of pores regardless of the type of SNs, while hot air dried (A-b, B-b, C-b) and vacuum dried (A-c, B-c, C-c) SN showed different closed patterns, instead of pores on the surface of the noodles. Among the freeze-dried samples, SN1 (A-a) retained more pores than SN2 (B-a) or SN3 (C-a), and showed a different pattern from the other two samples. Wang et al. (2010) reported that potato SNs were tighter, and thus absorbed less water. From the above results, it seems advantageous to use potato (SN2, B-a) or mung bean (SN3, C-a) SNs for their rehydration stability because they are able to inhibit the swelling of starchy tissues due to their reduced water absorption.

**Table 4. Changes in the length of PI at each step of treatment**

Samples	Treatment steps (mm)					Remarks
	Before treatment	Soaking	Boiling	Freezing drying	Rehydration	
PI1 <sup>1)</sup>	10.00 <sup>a</sup>	24.77±3.81 <sup>ax</sup>	24.16±3.81 <sup>bx</sup>	21.97±4.73 <sup>bx</sup>	15.63±1.69 <sup>bx</sup>	5.63±1.69 <sup>x</sup>
PI2	10.00 <sup>a</sup>	25.90±2.37 <sup>abx</sup>	25.31±2.39 <sup>abx</sup>	23.64±1.65 <sup>abx</sup>	21.02±3.76 <sup>bxy</sup>	10.96±3.76 <sup>xy</sup>
PI3	10.00 <sup>a</sup>	42.15±6.67 <sup>by</sup>	41.23±6.60 <sup>cy</sup>	37.35±6.43 <sup>cy</sup>	26.29±5.54 <sup>cy</sup>	16.29±5.54 <sup>y</sup>


<sup>1)</sup> PI1, distilled water; PI2, 2% citric acid; PI3, 2% sodium pyrophosphate.

<sup>a-c</sup> Means on the same row with different letters are significantly different ( $p < 0.05$ ).

<sup>x-y</sup> Means on the same column with different letters are significantly different ( $p < 0.05$ ).

PI, porcine intestine.





**Fig. 1.** SEM images of dried SN and PI. A, SN1; B, SN2; C, SN3; D, PI1; E, PI2; F, PI3; a, freeze dried; b, hot air dried; c, vacuum dried; SEM, scanning electron microscopy; SN, starch noodles; PI, porcine intestine; SN1, sweet potato starch; SN2, potato starch; SN3, mung bean starch; PI1, distilled water; PI2, 2% citric acid; PI3, 2% sodium pyrophosphate.

Overall, the freeze-dried PI samples also show a similar pattern to the SEM images of the dried SN ones. However, the size of the pores of the PI2 (E-a) and PI3 (F-a) samples is relatively larger than that of the PI1 (D-a, distilled water treatment) samples. The water-holding capacity of meat is lowest at pH 5.4–5.6 near the isoelectric point (pI), and increases as the pH shifts further away from the isoelectric point (Hong et al., 2013; Kim et al., 2000; Korea Food Research Institute, 2020). The tissue of PI treated with citric acid and sodium pyrophosphate is supposed to become looser and cause concomitant swelling, and keeps tissue in a loose state even after drying. As shown in Fig. 1, the larger pores are observed from the PI2 (E-a) and PI3 (F-a). This means that citric acid or sodium pyrophosphate affects the water-holding capacity of PI, which is responsible for the production of pores after freeze-drying. On the other hand, no clear pores were observed in the cross sections of any of the samples after air-drying and vacuum drying. Son et al. (2014) reported that the better the water-holding capacity of a sample, the better its rehydration stability after freeze-drying.

From the above results, the freeze-drying method seems to be advantageous for the production of dried *sundae* with improved rehydration stability.

### Rehydration stability

Based on the above results, freeze-dried *sundae* was manufactured using SN2 composed of potato starch and PI treated with various concentrations of sodium pyrophosphate in order to obtain the optimum concentration available for commercial use.

In consideration of the Korean Food Standard Codex (2020), the PI was treated at a maximum concentration of up to 2.0% sodium pyrophosphate and the rehydration stability of *sundae* was measured by the relative width changes of PI, compared to control.

As shown in Table 5, unlike the control (S1) showing a lower rehydration level (72.5% by 180 s), all of the treated samples maintained a rehydration stability of more than 91.5%, and this increased to 95.5% at 180 s when the concentration of sodium pyrophosphate was increased to 2.0%.

Judging from the above mentioned results, it is concluded that SN2 made with PI treated with more than 0.5% sodium pyrophosphate is advantageous for the production of *sundae* with desirable rehydration stability.

**Table 5.** Relative rehydration stability of *sundae* formulated with various levels of sodium pyrophosphate based on the length changes of PI

Samples	Relative rehydration (%)			
	0 s	60 s	120 s	180 s
S1 <sup>1)</sup>	100 <sup>a</sup>	76.2±5.7 <sup>bx</sup>	74.8±4.5 <sup>bx</sup>	72.5±4.4 <sup>bx</sup>
S2	100 <sup>a</sup>	92.6±4.8 <sup>aby</sup>	91.7±4.4 <sup>by</sup>	91.5±4.5 <sup>by</sup>
S3	100 <sup>a</sup>	93.0±4.5 <sup>by</sup>	92.7±1.6 <sup>by</sup>	92.5±1.9 <sup>by</sup>
S4	100	95.5±3.5 <sup>y</sup>	95.0±3.3 <sup>y</sup>	95.0±3.3 <sup>y</sup>
S5	100 <sup>a</sup>	96.9±2.1 <sup>aby</sup>	95.7±1.8 <sup>by</sup>	95.5±2.0 <sup>by</sup>

<sup>1)</sup> S1, *Sundae* (PI with 0% Na-pyrophosphate); S2, *Sundae* (PI with 0.5% Na-pyrophosphate); S3, *Sundae* (PI with 1.0% Na-pyrophosphate); S4, *Sundae* (PI with 1.5% Na-pyrophosphate); S5, *Sundae* (PI with 2.0% Na-pyrophosphate).

<sup>a,b</sup> Means on the same row with different letters are significantly different ( $p < 0.05$ ).

<sup>x,y</sup> Means on the same column with different letters are significantly different ( $p < 0.05$ ).

PI, porcine intestine.



## Conclusion

Expansion of SNs and contraction of PI were influenced by starch sources and pH difference, respectively. The SN from potato (SN2) and alkaline PI with sodium pyrophosphate (PI3, pH 9.18) showed relatively lower expansion and higher expansion ratio during rehydration, respectively. The rehydration stability was achieved more than 91.5% through application of SN2 and PI (0.5%–2.0% Na-pyrophosphate) to *sundae* manufacturing. Therefore, use of potato SN and PI treated with sodium pyrophosphate seems to be advantageous for strengthening rehydration stability of freeze dried *sundae*.

## Conflicts of Interest

The authors declare no potential conflicts of interest.

## Acknowledgements

This research was supported by the main research program of the Korea Food Research Institute (E0201100-01) and by a grant from Ministry of SMEs and Startups (GN181100-01).

## Author Contributions

Conceptualization: Hong SP. Data curation: Kim YM, Jang HJ. Formal analysis: Lim SD. Methodology: Kim YM. Writing - original draft: Kim YM. Writing - review & editing: Kim YM, Jang HJ, Lim SD, Hong SP.

## Ethics Approval

This article does not require IRB/IACUC approval because there are no human and animal participants.

## References

- Baslar M, Kılıçlı M, Toker OS, Sağdıç O, Arici M. 2014. Ultrasonic vacuum drying technique as a novel process for shortening the drying period for beef and chicken meats. *Innov Food Sci Emerg Technol* 26:182-190.
- Caldironi HA, Ockerman HW. 1982. Incorporation of blood proteins into sausage. *J Food Sci* 47:405-408.
- Celma AR, Cuadro F, López rodríguez F. 2009. Characterization of industrial tomato by-products from infrared drying process. *Food Bioprod Process* 87:282-291.
- Chen Z. 2003. Physicochemical properties of sweet potato starches and their application in noodle products. Ph.D. dissertation, Wageningen Univ., Wageningen, Netherlands.
- Chen Z, Sagis L, Legger A, Linssen JPH, Schols HA, Voragen AGJ. 2002. Evaluation of starch noodles made from three typical Chinese sweet potato starches. *J Food Sci* 67:3342-3347.
- Choe JH, Kim HY. 2016. Effects of swelled pig skin with various natural vinegars on quality characteristics of traditional Korean blood sausages (*Sundae*). *Food Sci Biotechnol* 25:1605-1611.
- Choi YS, Sung JM, Jeon KH, Choi HW, Seo DH, Kim CJ, Kim HW, Hwang KE, Kim YB. 2015. Quality characteristics on adding blood levels to blood sausage. *Korean J Food Cook Sci* 31:741-748.

- Hong SP, Kim YH, Lee NH, Heo YU. 2013. The physicochemical characteristics of marinated beef Galbi under different cooking conditions. *Korean J Food Cult* 28:78-88.
- Jung SH, Shin GJ, Choi CU. 1991. Comparison of physicochemical properties of corn, sweet potato, wheat and mungbean starches. *Korean J Food Sci Technol* 23:272-275.
- Karam MC, Petit J, Zimmer D, Djantou EB, Scher J. 2016. Effects of drying and grinding in production of fruit and vegetable powders: A review. *J Food Eng* 188:32-49.
- Kim CJ, Lim SC. 1994. Effects of tumbling condition and curing method on the quality of turkey drumstick. *Korean J Food Sci Anim Resour* 14:37-40.
- Kim YB, Rho JH, Richardson I, Wood J. 2000. Comparison of physicochemical properties of pork from 4 different pig breeds. *Korean J Anim Sci* 42:195-202.
- Kim YM, Lee SR, Jang HJ, Lim SD, Oh YC, Hong SP. 2019. Study on the development of HMR (RTE) product using *Sundae* (Korean sausage). 2019 KFN International Symposium and Annual Meeting of the Korean Society of Food Science and Nutrition, ICC, Jeju, Korea. p 328.
- Korean Food Research Institute. KFRI u-food quality information system. Available from: <http://db.ufood.kr/index.jsp>. Accessed at Jan 29, 2020.
- Ledward DA. 1985. Post-slaughter influences on the formation of metmyoglobin in beef muscles. *Meat Sci* 15:149-171.
- Lee YC, Oh SW, Han SB, Han SD, Kang N. 2002. Properties of dangmyuns using different starches and freeze dried dangmyuns. *Korean J Food Sci Technol* 34:24-29.
- Ministry of Food and Drug Safety. Korean Food Additives Standard Codex. Available from: [https://www.foodsafetykorea.go.kr/foodcode/04\\_03.jsp?idx=4](https://www.foodsafetykorea.go.kr/foodcode/04_03.jsp?idx=4). Accessed at Jan 29, 2020.
- Oh JH, Lee EJ, Kim KH, Yook HS. 2012. Characteristics of Korean blood sausages (*Soondae*) for globalization. *Food Ind Nutr* 17:23-26.
- Oh SD. 2012. A literature review on the types and cooking methods of *Soondae* during the Joseon dynasty. *Korean J Food Cult* 27:340-345.
- Owen JE, Lawrie RA. 1975. The effect of an artificially induced high pH (hydrogen ion concentration) on the susceptibility of minced porcine muscle to undergo oxidative rancidity under frozen storage. *J Food Sci Technol* 10:169-180.
- Park KL. 2017. A literature review of the recipes for pork: A focus on recipe data published in Korea from the Joseon dynasty to modern times. *Korean J Human Ecol* 26:471-493.
- Silva FAP, Amaral DS, Guerra ICD, Dalmás PS, Arcanjo NMO, Bezerra TKA, Beltrão Filho EM, Moreira RT, Mandruga MS. 2013. The chemical and sensory qualities of smoked blood sausage made with edible by-products of goat slaughter. *Meat Sci* 94:34-38.
- Son JW, Yum CA, Kim CJ. 1999. Effects of binding materials on sensory and cooking properties of Soondae. *Korean J Food Cook Sci* 15:191-196.
- Son YJ, Choi SY, Lee KY, Huang Y, Yoo KM, Hwang IK. 2014. Study on major Korean protein sources before and after freeze drying processing. *Korean J Food Cook Sci* 30:64-75.
- Wang M, Chen C, Sun G, Wang W, Fang H. 2010. Effects of curdlan on the color, syneresis, cooking qualities, and textural properties of potato starch noodles. *Starch Stärke* 62:429-434.
- Wu L, Orikasa T, Ogawa Y, Tagawa A. 2007. Vacuum drying characteristics of eggplants. *J Food Eng* 83:422-429.
- Yim DG, Yang MR, No GR, Choi DS, Jang HM, Kim TY, Jo JW, Yang SC, Kim SW, Kim IS. 2016. Quality characteristics

of the meat products reached expiration date in Korean market. *J Agric Life Sci* 50:169-177.

Yook C, Lee WK. 2001. Production of starch vermicelli (*Dangmyun*) by using modified corn starches (I): Physicochemical properties of domestic and foreign starch vermicelli (*Dangmyun*). *Korean J Food Sci Technol* 33:60-65.