

REVIEW

Effects of Pulsed Electric Field on Meat Tenderization and Microbial Decontamination: A Review

Se-Ho Jeong, Han-Beak Lee, and Dong-Un Lee*

Department of Food Science and Biotechnology, Chung-Ang University, Anseong 17546, Korea

 OPEN ACCESS

Received October 4, 2023

Revised November 23, 2023

Accepted December 19, 2023

***Corresponding author** : Dong-Un Lee
 Department of Food Science and Biotechnology, Chung-Ang University, Anseong 17546, Korea
 Tel: +82-31-670-3034
 E-mail: dong-un.lee@cau.ac.kr

***ORCID**
 Se-Ho Jeong
<https://orcid.org/0000-0003-4158-1203>
 Han-Beak Lee
<https://orcid.org/0009-0002-0475-0311>
 Dong-Un Lee
<https://orcid.org/0000-0002-7967-1792>

Abstract This review sought to categorize studies on meat tenderization and safety through pulsed electric field (PEF) treatment, with a particular focus on reconciling conflicting findings regarding the tenderization effect (i.e., the primary outcome of PEF treatment) and to discuss the underlying mechanisms of these effects. While the tenderization effect may vary depending on the homogeneity of PEF treatment and variations in the conditions of texture measurements, the protein associated with tenderization was degraded by PEF treatment in most studies. PEF technology enables the delivery of a high voltage for a brief duration, typically in the microsecond range, making it a non-thermal technology. One of the distinct advantages of PEF is its ability to preserve the freshness of meat due to its exceptionally short treatment time. While PEF studies have traditionally centered on pasteurizing liquid foods, research on its application to meat is steadily expanding. Therefore, this review aims to elucidate the mechanisms of PEF and provide current insights into the applications of this technology for meat tenderization and microbial inactivation.

Keywords pulsed electric field, meat, tenderization, proteolysis, meat safety

Introduction

Pulsed electric field (PEF) treatment is a non-thermal technology used in food processing, which is characterized by low heat generation, short treatment duration, and low energy consumption (Knorr and Angersbach, 1998). Unlike other electric-based technologies such as ohmic heating, PEF selectively affects the cell membrane during microsecond pulses in a square wave form (Gavahian and Farahnaky, 2018; Kumar et al., 2015). These properties allow for gentle food processing without the adverse effects of heat generation (Barsotti and Cheftel, 1999). Therefore, PEF offers advantages in preserving food characteristics that are generally associated with freshness such as flavor, taste, color, and nutrients.

Various non-thermal technologies have been extensively used in the food industry, including gamma irradiation, plasma, UV-light, pulsed light (PL), and high-pressure

processing (HPP). From a bactericidal perspective, gamma irradiation is considered an ideal technology, albeit with some limitations, such as negative public perception and the development of persistent rancidity in high-fat foods (Farkas et al., 2002). Furthermore, the bactericidal effects of plasma, UV-light, and PL are limited to surface-attached microbes (Levy et al., 2012; Soro et al., 2023; Sruthi et al., 2022). HPP is among the most widely commercialized non-thermal technologies and can be applied to both liquid and solid foods, even after packaging (Denoya et al., 2015; Perera et al., 2010). This technology also offers advantages in minimizing heat-induced quality deterioration and has a short treatment duration (Nath et al., 2023). However, HPP has several limitations compared to PEF in terms of productivity because it operates only in batch mode and requires a huge installation space and costs (Balasubramaniam et al., 2015; Sampedro et al., 2014). PEF, on the other hand, can be used in continuous mode and can easily switch to batch mode while retaining the advantages of HPP. Therefore, PEF has emerged as a next-generation non-thermal technology.

PEF can be employed for various purposes, such as controlling texture, facilitating mass transfer, aiding in extraction, and pasteurization, depending on the treatment intensity (Naliyadhara et al., 2022; Zare et al., 2023; Zhang et al., 2023). While most previous research on PEF has focused on liquid and plant-based foods, studies on its application to meat have been relatively limited. The limited application of PEF to meat compared to plant-based products is largely due to the high fat and protein content in meat. Nevertheless, the meat industry continually seeks novel technologies to ensure high-quality meat products. Meat tenderization and safety are key factors influencing meat quality and consumer acceptability. However, the effects of PEF on meat remains controversial due to variations in outcomes depending on the treatment conditions and evaluation methods. Therefore, this review sought to classify findings related to meat tenderization and safety and discuss the discrepancies between these findings.

General Description of Pulsed Electric Field

Electroporation induced by pulsed electric field treatment

The tenderizing and bactericidal effects of PEF are generally attributed to cell membrane breakdown induced by electroporation (Zimmermann, 1996). When a cell is exposed to an electric field, micropores are simultaneously created in the phospholipid bilayer of the cell membrane, resulting in increased cell membrane permeability. The cell membrane, which consists of a phospholipid bilayer naturally has a specific transmembrane potential of approximately 10 mV between its inner and outer layers. When the membrane is exposed to an external electric field, the potential induced by the electric field increases until it reaches a critical value. As the potential increases, the membrane reorganizes the charges on both sides, leading to a viscoelastic deformation of the membrane due to the attraction of charges with opposite polarity. This process continues until the critical value is reached. If the external electric field surpasses the critical value of the transmembrane potential, an electro-compression force is generated on both sides of the membrane, initiating the formation of pores.

Pulsed electric field treatment system

Detailed information on the PEF system and its components is provided in Fig. 1A. The PEF system typically consists of two main components: the pulse generator and the treatment chamber. The inclusion of other subcomponents depends on whether the system operates in continuous mode (for both liquid and solid samples) or batch mode (single batch). In the continuous mode, a pump or conveyor is necessary to transport the sample to the treatment chamber. Treatment chambers come in three common types: parallel plates, co-axial, and co-linear chambers (Arshad et al., 2020). For the treatment of




Fig. 1. Description of PEF system and components (A) and parameters of PEF treatment and oscilloscope image (B). PEF, pulsed electric field.

liquid food in continuous mode, heat exchangers are additionally used to preheat the sample before treatment and cool it down afterward (Toepfl et al., 2014). In contrast, the single batch mode does not require these additional components. The batch chamber is similar to the parallel plates chamber but lacks the flow of the sample. This type of chamber is typically used for solid foods and is suitable for bench-scale applications. PEF treatment for solid foods is conducted in the batch chamber, which is filled with a low-conductive medium such as tap water. Unlike continuous chambers, batch chambers have been modified in various forms (Jeong et al., 2023; Li et al., 2020; Mok et al., 2017). The adjustable chamber illustrated in Fig. 1A was designed to allow for the adjustment of electrode gap (Jeong et al., 2023). This setup can be used to treat samples

without the need for a medium and enables direct contact with the chamber without space between the electrode and the sample. While the use of batch chambers is advantageous for tailoring conditions at the bench scale, it is important to note that, in contrast to the continuous mode, treatment in batch mode is more susceptible to raising the sample temperature due to the high electric field strength (Toepfl et al., 2014). Therefore, batch treatment should be carried out at an acceptable level of PEF intensity to minimize heating.

Parameters of pulsed electric field treatment

The effect of PEF on tissue is determined by both the properties of the sample and the PEF parameters. From the perspective of suitability for PEF processing, the electrical conductivity of the samples is a crucial factor because it influences temperature increases (Athmaselvi et al., 2014). The parameters in the PEF system are detailed in Fig. 1B. The electric field strength (E, kV/cm) is a primary factor that determines the extent of the PEF effect. The electric field strength is calculated using the following Eq. (1):

$$\text{Electric field strength (E)} = \frac{\text{Output voltage (kV)}}{\text{Distance of electrodes gap (cm)}} \quad (1)$$

A sufficient field strength is necessary to reach the threshold for entering an electroporated state (irreversible state). The required level varies for each sample and depends on the cell size. Solid foods with large cells typically require a relatively lower field strength for electroporation compared to microbes. The pulse width, measured in microseconds, represents the duration during which the high voltage is maintained for each pulse in the form of a square wave. Short pulse widths are beneficial in preventing temperature increases during PEF treatment. The pulse frequency is a factor that determines the treatment time in terms of the number of pulses. These parameters can be expressed as a specific energy (Q, kJ/kg) using the following Eq. (2) (Zhang et al., 1995):

$$\text{Specific energy (Q)} = \frac{V^2 t}{Rm} \quad (2)$$

where V is the output voltage (kV), t is the treatment time (number of pulses/frequency), R is the resistance (ohm), and m is the mass of sample (kg). In the case of PEF treatments for solid foods, field strength is commonly used as a variable rather than specific energy, as the pulse width and treatment time are often held at fixed values.

Action of Pulsed Electric Field on the Tissue

Rupture of tissue by pulsed electric field

The mechanism of electroporation can be theoretically explained; however, the evidence of cell membrane permeability is not entirely clear at the cellular level. This uncertainty arises because, while we can observe the irreversible state of the cell disrupted by PEF, we cannot easily observe the reversible state. At the tissue level, such as in vegetables or meat with a heterogeneous distribution of cell sizes, observing the reversible state can be quite challenging. The effects of PEF on tissues are observed as macroscopic phenomena, unlike at the cellular level, and these phenomena are dependent on both PEF

intensity and tissue properties. When tissue is exposed to PEF, some cells within the tissue are damaged by the electric field. The distribution of damaged cells varies depending on the intensity and cell size. In other words, damaged and intact cells coexist heterogeneously. The ratio of damaged cells can be assessed by measuring an electric signal that results from ion release from the cells. This signal change is expressed in terms of electric conductivity (S/m). The PEF treatment selectively affects the cell membrane inside the tissue, leading to changes in ion release. Morphologically, it can be challenging to distinguish between intact and treated tissue. For example, in the microstructure of PEF-treated tissue, the cell wall may appear intact, whereas the cell membrane and organelles may exhibit some degree of damage (Kim et al., 2019). Similarly, in muscle tissue where there is no cell wall, morphological differences can be difficult to discern. This selective effect can result in a more flexible texture, caused by the release of turgor pressure within the cell. Therefore, PEF can soften the texture and improve processing suitability. Furthermore, the damaged cell membrane can create porosity in the tissue, increasing mass transport channels and accelerating mass transfer. Damaged organelles, such as vacuoles, can also facilitate the extraction of bio compounds.

Evaluation of permeability

Lebovka et al. (2002) introduced an effective tool for assessing the degree of tissue rupture using conductivity. The authors proposed evaluating the degree of rupture caused by PEF treatment through the cell disintegration index Z (CDI), which compares intact cells with completely ruptured cells. This tool has gained widespread acceptance in evaluating tissue permeability in numerous studies. Similar to the CDI concept, other evaluation tools have been developed to assess relative disintegration of mass transfer, texture, and acoustics. These methods provide a representation of macroscopic change of the tissue rather than focusing on the cellular level.

The underlying mechanisms through which PEF influences microscopic phenomena, as well as at the cellular level, remain unclear. However, recent efforts have been made to substantiate existing hypotheses. Genovese et al. (2021) utilized non-destructive MRI analysis to observe changes in moisture distribution in plant tissues after PEF treatment. Plant tissues exhibit heterogeneous moisture distribution due to variations in cell size. This heterogeneity leads to varying effects of PEF on each cell. Irreversible deformation induced by PEF results in a homogeneous distribution of moisture, which is confirmed through T_2 mapping and MRI tomography (Genovese et al., 2023). Some ruptured cells release cytoplasm into intracellular voids, leading to a homogeneous moisture distribution as the cytoplasm fills the voids. This phenomenon alters the osmotic pressure in adjacent cells. This non-destructive analysis significantly enhances our understanding of cell permeabilization and complements conductivity analysis.

Tenderization Effects of Pulsed Electric Field Treatment

Conflicting views on the tenderization effects of pulsed electric field treatment

Tenderness is a critical factor when evaluating sensory quality, and the primary effect of PEF on meat is its tenderization. The degree of tenderization at a macroscopic scale is typically assessed through mechanical measurements such as shear force or meat hardness. However, various studies utilizing different meat materials, PEF intensities, and sample sizes have produced conflicting results regarding the tenderization effects of PEF. Tables 1 and 2 categorize these studies based on the presence or absence of a tenderization effect following PEF treatment. The studies listed in Table 1 demonstrate that PEF treatment did not lead to a significant tenderization effect. These studies share common characteristics, including the use of

Table 1. Effect of PEF on the tenderization of different muscles, with little effects observed

Meat materials	PEF condition	Sample status	Findings	References
<i>Semimembranosus</i> (beef)	20 μ s, 90 Hz, 0.36 kV/cm 20 μ s, 20 Hz, 0.60 kV/cm The treatment time or number of pulses were not provided	Post-rigor, sample size was trimmed fitted in the batch chamber (13 \times 8 \times 5 cm), direct contact with the electrodes	No significant reduction in shear force	Bhat et al. (2019a)
<i>Longissimus et lumborum</i> (beef)	20 μ s, 200 Hz, 0.23 kV/cm 20 μ s, 200 Hz, 0.68 kV/cm Total treatment time: 30 s	Post-rigor, sample size was trimmed fitted in the batch chamber (13 \times 8 \times 5 cm), direct contact with the electrodes	No significant reduction in shear force	Khan et al. (2017)
<i>Semitendinosus</i> (beef)	20 μ s, 50 Hz, 1.40 kV/cm, 1,032 pulses	Post-rigor, sample size was trimmed fitted in a triangular batch chamber (6 \times 4 \times 6 cm), direct contact the electrodes	No significant reduction in shear force	Faridnia et al. (2015)
Breast meat (turkey)	20 μ s, 10 to 110 Hz, 1.25 to 2.0 kV/cm, 100 to 300 pulses	Post-rigor, sample size was trimmed to 6 \times 2 \times 2 cm, Direct contact with the electrodes	No significant reduction in shear force	Arroyo et al. (2015a)
<i>Longissimus thoracis et lumborum</i> (beef)	20 μ s, 10 Hz, 1.40 kV/cm, 300 and 600 pulses	Post-rigor, sample size was trimmed to 6 \times 2 \times 2 cm, direct contact with the electrodes	No significant reduction in shear force	Arroyo et al. (2015b)
<i>Longissimus thoracis</i> (beef)	20 μ s, 1 to 50 Hz, 0.20 to 0.56 kV/cm, 30 to 1,528 pulses	Post-rigor, sample size was trimmed to 10 \times 8 \times 4 cm, direct contact with the electrodes	No significant reduction in shear force	Faridnia et al. (2014)
<i>Longissimus thoracis et lumborum</i> (pork)	20 μ s, 100 and 200 Hz, 1.20 and 2.30 kV/cm, 150 and 300 pulses	Post-rigor, sample size was trimmed to 6 \times 2 \times 2 cm, direct contact with the electrodes	No significant reduction in hardness and chewiness	McDonnell et al. (2014)
<i>Semitendinosus</i> (beef)	20 μ s, 152 Hz, 1.10 kV/cm, 152 pulses 20 μ s, 200 Hz, 1.50 kV/cm, 200 pulses 20 μ s, 65 Hz, 1.90 kV/cm, 250 pulses 20 μ s, 5 Hz, 2.80 kV/cm, 300 pulses	Post-rigor, sample size was trimmed to 6 \times 2 \times 2 cm, direct contact with the electrodes	No significant reduction in shear force	O'Dowd et al. (2013)

PEF, pulsed electric field.

commercial PEF systems and direct contact with the electrodes. Interestingly, low PEF intensity did not result in a tenderization effect (Bhat et al., 2019a; Khan et al., 2017). Similarly, studies employing high-intensity PEF treatment, which should be sufficient for cell disruption, did not impact meat tenderness (Arroyo et al., 2015a; Arroyo et al., 2015b; Faridnia et al., 2014; Faridnia et al., 2015; McDonnell et al., 2014; O'Dowd et al., 2013). In contrast, in the case of plant tissue, treatment with an electric field strength of approximately 1.0 kV/cm resulted in increased flexibility and softness in texture (Fauster et al., 2018; Lee et al., 2022). These results may be because the PEF effect is masked when using excessively large sample sizes for treatment and measuring texture at excessively high temperatures.

Large sample sizes require a considerable amount of energy to reach the necessary PEF intensity for tenderization. Additionally, the outcome is influenced by sample properties such as fat composition or connective tissue. Bhat et al. (2019a)

Table 2. Effect of PEF on the tenderization of different muscles

Meat materials	PEF condition	Sample status	Findings	References
<i>Semitendinosus</i> (beef)	30 μ s, 50 Hz, 0.5 to 2.0 kV/cm, 100 pulses	Post-rigor, sample size was trimmed to 2 \times 1 \times 1 cm, direct contact with the electrodes	Significant reduction of shear force, hardness, and chewiness of raw meat	Jeong et al. (2023)
<i>Transversus thoracis</i> muscle (beef)	20 μ s, 50 Hz, 0.7 kV/cm, 1,600 and 5,200 pulses	Post-rigor, sample was cut into 5 cm width pieces, treated in a batch chamber filled with agar pads as medium	Significant reduction of hardness in high-intensity PEF-treated meat after sous-vide cooking	Karki et al. (2023)
<i>Semitendinosus</i> (beef)	20 μ s, 50 Hz, 1.0 to 2.0 kV/cm, 200 pulses	Post-rigor, sample size was trimmed to 6 \times 3 \times 3 cm, treated in a batch chamber filled with tap water	Significant reduction of shear force and hardness both in raw and sous-vide cooked meat	Jeong et al. (2020)
<i>Longissimus lumborum</i> and <i>semimembranosus</i> (beef)	Repeated treatment (1 \times , 2 \times , 3 \times) 20 μ s, 90 Hz, 1.25 kV/cm Total treatment time: 30s	Pre-rigor, sample size was trimmed fitted in the batch chamber (13 \times 8 \times 5 cm), direct contact with the electrodes	Significant reduction of shear force on the <i>semimembranosus</i>	Bekhit et al. (2016)
<i>Biceps femoris</i> (beef)	20 μ s, 50 Hz, 1.7 kV/cm	Post-rigor, sample size was trimmed fitted in a triangular batch chamber (6 \times 4 \times 6 cm), direct contact with the electrodes	Significant reduction of shear force at all aging times	Faridnia et al. (2016)
<i>Longissimus lumborum</i> and <i>semimembranosus</i> (beef)	Repeated treatment (1 \times , 2 \times , 3 \times) 20 μ s, 90 Hz, 1.25 kV/cm Treatment time or number of pulses are not described	Post-rigor, sample size was trimmed fitted in the batch chamber (13 \times 8 \times 5 cm), direct contact with the electrodes	Significant reduction of shear force on the <i>Longissimus lumborum</i>	Suwandy et al. (2015c)
<i>Longissimus lumborum</i> and <i>semimembranosus</i> (beef)	Repeated treatment (1 \times , 2 \times , 3 \times) 20 μ s, 90 Hz, 1.25 kV/cm Total treatment time: 30s	Pre-rigor, sample size was trimmed fitted in the batch chamber (13 \times 8 \times 5 cm), direct contact with the electrodes	Significant reduction of shear force at all aging times	Suwandy et al. (2015b)
<i>Longissimus lumborum</i> and <i>semimembranosus</i> (beef)	Repeated treatment (1 \times , 2 \times , 3 \times) 20 μ s, 90 Hz, 0.625 and 1.25 kV/cm	Post-rigor sample size was trimmed fitted in the batch chamber (13 \times 8 \times 5 cm), direct contact with the electrodes	Significant reduction of shear force at all aging times	Suwandy et al. (2015a)
<i>Longissimus lumborum</i> and <i>semimembranosus</i> (beef)	Repeated treatment (1 \times , 2 \times , 3 \times) 20 μ s, 20 to 90 Hz, 0.27 to 0.56 kV/cm 606 to 2,724 pulses	Post-rigor, sample size was trimmed fitted in the batch chamber (13 \times 8 \times 5 cm), direct contact with the electrodes	Significant reduction of shear force at all aging times	Bekhit et al. (2014)

PEF, pulsed electric field.

and Khan et al. (2017) did not observe a significant reduction in shear force due to insufficient PEF intensity. However, Suwandy et al. (2015c) observed a tenderization effect with a sufficient PEF intensity, even when using samples of the same size as the former studies. A sufficient PEF intensity can ensure that the cell permeability induced by PEF is homogeneous. Generally, the presence or absence of a PEF effect is indirectly validated through changes in electric signals. Sufficient PEF intensity increases the electric conductivity of muscle tissue because damaged cells release intracellular fluid, which is an electrolyte. However, it can be challenging to determine whether a homogeneous PEF effect has been achieved for each sample immediately after PEF treatment, as some previous studies did not provide this information (Arroyo et al., 2015a; Arroyo et al., 2015b; McDonnell et al., 2014). Conversely, in a recent study, Jeong et al. (2023) used smaller sample sizes and clearly observed a positive linear relationship between PEF intensity and electric conductivity. While this relationship can be observed through changes in texture, an increase in conductivity can also occur due to changes in microstructure, regardless of the tenderization effect. According to O'Dowd et al. (2013), PEF treatment reduced the diameter of muscle fiber bundles and increased conductivity. Similar results were reported by Khan et al. (2017).

Most of the studies listed in Tables 1 and 2 heated the meat samples for measuring shear force or texture profiles. While Jeong et al. (2023) assessed the texture of raw beef *Semitendinosus* muscle, it is important to note that eating raw meat is not a common practice worldwide, except in some East Asian countries. Additionally, the texture of raw meat can vary depending on its temperature. Due to these factors, meat texture is generally evaluated using cooked meat, typically prepared in a water bath at 80°C until the internal temperature reaches approximately 70°C. However, this cooking process can potentially negate the effects of PEF treatment. Connective tissues in meat tend to contract at temperatures over 65°C (Tornberg, 2005), resulting in a tougher texture. Tenderization by PEF may not be observed when the PEF effect is not uniformly distributed throughout the meat. This phenomenon can be confirmed by comparing the studies listed in Tables 1 and 2, as they provide evidence of this effect. Furthermore, studies that have explored the combination of PEF and sous-vide cooking have observed a tenderization effect (Jeong et al., 2020; Karki et al., 2023). According to Jeong et al. (2020), the PEF effect observed in raw beef *Semitendinosus* muscle was maintained even after sous-vide cooking (60°C, 1 h to 24 h). Karki et al. (2023) conducted the PEF treatment and sous-vide cooking (60°C, 24 h and 36 h) on short ribs. The high-energy PEF-treated (0.7 kV/cm, 5,200 pulses) and longer-cooked (36 h) samples exhibited a significant reduction in tenderness compared to the control ($p < 0.001$).

Proteolysis action induced by pulsed electric field treatment

Several studies have demonstrated that proteolysis is facilitated by PEF treatment, regardless of the mechanical texture measurements. Table 3 summarizes the findings of studies on proteolysis induced by PEF, which were approached from three different perspectives: the level of myofibril structure (Chian et al., 2019; Mungure et al., 2020; O'Dowd et al., 2013), structural proteins (Bhat et al., 2019b; Bhat et al., 2019c; Jeong et al., 2023; Suwandy et al., 2015a; Suwandy et al., 2015b), and myofibrillar protein (MP) extracts (Dong et al., 2020; Wang et al., 2022).

In a study by O'Dowd et al. (2013), myofibrils were extracted through a four-step filtration process, and their particle size was measured. The final filtration fraction, from which connective tissues and debris had been removed, exhibited a smaller particle size in the PEF-treated sample compared to the control and heated samples. PEF treatment also resulted in the rupture and fragmentation of myofibril structure (Chian et al., 2019; Mungure et al., 2020). While these phenomena could be observed under a microscope, further measurements are needed to examine finer changes at the level of structural proteins. The degradation of troponin-T and desmin has been associated with tenderization (Marino et al., 2013; Yates et al., 1983). Calpain activity also plays a crucial role in meat protein degradation (Coria et al., 2018). Jeong et al. (2023) investigated the

Table 3. Effect of PEF on the proteolysis of different muscles

Meat materials	PEF condition	Target protein	Findings	References
<i>Semitendinosus</i> (beef)	30 μ s, 50 Hz, 0.5 to 2.0 kV/cm, 100 pulses	μ -Calpain, troponin-T	μ -Calpain was not reduced regardless of PEF intensity, however, troponin-T was proportionally reduced by increases of PEF intensity.	Jeong et al. (2023)
Myofibrillar protein (MP) extraction of <i>longissimus lumborum</i> (pork)	6 μ s, 500 Hz, 2.5 to 7.5 kV/cm, 15,000 to 45,000 pulses	MP	Microstructure of MP was deformed, and the hydrophobicity of MP surface was enhanced by PEF treatment.	Wang et al. (2022)
MP extraction from the <i>pectoralis</i> (chicken)	600 to 1,000 Hz, 8 to 28 kV/cm, pulse width and number of pulses are not described	MP	The solubility and hydrophobicity of MP were increased, and the dynamic rheological properties were enhanced by PEF treatment.	Dong et al. (2020)
<i>Longissimus lumborum</i> (deer)	20 μ s, 50 Hz, 2.5 to 10 kV, treatment time or number of pulses are not described	Myofibril	The myofibril was fragmented and ruptured along the z-line by PEF treatment.	Mungure et al. (2020)
<i>Longissimus thoracis</i> (beef)	20 μ s, 50 Hz, 1.0 and 1.25 kV/cm, 500 and 2,000 pulses	Myosin, actin, collagen, myofibril	Thermal profiles of the myosin, actin, and collagen were not affected by PEF treatment. The myofibrillar structure was ruptured by PEF treatment.	Chian et al. (2019)
<i>Longissimus dorsi</i> (deer)	20 μ s, 50 and 90 Hz, 0.2 and 0.5 kV/cm, 1,500 and 2,700 pulses	Calpain, desmin, troponin-T	The calpain activity was slightly increased by PEF treatment. The degradation of desmin and troponin-T was observed in PEF treated sample.	Bhat et al. (2019c)
<i>Longissimus dorsi</i> (deer)	20 μ s, 50 and 90 Hz, 0.3125 and 1.25 kV/cm, 1,500 and 2,700 pulses	Meat protein Free amino acids	The digestibility and solubility of meat protein were increased by PEF treatment. The concentration of free amino acids was higher in the PEF treated sample.	Bhat et al. (2019b)
<i>Longissimus lumborum</i> and <i>semimembranosus</i> (beef)	20 to 90 Hz, 0.625 and 1.25 kV/cm, pulse width and number of pulses are not provided	Troponin-T Desmin	The degradation of desmin and troponin-T was increased by PEF treatment.	Suwandy et al. (2015a)
<i>Longissimus lumborum</i> and <i>semimembranosus</i> (beef)	20 to 90 Hz, 5 and 10 kV/cm, pulse width and number of pulses are not provided	Troponin-T Desmin	The degradation of desmin and troponin-T was increased by PEF treatment.	Suwandy et al. (2015b)
<i>Semitendinosus</i> (beef)	20 μ s, 5 to 200 Hz, 1.1 to 2.8 kV/cm, 152 to 300 pulses	Myofibril fragment	The particle size of filtered myofibril fragment was decreased by PEF treatment.	O'Dowd et al. (2013)

PEF, pulsed electric field.

proteolysis effect of PEF treatment (0.5 to 2.0 kV/cm) and found that PEF treatment did not significantly affect μ -calpain but resulted in a significant decrease in troponin-T. Calpain activity was slightly increased by PEF treatment in deer *longissimus lumborum* muscle (Bhat et al., 2019c). Likewise, troponin-T and desmin were degraded by PEF on different muscles (Bhat et al., 2019b; Bhat et al., 2019c; Suwandy et al., 2015a; Suwandy et al., 2015b). Dong et al. (2020) and Wang et al. (2022) used MP extracts to observe structural changes in the protein, rather than focusing on a specific protein. Unlike other studies, they

treated the samples with high-intensity PEF exceeding 10 kV/cm. Their results indicated that the microstructure of MP was deformed, and the hydrophobicity of the MP surface was enhanced by PEF. These structural changes induced modifications in the rheological properties of MP (Wang et al., 2022).

In addition to meat protein degradation, changes in structure and increased permeability induced by PEF treatment can facilitate the aging process. As evidence of the tenderization effect, PEF treatment may not produce immediate results in the absence of an aging process (O'Dowd et al., 2013), but its effect becomes evident with extended aging periods (Suwandy et al., 2015a; Suwandy et al., 2015b). Therefore, in addition to mechanical texture properties, PEF treatment can facilitate tenderization through proteolysis. Further research should explore approaches in which PEF improves end-product properties as a pre-treatment and investigates whether the PEF effect persists after subsequent processes or enhances the efficiency of the process.

Use of Pulsed Electric Field for the Microbial Inactivation of Meat

The majority of studies on the microbial inactivation effect of PEF have been conducted on liquid food due to the higher intensity of PEF required for electrical breakdown of microorganisms compared to solid food. Smaller cell sizes in microorganisms require higher intensities to induce membrane breakdown. Typically, intensities exceeding 10 kV/cm are required for liquid food. However, meat can begin to cook when such high intensities are applied, which is similar to ohmic heating. Although the microbicidal effect on meat is limited for this reason, various studies on meat safety have consistently been conducted and are summarized in Table 4. The studies in Table 4 examined microbial stability over different storage periods (Aşık-Canbaz et al., 2022; Faridnia et al., 2015), contamination that may occur during PEF treatment (Bhat et al., 2020), and the direct effects of pasteurization (Alahakoon et al., 2019; Haughton et al., 2012; McDonnell et al., 2014; Stachelska et al., 2012).

Aşık-Canbaz et al. (2022) investigated the impact of moderate-intensity PEF (MIPEF) that is under 10 kV/cm level of electric field strength on the cold storage stability of chicken breast. MIPEF treatment extended the limit for total mesophilic aerobic bacteria counts by up to two more days. Moreover, total coliform bacteria were reduced by 2 Log CFU/g in MIPEF-treated samples. However, the counts of *Listeria monocytogenes* and *Pseudomonas aeruginosa* remained equivalent to the control group. Similarly, Faridnia et al. (2015) found that the growth of aerobic bacteria in fresh semitendinosus meat did not differ between PEF-treated and control samples over 7 days. However, in frozen-thawed meat, aerobic bacteria increased by 2 Log CFU/g in PEF-treated samples at 7 days. No microbial contamination occurred due to PEF treatment (Bhat et al., 2020), suggesting that PEF is a suitable technology for meat processing in terms of safety.

Although previous studies have shown stability in terms of storage and contamination following PEF treatment, the lethal effect on microorganisms was either not observed or only minimally apparent. Alahakoon et al. (2019), Haughton et al. (2012), and McDonnell et al. (2014) did not observe significant microbial inactivation effects. Alahakoon et al. (2019) and McDonnell et al. (2014) conducted their studies with insufficient PEF intensity for microbial inactivation. Haughton et al. (2012) inoculated *Yersinia enterocolitica* in meat (approximately 5–7 Log CFU/g) and applied relatively high PEF voltages (3.75 and 15 kV/cm). Despite applying a sufficient PEF intensity, the reduction in microbial counts was less than 1 Log CFU/g. Since PEF treatment for meat is typically performed at cold temperatures, achieving a direct pasteurization effect is challenging. Even in liquid food, the pasteurization effect diminishes at input temperatures below 25°C. Therefore, the microbial inactivation effect of PEF on meat is relatively lower than that observed in liquid food due to limited available voltage and the requirement for the meat to be in a cold state.

Table 4. Effect of PEF on the microbial inactivation of different muscles

Meat materials	PEF condition	Target microbes	Findings	References
Chicken breast fillets	0.2 ms, 4.67, and 7 kV/cm Continuous treatment during storage period	Total mesophilic aero bacteria (TMAB), total coliform bacteria (TCB) <i>Listeria monocytogenes</i> , <i>Pseudomonas aeruginosa</i>	The limit for TMAB counts was exceeded in PEF-treated samples 2 days later than the control samples. The TCB counts were reduced with 2 Log CFU/g in PEF-treated samples. The counts of <i>L. monocytogenes</i> and <i>P. aeruginosa</i> were maintained in 7 kV/cm PEF treated samples during the storage periods.	Aşık-Canbaz et al. (2022)
Beef jerky	20 µs, 20 Hz, 0.52 kV/cm, 606 pulses	Total plate count, coliform yeast and mold	The microbial contamination by PEF treatment was not observed.	Bhat et al. (2020)
Deep and superficial pectoral (beef)	20 µs, 50 Hz, 0.7 and 1.5 kV/cm, 1,030 to 6,400 pulses	Lactic acid bacteria Total aerobic bacteria	The microbial inactivation effect was not observed after PEF pretreatment.	Alahakoon et al. (2019)
<i>Semitenidinosus</i> (beef)	20 µs, 50 Hz, 1.4 kV/cm, 1,032 pulses	Total aerobic bacteria	The growth of aerobic bacteria in fresh meat was not different between PEF-treated and untreated meat over the course of 7 days. The growth of aerobic bacteria in frozen-thawed meat increased 2 Log CFU/g in PEF treated meat at 7 days.	Faridnia et al. (2015)
<i>Longissimus thoracis et lumborum</i> (pork)	20 µs, 100 and 200 Hz, 1.2 and 2.3 kV/cm, 150 and 300 pulses	Total viable count (TVC)	The TVC was not affected by PEF treatment.	McDonnell et al. (2014)
Minced beef meat	28 to 2,800 MHz, 300 V/m, treatment time: 15 mins, pulse width not described	<i>Yersinia enterocolitica</i>	The <i>Y. enterocolitica</i> inoculated in the meat was reduced under 1 Log CFU/g.	Stachelska et al. (2012)
Chicken breast	10 µs, 5 Hz, 3.75 and 15 kV/cm, treatment time: 10 to 30 sec	<i>Campylobacter jejuni</i> , <i>Escherichia coli</i> , <i>Salmonella enteridis</i> , <i>Enterobacteriaceae</i> , total viable counts	The microbes were unaffected by PEF treatment.	Haughton et al. (2012)

PEF, pulsed electric field.

Considerations of Pulsed Electric Field Treatment on the Meat

PEF technology is considered a non-thermal processing method. However, it can result in slight temperature increases during the process. Therefore, most PEF studies on meat have focused on changes in meat quality following PEF treatment. Various factors affecting meat quality, such as color, purge loss, cooking loss, and lipid oxidation, have been the primary areas of investigation. In most studies, meat color was not significantly affected by PEF treatment, and although some studies observed a reduction in CIE a*, these changes were not distinguishable to the naked eye (Jeong et al., 2020; Jeong et al., 2023). Purge loss and cooking loss experienced slight increases due to changes in microstructure induced by PEF treatment (McDonnell et al., 2014; O'Dowd et al., 2013). Additionally, lipid oxidation was found to increase with higher PEF intensities in some studies (Alahakoon et al., 2019; Kantono et al., 2021). These observations highlight the importance of determining an appropriate PEF intensity level that does not compromise meat quality.

Conclusion

PEF technology can have various effects on food depending on the intensity used. In the case of meat, PEF studies have primarily focused on the textural effects of tenderization, although there have been conflicting findings in this regard. These discrepancies can be attributed to the homogeneity of PEF treatment and variations in the conditions of mechanical measurements. However, despite these conflicting results, most studies have seem to agree that PEF can facilitate tenderization through proteolysis. Compared to its effect on liquid foods, the microbial inactivation effect of PEF on meat is relatively limited because the required voltage to achieve a lethal effect is often not reached due to the inherent properties of meat. Additionally, PEF intensity must be carefully modulated to prevent the loss of meat quality, such as lipid oxidation. To overcome these limitations, future research in this area should focus on developing chambers or systems that can more consistently and stably process meat via PEF treatment. Additionally, establishing reference points to ensure the homogeneity of PEF treatment in meat processing will be important to promote the widespread adoption of this technology in the meat industry.

Conflicts of Interest

The authors declare no potential conflicts of interest.

Acknowledgements

This research was supported by the Chung-Ang University Research Scholarship Grants in 2021. This work was support by Korea Institute of Planning and Evaluation for Technology in Food, Agriculture, Forestry (IPET) through High Value-added Food Technology Development Program funded by Ministry of Agriculture, Food, and Rural Affairs (MAFRA) (332018-04).

Author Contributions

Conceptualization: Jeong SH, Lee DU. Investigation: Jeong SH, Lee HB. Writing - original draft: Jeong SH. Writing - review & editing: Jeong SH, Lee HB, Lee DU.

Ethics Approval

This article does not require IRB/IACUC approval because there are no human and animal participants.

References

- Alahakoon AU, Oey I, Bremer P, Silcock P. 2019. Quality and safety considerations of incorporating post-PEF ageing into the pulsed electric fields and sous vide processing chain. *Food Bioprocess Technol* 12:852-864.
- Arroyo C, Eslami S, Brunton NP, Arimi JM, Noci F, Lyng JG. 2015a. An assessment of the impact of pulsed electric fields processing factors on oxidation, color, texture, and sensory attributes of turkey breast meat. *Poult Sci* 94:1088-1095.
- Arroyo C, Lascorz D, O'Dowd L, Noci F, Arimi J, Lyng JG. 2015b. Effect of pulsed electric field treatments at various stages

- during conditioning on quality attributes of beef *longissimus thoracis et lumborum* muscle. *Meat Sci* 99:52-59.
- Arshad RN, Abdul-Malek Z, Ahmad MH, Buntat Z, Nawawi Z, Kumara CLGP, Abdulameer AZ, Sidik MAB. 2020. Coaxial treatment chamber for liquid food treatment through pulsed electric field. *Indones J Electr Eng Comput Sci* 19:1169-1176.
- Aşık-Canbaz E, Çömlekçi S, Can Seydim A. 2022. Effect of moderate intensity pulsed electric field on shelf-life of chicken breast meat. *Br Poult Sci* 63:641-649.
- Athmaselvi KA, Viswanathan R, Balasubramanian M, Roy I. 2014. The effects of concentration and type of electrode on electrical conductivity of guava pulp during ohmic heating. *J Food Res Technol* 2:113-123.
- Balasubramaniam VM, Martinez-Montegudo SI, Gupta R. 2015. Principles and application of high pressure-Based technologies in the food industry. *Annu Rev Food Sci Technol* 6:435-462.
- Barsotti L, Cheftel JC. 1999. Food processing by pulsed electric fields: II. Biological aspects. *Food Rev Int* 15:181-213.
- Bekhit AEDA, Suwandy V, Carne A, van de Ven R, Hopkins DL. 2016. Effect of repeated pulsed electric field treatment on the quality of hot-boned beef loins and topsides. *Meat Sci* 111:139-146.
- Bekhit AEDA, van de Ven R, Suwandy V, Fahri F, Hopkins DL. 2014. Effect of pulsed electric field treatment on cold-boned muscles of different potential tenderness. *Food Bioprocess Technol* 7:3136-3146.
- Bhat ZF, Morton JD, Mason SL, Bekhit AEDA. 2019a. Does pulsed electric field have a potential to improve the quality of beef from older animals and how? *Innov Food Sci Emerg Technol* 56:102194.
- Bhat ZF, Morton JD, Mason SL, Bekhit AEDA. 2020. The application of pulsed electric field as a sodium reducing strategy for meat products. *Food Chem* 306:125622.
- Bhat ZF, Morton JD, Mason SL, Bekhit AEDA, Mungure TE. 2019b. Pulsed electric field: Effect on *in-vitro* simulated gastrointestinal protein digestion of deer *Longissimus dorsi*. *Food Res Int* 120:793-799.
- Bhat ZF, Morton JD, Mason SL, Mungure TE, Jayawardena SR, Bekhit AEDA. 2019c. Effect of pulsed electric field on calpain activity and proteolysis of venison. *Innov Food Sci Emerg Technol* 52:131-135.
- Chian FM, Kaur L, Oey I, Astruc T, Hodgkinson S, Boland M. 2019. Effect of pulsed electric fields (PEF) on the ultrastructure and *in vitro* protein digestibility of bovine *longissimus thoracis*. *LWT-Food Sci Technol* 103:253-259.
- Coria MS, Carranza PG, Palma GA. 2018. Calpain system in meat tenderization: A molecular approach. *Rev MVZ Córdoba* 23:6523-6536.
- Denoya GI, Vaudagna SR, Polenta G. 2015. Effect of high pressure processing and vacuum packaging on the preservation of fresh-cut peaches. *LWT-Food Sci Technol* 62:801-806.
- Dong M, Xu Y, Zhang Y, Han M, Wang P, Xu X, Zhou G. 2020. Physicochemical and structural properties of myofibrillar proteins isolated from pale, soft, exudative (pse)-like chicken breast meat: Effects of pulsed electric field (pef). *Innov Food Sci Emerg Technol* 59:102277.
- Faridnia F, Bekhit AEDA, Niven B, Oey I. 2014. Impact of pulsed electric fields and post-mortem vacuum ageing on beef *longissimus thoracis* muscles. *Int J Food Sci Technol* 49:2339-2347.
- Faridnia F, Bremer P, Burritt DJ, Oey I. 2016. Effects of pulsed electric fields on selected quality attributes of beef outside flat (*Biceps femoris*). 1st World Congress on Electroporation and Pulsed Electric Fields in Biology, Medicine and Food & Environmental Technologies, Portorož, Slovenia. pp 51-54.
- Faridnia F, Ma QL, Bremer PJ, Burritt DJ, Hamid N, Oey I. 2015. Effect of freezing as pre-treatment prior to pulsed electric field processing on quality traits of beef muscles. *Innov Food Sci Emerg Technol* 29:31-40.
- Farkas J, Polyák-Fehér K, Andrásy É, Mészáros L. 2002. Improvement of microbiological safety of sous-vide meals by

- gamma radiation. *Radiat Phys Chem* 63:345-348.
- Fauster T, Schlossnikl D, Rath F, Ostermeier R, Teufel F, Toepfl S, Jaeger H. 2018. Impact of pulsed electric field (PEF) pretreatment on process performance of industrial french fries production. *J Food Eng* 235:16-22.
- Gavahian M, Farahnaky A. 2018. Ohmic-assisted hydrodistillation technology: A review. *Trends Food Sci Technol* 72:153-161.
- Genovese J, Kranjc M, Serša I, Petracci M, Rocculi P, Miklavčič D, Mahnič-Kalamiza S. 2021. PEF-treated plant and animal tissues: Insights by approaching with different electroporation assessment methods. *Innov Food Sci Emerg Technol* 74:102872.
- Genovese J, Stručić M, Serša I, Novickij V, Rocculi P, Miklavčič D, Mahnič-Kalamiza S, Kranjc M. 2023. PEF treatment effect on plant tissues of heterogeneous structure no longer an enigma: MRI insights beyond the naked eye. *Food Chem* 405:134892.
- Houghton P, Lyng J, Cronin DA, Morgan DJ, Fanning S, Whyte P. 2012. Efficacy of pulsed electric fields for the inactivation of indicator microorganisms and foodborne pathogens in liquids and raw chicken. *Food Control* 25:131-135.
- Jeong SH, Jung YM, Kim S, Kim JH, Yeo H, Lee DU. 2023. Tenderization of beef *semitendinosus* muscle by pulsed electric field treatment with a direct contact chamber and its impact on proteolysis and physicochemical properties. *Foods* 12:430.
- Jeong SH, Kim EC, Lee DU. 2020. The impact of a consecutive process of pulsed electric field, sous-vide cooking, and reheating on the properties of beef *semitendinosus* muscle. *Foods* 9:1674.
- Kantono K, Hamid N, Ma Q, Oey I, Farouk M. 2021. Changes in the physicochemical properties of chilled and frozen-thawed lamb cuts subjected to pulsed electric field processing. *Food Res Int* 141:110092.
- Karki R, Oey I, Bremer P, Silcock P. 2023. Understanding the effect of meat electrical conductivity on pulsed electric field (PEF) process parameters and the ability of PEF to enhance the quality and shorten sous vide processing for beef short ribs. *Food Res Int* 163:112251.
- Khan AA, Randhawa MA, Carne A, Ahmed IAM, Barr D, Reid M, Bekhit AEDA. 2017. Effect of low and high pulsed electric field on the quality and nutritional minerals in cold boned beef *M. Longissimus et lumborum*. *Innov Food Sci Emerg Technol* 41:135-143.
- Kim YN, Kwon HJ, Lee DU. 2019. Effects of pulsed electric field (PEF) treatment on physicochemical properties of *Panax ginseng*. *Innov Food Sci Emerg Technol* 58:102232.
- Knorr D, Angersbach A. 1998. Impact of high-intensity electric field pulses on plant membrane permeabilization. *Trends Food Sci Technol* 9:185-191.
- Kumar Y, Patel KK, Kumar V. 2015. Pulsed electric field processing in food technology. *Int J Eng Stud Tech Approach* 1:6-17.
- Lebovka N, Bazhal MI, Vorobiev E. 2002. Estimation of characteristic damage time of food materials in pulsed-electric fields. *J Food Eng* 54:337-346.
- Lee SH, Shahbaz HM, Jeong SH, Hong SY, Lee DU. 2022. Effect of pulsed electric field treatment on cell-membrane permeabilization of potato tissue and the quality of French fries. *Ital J Food Sci* 34:13-24.
- Levy C, Aubert X, Lacour B, Carlin F. 2012. Relevant factors affecting microbial surface decontamination by pulsed light. *Int J Food Microbiol* 152:168-174.
- Li J, Shi J, Huang X, Zou X, Li Z, Zhang D, Zhang W, Xu Y. 2020. Effects of pulsed electric field on freeze-thaw quality of Atlantic salmon. *Innov Food Sci Emerg Technol* 65:102454.

- Marino R, Albenzio M, della Malva A, Santillo A, Loizzo P, Sevi A. 2013. Proteolytic pattern of myofibrillar protein and meat tenderness as affected by breed and aging time. *Meat Sci* 95:281-287.
- McDonnell CK, Allen P, Chardonnerau FS, Arimi JM, Lyng JG. 2014. The use of pulsed electric fields for accelerating the salting of pork. *LWT-Food Sci Technol* 59:1054-1060.
- Mok JH, Her JY, Kang T, Hoptowitz R, Jun S. 2017. Effects of pulsed electric field (PEF) and oscillating magnetic field (OMF) combination technology on the extension of supercooling for chicken breasts. *J Food Eng* 196:27-35.
- Mungure TE, Farouk MM, Birch EJ, Carne A, Staincliffe M, Stewart I, Bekhit AEDA. 2020. Effect of PEF treatment on meat quality attributes, ultrastructure and metabolite profiles of wet and dry aged venison *Longissimus dorsi* muscle. *Innov Food Sci Emerg Technol* 65:102457.
- Naliyadhara N, Kumar A, Girisa S, Daimary UD, Hegde M, Kunnumakkara AB. 2022. Pulsed electric field (PEF): Avant-garde extraction escalation technology in food industry. *Trends Food Sci Technol* 122:238-255.
- Nath KG, Pandiselvam R, Sunil CK. 2023. High-pressure processing: Effect on textural properties of food: A review. *J Food Eng* 351:111521.
- O'Dowd LP, Arimi JM, Noci F, Cronin DA, Lyng JG. 2013. An assessment of the effect of pulsed electrical fields on tenderness and selected quality attributes of post rigour beef muscle. *Meat Sci* 93:303-309.
- Perera N, Gamage TV, Wakeling L, Gamlath G, Versteeg C. 2010. Colour and texture of apples high pressure processed in pineapple juice. *Innov Food Sci Emerg Technol* 11:39-46.
- Sampedro F, McAloon A, Yee W, Fan X, Geveke DJ. 2014. Cost analysis and environmental impact of pulsed electric fields and high pressure processing in comparison with thermal pasteurization. *Food Bioprocess Technol* 7:1928-1937.
- Soro AB, Shokri S, Nicolau-Lapeña I, Ekhlash D, Burgess CM, Whyte P, Bolton DJ, Bourke P, Tiwari BK. 2023. Current challenges in the application of the UV-LED technology for food decontamination. *Trends Food Sci Technol* 131:264-276.
- Sruthi N, Josna K, Pandiselvam R, Kothakota A, Gavahian M, Khaneghah AM. 2022. Impacts of cold plasma treatment on physicochemical, functional, bioactive, textural, and sensory attributes of food: A comprehensive review. *Food Chem* 368:130809.
- Stachelska MA, Stankiewicz-Szymczak W, Jakubczak A, Świsłocka R, Lewandowski W. 2012. Influence of pulsed electric field on the survival of *Yersinia enterocolitica* in minced beef meat: Assessment of microbiological activity of selected cell lines of bacteria under influence of physical-chemical factors. *Apar Badaw Dydakt* 17:13-17.
- Suwandy V, Carne A, van de Ven R, Bekhit AEDA, Hopkins DL. 2015a. Effect of pulsed electric field on the proteolysis of cold boned beef *M. Longissimus lumborum* and *M. Semimembranosus*. *Meat Sci* 100:222-226.
- Suwandy V, Carne A, van de Ven R, Bekhit AEDA, Hopkins DL. 2015b. Effect of pulsed electric field treatment on hot-boned muscles of different potential tenderness. *Meat Sci* 105:25-31.
- Suwandy V, Carne A, van de Ven R, Bekhit AEDA, Hopkins DL. 2015c. Effect of repeated pulsed electric field treatment on the quality of cold-boned beef loins and topsides. *Food Bioprocess Technol* 8:1218-1228.
- Toepfl S, Siemer C, Saldaña-Navarro G, Heinz V. 2014. Overview of pulsed electric fields processing for food. In *Emerging technologies for food processing*. 2nd ed. Sun DW (ed). Academic Press, Cambridge, MA, USA. pp 93-114.
- Tornberg E. 2005. Effects of heat on meat proteins: Implications on structure and quality of meat products. *Meat Sci* 70:493-508.
- Wang Q, Wei R, Hu J, Luan Y, Liu R, Ge Q, Yu H, Wu M. 2022. Moderate pulsed electric field-induced structural unfolding

- ameliorated the gelling properties of porcine muscle myofibrillar protein. *Innov Food Sci Emerg Technol* 81:103145.
- Yates LD, Dutson TR, Caldwell J, Carpenter ZL. 1983. Effect of temperature and pH on the post-mortem degradation of myofibrillar proteins. *Meat Sci* 9:157-179.
- Zare F, Ghasemi N, Bansal N, Hosano H. 2023. Advances in pulsed electric stimuli as a physical method for treating liquid foods. *Phys Life Rev* 44:207-266.
- Zhang C, Lyu X, Arshad RN, Aadil RM, Tong Y, Zhao W, Yang R. 2023. Pulsed electric field as a promising technology for solid foods processing: A review. *Food Chem* 403:134367.
- Zhang Q, Barbosa-Cánovas GV, Swanson BG. 1995. Engineering aspects of pulsed electric field pasteurization. *J Food Eng* 25:261-281.
- Zimmermann U. 1996. The effect of high intensity electric field pulses on eukaryotic cell membranes: Fundamentals and applications. In *Electromanipulation of cells*. Zimmermann U, Neil GA (ed). CRC Press, Boca Raton, FL, USA. pp 1-106.