

ARTICLE

Comparison of Muscle Fiber and Meat Quality Characteristics of Beef Strip Loin, Tenderloin, and Round Cuts among Jeju Black Cattle, Hanwoo, and Their Crossbreeds

 Soo-Hyun Cho¹, Van-Ba Hoa¹, Dong-Heon Song¹, Dong Kyun Kim¹, Yun-Seok Kim¹, Hyun-Wook Kim¹, In-Seon Bae¹, Pil Nam Sung¹, Junyoung Park², Sumin Song², Huilin Cheng², Lixin Du², Choeun Im³, and Gap-Don Kim^{2,4,*}
 OPEN ACCESS

Received July 5, 2024
Revised July 27, 2024
Accepted July 29, 2024

*Corresponding author : Gap-Don Kim
 Graduate School of International Agricultural Technology, Seoul National University, Pyeongchang 25354, Korea
 Tel: +82-33-339-5778
 Fax: +82-33-339-5779
 E-mail: gapdonkim@snu.ac.kr

***ORCID**

Soo-Hyun Cho
<https://orcid.org/0000-0002-8073-8771>
 Van-Ba Hoa
<https://orcid.org/0000-0001-8725-1504>
 Dong-Heon Song
<https://orcid.org/0000-0002-4670-3321>
 Dong Kyun Kim
<https://orcid.org/0000-0002-8416-0436>
 Yun-Seok Kim
<https://orcid.org/0000-0003-4186-7877>
 Hyun-Wook Kim
<https://orcid.org/0000-0002-2979-345X>
 In-Seon Bae
<https://orcid.org/0000-0003-3543-8785>
 Pil Nam Sung
<https://orcid.org/0000-0003-2915-1059>
 Junyoung Park
<https://orcid.org/0000-0003-2569-6422>
 Sumin Song
<https://orcid.org/0000-0001-7115-2253>
 Huilin Cheng
<https://orcid.org/0000-0003-0628-3358>
 Lixin Du
<https://orcid.org/0000-0002-3287-5018>
 Choeun Im
<https://orcid.org/0000-0003-3564-7069>
 Gap-Don Kim
<https://orcid.org/0000-0001-5870-8990>

¹Animal Products Utilization Division, National Institute of Animal Science, RDA, Wanju 55365, Korea

²Graduate School of International Agricultural Technology, Seoul National University, Pyeongchang 25354, Korea

³Agricultural Cooperative Federation Livestock Research Institute, Seoul 04516, Korea

⁴Institutes of Green Bio Science & Technology, Seoul National University, Pyeongchang 25354, Korea

Abstract This study assessed and compared meat quality and fiber characteristics of *longissimus lumborum* (LL), *psaos major* (PM), and *semimembranosus* muscles among Hanwoo (HW), Jeju black (BL), and their crossbred (BH) cattle. Twelve carcasses from each breed (36 in total) were used in this study. BL and BH had higher moisture and crude ash contents and lower crude fat and protein contents than HW, regardless of the muscle type. BL had higher CIE a*, cooking loss, and shear force values than did the other breeds for all muscle types. The muscle fiber size (cross-sectional area) of BL and BH was larger than that of HW for all muscle types. Type IIX was the dominant muscle fiber type in both BL and BH, regardless of muscle type; however, HW had the highest composition of type I compared to the other types (IIA, IIAX, and IIX) in PM. Higher total fiber density was observed in the LL and PM muscles of HW than in those of BL and BH. Meat quality and muscle fiber characteristics of BL and BH were distinct from those of HW.

Keywords Jeju black cattle, Hanwoo, meat quality, muscle fiber characteristics

Introduction

Hanwoo (HW), a representative beef cattle in Republic of Korea, is known for its high marbling and palatability. The name ‘Hanwoo’ consists of Han (meaning Korea) and Woo (meaning cattle) and refers to Korean cattle. The coat color of HW is generally

brown. However, HW comprises four different breeds: Chickso (brindle with black and brown coat color), Heukwoo (coat color is black but occasionally mixed brown on the face and back), and Jeju black (BL; whole coat color is black; Lee et al., 2014). In the National Report on the State of Animal Genetic Resources from the Ministry for Food, Agriculture, Forestry and Fisheries (2004), cattle in the Republic of Korea comprised 67% brown cattle, 37% Holstein, and the rest included Chickso, Heukwoo, and BL. BL has been a natural monument 546 in the Republic of Korea after being formally registered with the FAO in this report. The population of BL was 1,087 in 2023, and the concerned body of Jeju Island has planned to increase this population to 2,000 by 2030. Overall, the BL breed still faces the threat of extinction, probably owing to the lack of consumer knowledge and need for this breed.

Studies on BL are scanty. Dadi et al. (2014) analyzed cattle genes distributed in East Asia and reported specific mitochondrial DNA sequences of BL, brown HW, and brindle HW. Alam et al. (2021) reported lower genetic heterozygosity in BL than in brown HW and Holstein in the Republic of Korea. Han et al. (2011) reported that the melanocortin-1 receptor alleles ED, E+, and e are found in BL and that these traits are directly related to the coat color phenotype of HW, BL, and their crossbreeds. Although BL exhibits lower average daily gain, live weight, and carcass weight, a higher feed efficiency has been observed than that in Charolles and the crossbreeds of BL and Charolles raised for 18 months (Oh et al., 2008). Lee et al. (2007) observed that pure BL shows a low average daily gain until 16 months, which increases during the period of 17–24 months, when compared with the those of other breeds. In the same study, although the carcass weight of pure BL was low, no significant difference in the carcass yield was observed. Moon et al. (2012) have compared same-quality grade brown HW, BL, and imported Australian beef and found no significant differences in the proximate composition; however, higher CIE a* and lower melting point were observed for BL than for the other two breeds. Additionally, the fatty acid composition of BL showed low levels of saturated and high levels of unsaturated fatty acids (Hoa et al., 2024; Lee et al., 2019; Moon et al., 2012; Song and Hwang, 2023; Tuell et al., 2022).

BL is a traditional cattle breed raised on Jeju Island in the Republic of Korea, and is served at national events such as the birthday of the King or memorial dates of the Royal Family (Lee et al., 2019). Although some studies have focused on the characteristics of BL, these were restricted to only genetics; additionally, meat quality has been evaluated only in loins. Therefore, this study aimed to evaluate the meat quality and fiber characteristics of *longissimus lumborum*, *psaos major*, and *semimembranosus* muscles from BL, brown HW, and their crossbreeds.

Materials and Methods

Sample preparation

A total of 36 steers (12 steers per breed) of HW (31.8±1.0 months of age, 620.3±6.5 kg carcass weight), BL (33.2±2.4 months of age, 583.3±17.9 kg carcass weight), and crossbreeds (BH; 31.3±1.4 months of age, 720.1±23.8 kg carcass weight), which were raised under the same conditions and fattened on a cattle farm located in Jeju Island, were randomly selected and transported to the abattoir of the National Institute of Animal Science (NIAS), Wanju, Republic of Korea. Feeding was conducted according to the Korean Feeding Standard for Hanwoo (NIAS, 2020). Livestock transport, preslaughter management, and slaughter were conducted in accordance with the Livestock Products Sanitation Control Act. Beef carcasses were chilled at 1°C for 48 h after slaughter; then, three muscles (*M. longissimus lumborum*, *M. psaos major*, and *M. semimembranosus*) were removed from the left sides of the carcasses. For immunohistochemistry, muscle tissues (1.5×1.5×2.0 cm) were immediately frozen in 2-methylbutane chilled with liquid nitrogen and stored at –80°C until analysis. The

remaining muscle samples were used for analyzing physicochemical properties (proximate composition, pH, color, cooking loss, and shear force).

Physicochemical properties

Moisture, crude protein, and crude ash contents were analyzed according to the guidelines of the Association of Official Agricultural Chemists (AOAC, 2000), with some modifications. Briefly, moisture content was determined by drying 10 g sample for 24 h in a dry oven at 105°C and expressed as the percentage of total weight by calculating the difference in weight before and after drying. To determine the crude protein content, 1.0 g sample was digested with sulfuric acid and distilled ammonia into boric acid following the Kjeldahl method (AOAC, 2000). Borate ions were titrated with hydrochloric acid, and crude protein content was calculated from the total nitrogen content. Crude ash contents were calculated by measuring the weights of samples before and after burning the muscles at 200°C, 400°C, 600°C, and 800°C for 2 h after pre-drying at 100°C for 2 h; the ash content was expressed as a percentage. Crude fat content was measured as described by Folch et al. (1957). Briefly, approximately 5.0 g sample was homogenized in 30 mL extraction buffer (chloroform:methanol at a 2:1 v/v ratio) and filtered through a Whatman No. 1 filter paper (Merck, Darmstadt, Germany). The filtrate was mixed with 0.88% sodium chloride and incubated at room temperature for 2 h. Once the upper layer was removed using an evaporator, 10 mL of the lower layer was dried in pre-weighed aluminum dish to obtain crude fat. Crude fat content was expressed as the percentage of the sample. The proximate composition was obtained from triplicate measurements for each sample.

For measuring the pH, 3.0 g sample was homogenized with 27 mL deionized water, and the pH was measured using a pH meter (S220, Mettler Toledo, Greifensee, Switzerland) calibrated with standard buffer solutions (pH 4.01, 7.00, and 9.21). The pH values were determined from three replicates of each sample.

Meat color intensity on the sample surface was measured after exposure to air for 20 min at room temperature. A colorimeter (CR-400, Konica Minolta, Tokyo, Japan) was set up with a D65 light source, an 8° illumination angle, and an 8-mm measuring aperture after calibration with a standard plate (Y, 93.5; x, 0.3132; y, 0.3198). Color values for CIE L*, CIE a*, and CIE b* were represented according to the Commission Internationale de l'Éclairage System (CIE, 1978). Meat color intensity was measured in five different regions of each sample.

Approximately 25 g sample was cooked in a water bath (WB-22, Daihan Scientific, Wonju, Korea) at 75°C to measure cooking loss. Samples for which the internal temperature reached 70°C were cooled to room temperature, and their weights were measured. The difference in weight before and after cooking was monitored. Cooking loss was calculated as the percentage of initial weight from triplicate measurements.

Cooked samples were used for measuring the shear force. Three cores (1.0 cm diameter) of each sample were obtained by cutting parallel to the orientation of muscle fiber. Shear force (N/cm²) was measured using a texture analyzer (TA1, AMETEK, Largo, FL, USA) equipped with a Warner-Bratzler shear blade at 3.0 mm/s and 50 kgf of load cell.

Muscle fiber characteristics

Immunohistochemistry was conducted to analyze muscle fiber characteristics according to the method described by Song et al. (2020), with some modifications. Briefly, cross-sections (10 µm thickness) obtained from each frozen sample were blocked with 10% normal goat serum (Cell Signalling Technology, Danvers, MA, USA) and incubated with primary antibodies (BA-F8, SC-71, and 6H1; DSHB, Iowa City, IA, USA). The cross-sections were then incubated with secondary antibodies (anti-IgG and anti-IgM) conjugated with Alexa Fluor 405, 488, or 594 (Thermo Fisher Scientific, Waltham, MA,

USA). Images of three different regions of each section were captured using a microscope (EVOS M5000, Thermo Fisher Scientific). Muscle fiber characteristics (cross-sectional area, μm^2 ; relative fiber area, %; fiber density, number/ mm^2) were evaluated using approximately 800 muscle fibers of each section. Image Pro Plus (Media Cybernetics, Rockville, MD, USA) was used to analyze each image.

Statistical analysis

All data are presented as mean \pm SE. Statistical analyses were conducted using SAS software v.9.4 (SAS Institute, Cary, NC, USA). Differences in physicochemical and muscle fiber characteristics within the same muscle among breeds were evaluated using a one-way analysis of variance (ANOVA), followed by Duncan's multiple range post-hoc test. Pearson's correlation coefficients between physicochemical properties and muscle fiber characteristics were analyzed to describe the relationship between muscle fiber characteristics and physicochemical properties, regardless of muscle type. Statistical significance was set at $p < 0.05$.

Results

The proximate composition and meat quality traits of BL, brown HW, and BH are shown in Figs. 1, 2, and 3. All characteristics, except the pH, were significantly different in *M. longissimus lumborum*. BL had relatively high moisture content and low crude fat and protein contents ($p < 0.05$; Figs. 1A, B, and C). BH showed intermediate traits, in which the moisture content was higher than that of HW, and a moderate level of crude fat content between those of HW and BL ($p < 0.05$). The crude ash content of BH was higher than that of HW ($p < 0.05$; Fig. 1D), but no significant difference was




Fig. 1. Comparison of proximate composition (A–D) and meat quality characteristics (E–J) of *M. longissimus lumborum* among Hanwoo (HW), Jeju black (BL), and their crossbred (BH) cattle. Data represent mean \pm SE (bar). ^{A–C} Significant differences ($p < 0.05$) among the breeds are represented with different letters.




Fig. 2. Comparison of proximate composition (A–D) and meat quality characteristics (E–J) of *M. psoas major* among Hanwoo (HW), Jeju black (BL), and their crossbred (BH) cattle. Data represent mean ± SE. ^{A–C} Significant differences (p < 0.05) among the breeds are represented with different letters.




Fig. 3. Comparison of proximate composition (A–D) and meat quality characteristics (E–J) of *M. semimembranosus* between Hanwoo (HW), Jeju black (BL), and their crossbred (BH) cattle. Data represent mean ± SE. ^{A–C} Significant differences (p < 0.05) among the breeds are represented with different letters.

observed when compared with that of BL ($p>0.05$). HW showed higher CIE L*, CIE a*, and CIE b* than did BH ($p<0.05$; Figs. 1F, G, and H). BL showed lower CIE L* than did HW and higher CIE a* and CIE b* than did BH ($p<0.05$). The cooking loss in BL was significantly lower than in HW and higher than in BH ($p<0.05$; Fig. 1I). The shear force was highest for BL, followed by that for BH and HW ($p<0.05$; Fig. 1J).

Most traits, except crude protein content and CIE L*, showed significant differences in *M. psoas major* among breeds. BL had higher moisture and crude ash contents and a lower crude fat content than those of HW ($p<0.05$; Figs. 2A, B, and D). The proximate composition of the *psoas major* muscle of BH was similar to that of BL, but the crude ash content was relatively higher in BL ($p<0.05$; Figs. 2A, B, and D). The pH value for BH was significantly lower than the values for HW and BL ($p<0.05$; Fig. 2E). BL showed greater CIE a* than did HW and BH; CIE b* was significantly different between BL and HW ($p<0.05$; Figs. 2G and H). Cooking loss in BH was significantly lower than that in BL and HW ($p<0.05$; Fig. 2I). The highest and lowest shear force values were observed in BL and HW, respectively ($p<0.05$; Fig. 2J).

Significant differences in all meat quality traits, except the pH, were observed in *M. semimembranosus* ($p<0.05$) among different breeds. The results of proximate composition showed significantly high moisture and crude ash contents in BH, and crude fat and crude protein content in HW ($p<0.05$; Figs. 3A, B, C, and D). The color of *M. semimembranosus* was similar to that of *M. longissimus lumborum*; HW showed higher CIE L* and lower CIE a* and CIE b* than did BL and BH ($p<0.05$; Figs. 3F, G, and H). BH showed lower cooking loss than did BL and HW ($p<0.05$; Fig. 3I), and the values for BL and HW were not significantly different ($p>0.05$). Significantly higher shear force was observed for BL than for HW ($p<0.05$; Fig. 3J), and no significant difference was observed in BH ($p>0.05$).

Representative immunofluorescent images of BL, HW, and BH for the analysis of muscle fiber characteristics are shown in Fig. 4. The cross-sectional area, relative area ratio, and density of muscle fibers were analyzed using those images; the results




Fig. 4. Immunohistochemical analysis of cross-sections of *M. longissimus lumborum*, *M. psoas major*, and *M. semimembranosus* of Hanwoo (HW), Jeju black (BL), and their crossbred (BH) cattle. Bar=300 μ m.

are shown in Fig. 5. Regarding the cross-sectional area of *M. longissimus lumborum*, the area was significantly higher in BL than in the other breeds, regardless of the muscle fiber type ($p < 0.05$; Fig. 4A). Specifically, muscle fiber types IIA and IIAX were not significantly different between BL and BH ($p > 0.05$); however, fiber types I and IIX and mean cross-sectional area were significantly higher in BL than in BH ($p < 0.05$). Type IIX fibers appeared to be largest in BL and BH, whereas type IIAX was largest in HW ($p < 0.05$). No significant difference was observed for type IIX ($p > 0.05$; Fig. 5A); BL had a higher area ratio of type I, and a lower area ratio of type IIA and IIAX than the other breeds ($p < 0.05$; Fig. 5B). Regardless of breed, the area ratio of IIX was highest and that of IIAX was lowest ($p < 0.05$). No significant difference was observed in type I among the breeds ($p > 0.05$; Fig. 5C). Relatively high fiber density was noticed for the other and total fiber types for *M. longissimus lumborum* in HW, whereas the density of type IIA fiber and total density in BL was lower than those in BH ($p < 0.05$).

The fiber characteristics of *M. psoas major* of the three different breeds are shown in Figs. 5D, E, and F. The cross-sectional area of BH was largest among all types, except for type IIAX and the mean area ($p < 0.05$). BL had significantly larger areas of types IIA, IIAX, and IIX than did HW ($p < 0.05$). Within breeds, BL and BH had the largest areas of type IIX, whereas type IIAX was largest in HW ($p < 0.05$). Regarding the relative area ratio, the highest ratio over 50% for type I was found in HW, and the area ratio of IIA was significantly high in HW ($p < 0.05$). No significant difference in type IIAX among the breeds ($p > 0.05$) was observed, but BL and HW showed a significantly high area ratio of type IIX ($p < 0.05$). HW had the




Fig. 5. Comparison of cross-sectional area (A, D, and G), relative fiber area (B, E, and H), and fiber density (C, F, and I) of *M. longissimus lumborum* (A, B, and C), *M. psoas major* (D, E, and F) and *M. semimembranosus* (G, H, and I) among Hanwoo (HW), Jeju black (BL), and their crossbred (BH) cattle. Data represent mean \pm SE. Significant differences ($p < 0.05$) among the breeds (X, Y, and Z) within the same muscle fiber type and between muscle fiber types (a–d) within the same muscle and breed are represented with different letters.

highest area ratio of type I, followed by that of IIA, IIX, and IIAX; BL and BH showed the highest area ratio of IIX, followed by that of I, IIA, and IIAX ($p < 0.05$). The highest densities of type I and IIA, and the highest total density were noticed in HW; type IIX in BL was significantly higher than that in the other breeds ($p < 0.05$). No significant difference was noticed for type IIAX among the different breeds ($p > 0.05$). Within the breeds, the density of type I was highest across all breeds, followed by that of IIX, IIA, and IIAX in BL and BH, and IIA, IIX, and IIAX in HW ($p < 0.05$).

The muscle fiber characteristics of *M. semimembranosus* are shown in Figs. 5G, H, and I. Unlike that in other muscles, a significant difference in cross-sectional area was observed only for type IIX fiber ($p < 0.05$). Within the breeds, muscle fiber type IIX was largest in BL and BH, and the cross-sectional area of type IIAX was higher than that of type IIA in HW ($p < 0.05$). The relative area ratio of *M. semimembranosus* was significantly higher for types I and IIA in HW and BL than in BH and that for IIX was higher in BH than in the other breeds ($p < 0.05$). A comparison of muscle fiber types within the same breed revealed high-to-low ratios in the order of type IIX, IIA, I, and IIAX, regardless of breed ($p < 0.05$). Significant differences in fiber density were noticed for types IIA, IIX, and total density; the density was highest in HW ($p < 0.05$). However, no significant differences were observed in type IIA and total density compared with those in BL ($p > 0.05$). Among the types within the same breed, all breeds showed the same order of trend: types IIA, IIX, I, and IIAX ($p < 0.05$).

The correlation coefficients between the muscle fiber characteristics and physicochemical properties are shown in Table 1. The cross-sectional area of each fiber was not significantly correlated with moisture and crude protein contents ($p > 0.05$), while that of type IIX showed negative and positive correlations with crude fat and crude ash contents, respectively ($p < 0.05$). CIE a* showed a correlation with the cross-sectional area, except in type IIAX fiber ($p < 0.05$), while CIE L* and CIE b* exhibited no significant correlations ($p > 0.05$). The cross-sectional area of type IIX showed a significant correlation with cooking loss ($p < 0.05$). Shear force exhibited a positive correlation with each muscle fiber type ($p < 0.05$). Relative fiber area of type I exhibited negative correlations with moisture, crude protein, crude ash contents, and shear force, while it showed a positive correlation with crude fat content ($p < 0.05$). The relative area of type IIA was positively correlated with crude protein content, CIE b*, cooking loss, and shear force ($p < 0.05$). Relative fiber area of type IIAX exhibited a negative correlation with moisture contents; however, it was positively correlated with crude fat content and CIE L* ($p < 0.05$). The relative area of type IIX exhibited positive correlations with moisture, crude ash, and shear force but a negative correlation with crude fat content

Table 1. Correlation coefficients between physicochemical properties and muscle fiber characteristics

Traits	Cross-sectional area					Relative fiber area				Fiber density				
	I	IIA	IIAX	IIX	Mean	I	IIA	IIAX	IIX	I	IIA	IIAX	IIX	Total
Moisture content	0.06	0.00	0.01	0.16	0.09	-0.39***	0.02	-0.41***	0.46***	-0.30**	-0.17	-0.41***	0.13	-0.21*
Crude protein content	0.15	0.12	0.10	-0.01	0.08	-0.32***	0.36***	0.06	0.09	-0.23**	-0.01	0.00	0.08	-0.12
Crude fat content	-0.15	-0.07	-0.06	-0.22*	-0.17	0.49***	-0.11	0.33***	-0.49***	0.39***	0.21*	0.36	-0.10	0.28**
Crude ash content	0.02	0.11	0.07	0.26**	0.16	-0.29**	-0.15	-0.05	0.38***	-0.24**	-0.28**	-0.10	-0.09	-0.24**
pH	0.01	-0.03	-0.04	-0.05	-0.03	0.15	-0.10	-0.06	-0.06	0.10	0.03	0.02	0.04	0.08
CIE L*	0.05	0.01	0.06	-0.11	-0.03	-0.15	0.19	0.23**	-0.01	-0.11	0.02	0.05	0.13	-0.03
CIE a*	-0.21*	-0.20*	-0.15	-0.19*	-0.21	-0.09	0.01	-0.03	0.09	0.01	0.10	0.04	0.27**	0.10
CIE b*	-0.06	-0.05	-0.05	-0.16	-0.10	-0.13	0.20*	0.05	0.00	-0.08	0.08	0.01	0.19*	0.02
Cooking loss	0.00	-0.17	-0.11	-0.23**	-0.15	-0.15	0.33***	-0.05	-0.04	-0.04	0.21*	0.01	0.18	0.08
Shear force	0.74*	0.65*	0.56*	0.69***	0.73	-0.52***	0.36***	-0.10	0.34***	-0.69***	-0.55***	-0.50***	-0.46***	-0.70***

* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

($p < 0.05$). Correlation of type I fiber density with physicochemical properties corresponded to that of relative type I fiber area ($p < 0.05$). Proximate composition showed significant correlations with types IIA, IIX, and total fiber density ($p < 0.05$). CIE a^* , CIE b^* , and cooking loss showed positive correlations with types IIX and IIA ($p < 0.05$). Shear force showed a negative correlation with all muscle fiber density, including total fiber density ($p < 0.05$). Among the physicochemical properties, only the pH did not show any significant correlation with muscle fiber characteristics ($p > 0.05$).

Discussion

In the present study, BL and BH had relatively high moisture contents and relatively low crude fat and protein contents in the three different muscles. In contrast, HW had a relatively low moisture content and relatively high crude fat and protein contents. BH had a relatively high crude ash content, regardless of the muscle type. The quality grade of beef is highly dependent on the intramuscular fat contents owing to the preference for grilling beef as thin slices in the Republic of Korea; the general intramuscular fat content of HW is 6.60%–23.85%. In the present study, the quality grade of HW seemed to be 1, and that of BH and BL between 1–2 (Gajaweera et al., 2020; Jo et al., 2012; Lee et al., 2015). The intramuscular fat content affects eating quality, particularly tenderness and flavor (Gotoh et al., 2018; Wheeler et al., 1994). The factors affecting the intramuscular fat contents of meat vary; nevertheless, consumer demands can be met by increasing the intramuscular fat content through breeding or feeding systems, and likewise, HW was originally used for farming to meet such demands (Avilés et al., 2015; Lee et al., 2014; Miller et al., 1987; Park et al., 2018). The pH significantly affects muscle fiber characteristics, meat color, and water-holding capacity (Ertbjerg and Puolanne, 2017; Hopkins et al., 2014; Ryu and Kim, 2006; Węglarz, 2010); however, it did not significantly affect other meat quality traits in the present study. The results of meat color showed higher CIE L^* , CIE a^* , and CIE b^* in the *M. longissimus lumborum* and *M. semimembranosus* of HW than in those of others. Wulf and Wise (1999) reported that color positively correlates with lean maturity and marbling score; and another study has reported significantly low CIE a^* of *M. longissimus thoracis* with low intramuscular fat in HW (Kim and Lee, 2003). However, a contrasting result for *M. psoas major* indicated that intramuscular fat did not affect meat color in the present study. Cho et al. (2010) and Jo et al. (2012) reported the lowest CIE a^* in quality grade 1 in HW, which increased following an increase in the intramuscular fat content. Furthermore, Lee et al. (2012) have reported a significant difference in CIE a^* in the intramuscular fat, ranging between 13.12%–18.69 %, as analyzed from the HW marbling score. *M. psoas major* used in the present study had higher intramuscular fat and lower moisture contents. These findings indicate that intramuscular fat affects meat color, and the extent depends on the fat content. In all the three muscles, cooking loss was higher in BH than in the other breeds, which is in line the results of a previous study that analyzed proximate composition by doneness, wherein the degree of change in crude fat and moisture contents was higher than that of crude protein and ash contents (Smith et al., 2011). Regardless of the muscle types, BL showed the highest and HW showed the lowest shear force values, which is in agreement with the findings of Wheeler et al. (1994), who reported that an increase in intramuscular fat content results in a reduction in the shear force value. However, shear force does not affect all traits of meat, and additional analyses, such as texture profile analysis or sensory evaluation, are necessary (Caine et al., 2003).

The aforementioned meat quality traits are dependent on individual trait and muscle fiber characteristics. The trends such as large cross-sectional area of glycolytic muscle fibers, small oxidative fibers, and intermediate fibers are intermediate have been reported previously (Maltin et al., 1998; Spindler et al., 1980; Xie et al., 2012). Regarding significant differences in the cross-sectional area, those of HW were smaller than those of the other breeds. No significant difference was found in the

cross-sectional area of *M. semimembranosus* between BL and BH, and *M. longissimus lumborum* in BL and *M. psoas major* in BH had larger cross-sectional areas than those of the other breeds. Although the carcass weight of BL was lowest, *M. longissimus lumborum* in BL and *M. psoas major* in BH exhibited largest cross-sectional area than those of other breeds, while the cross-sectional area of *M. semimembranosus* was not significantly different between BL and BH. These results are in accordance with previous findings showing a positive correlation with intramuscular fat and a negative correlation with CIE L* of the cross-sectional area of types I and IIA (Wegner et al., 2000). However, no significant correlation was observed in the present study. Similarly, a previous study has reported that relatively small fiber size and increased number of oxidative fibers promote metabolism, resulting in increased fat accumulation owing to energy supplementation, thereby affecting meat quality (Schiaffino and Reggiani, 1996). Although no significant correlation was observed between crude fat content and oxidative fibers, type IIX showed a negative correlation with crude fat content. Probably the increasing size resulted in increased shear force (Joo et al., 2017), while the diameter did not (Hwang et al., 2010; Ozawa et al., 2000).

Generally, the muscle fiber density of HW was higher, while the cross-sectional area was similar to or lower than those of the other breeds in the present study. This implies that the muscle fiber number in BL and BH is less than that in HW, and hyperplasia of HW or hypertrophy of BL and BH can be considered, indicating that the characteristics of each cell from the three breeds are different (Picard and Gagaoua, 2020). While only a few studies have compared muscle fiber characteristics of HW with those of other breeds, Kim et al. (2021) have reported similar results, showing that HW consists of an increased number of muscle fibers compared to that in Holstein raised for meat in the Republic of Korea; additionally, a relatively low shear force has been observed. Xie et al. (2012) have reported different muscle fiber characteristics among five different breeds raised in the same feeding system. Johnston et al. (1975) and Spindler et al. (1980) have observed different muscle fiber characteristics among breeds. Generally, the cross-sectional area of HW is smaller than that of the other breeds, whereas the cross-sectional areas of BL and BH are larger than the area of HW.

Muscle fiber composition, representing muscle characteristics, is affected by breed, similar to that in other species (Park et al., 2022; Picard and Gagaoua, 2020). The present study reported that each muscle has a different muscle-fiber composition. Unexpectedly, the compositions of types I and IIX did not vary much from those of other muscles and were found to be similar to those of *M. psoas major* of BL and BH. Some differences were observed between the muscles in terms of muscle fiber composition. HW showed results similar to those of previous studies (Hwang et al., 2010; Kim et al., 2016). Type IIX fibers are positively correlated with the shear force. A relatively low shear force value was observed for HW, which had a relatively low type IIX composition in the present study. A high number of glycolytic fibers results in a reduction in the pH via accumulation of lactic acid during glycolysis in the early postmortem period; however, this does not fully harmonize with the results of the present study (Yu et al., 2019). The high content of oxidative fibers (types I and IIA) results in an increase in CIE a* owing to their metabolic traits (Jeong et al., 2009; Lefaucheur, 2010); however, the present study did not show similar results, which may be due to the low fat content. Muscle fiber composition exhibited changes in proximate composition in terms of fat accumulation, as previously described by Schiaffino and Reggiani (1996). Moreover, a high ratio of type IIX fiber resulted in a high shear force, as also previously described by Hwang et al. (2010). However, whether fat contents, type IIX characteristics, or other factors cause this phenomenon should be clarified.

Conclusion

The meat quality traits and muscle fiber characteristics of three different muscles (*M. longissimus lumborum*, *M. psoas*

major, and *M. semimembranosus*) from three different breeds (BL, HW, and BH) were evaluated. BL showed lower intramuscular fat content and higher CIE L*, CIE a*, CIE b*, cooking loss, and shear force than the other breeds. The BH cattle had intermediate characteristics between those of BL and HW cattle. HW had smaller muscle fiber size and higher muscle fiber density than did the other breeds. In conclusion, differences in physicochemical and muscle fiber characteristics were confirmed, regardless of the type of muscle, among three breeds of Korean native cattle.

Conflicts of Interest

The authors declare no potential conflicts of interest.

Acknowledgements

This research work was carried out with the support of the “Cooperative Research Program for Agriculture Science and Technology Development (Project No. PJ01620103),” Rural Development Administration, Korea.

Author Contributions

Conceptualization: Cho SH, Kim GD. Data curation: Hoa VB, Song DH, Kim DK, Kim YS, Kim HW, Bae IS, Sung PN, Cheong H, Du L, Im C. Formal analysis: Park J, Song S. Methodology: Cho SH, Song DH, Park J, Kim GD. Writing - original draft: Cho SH, Park J, Song S, Kim GD. Writing - review & editing: Cho SH, Hoa VB, Song DH, Kim DK, Kim YS, Kim HW, Bae IS, Sung PN, Park J, Song S, Cheng H, Du L, Im C, Kim GD.

Ethics Approval

This article does not require IRB/IACUC approval because there are no human and animal participants.

References

- Alam MZ, Lee YM, Son HJ, Hanna LH, Riley DG, Mannen H, Sasazaki S, Park SP, Kim JJ. 2021. Genetic characteristics of Korean Jeju black cattle with high density single nucleotide polymorphisms. *Anim Biosci* 34:789-800.
- AOAC. 2000. Official methods of analysis. 18th ed. Association of Official Analytical Chemists, Washington, DC, USA.
- Avilés C, Martínez AL, Domenech V, Peña F. 2015. Effect of feeding system and breed on growth performance, and carcass and meat quality traits in two continental beef breeds. *Meat Sci* 107:94-103.
- Caine WR, Aalhus JL, Best DR, Dugan MER, Jeremiah LE. 2003. Relationship of texture profile analysis and Warner-Bratzler shear force with sensory characteristics of beef rib steaks. *Meat Sci* 64:333-339.
- Cho SH, Kim J, Park BY, Seong PN, Kang GH, Kim JH, Jung SG, Im SK, Kim DH. 2010. Assessment of meat quality properties and development of a palatability prediction model for Korean Hanwoo steer beef. *Meat Sci* 86:236-242.
- Commission Internationale de l'Eclairage [CIE]. 1978. Recommendations on uniform color spaces-color difference equations, psychometric color terms. Supplement No. 2 to CIE Publication No. 15 (E.-1.3. 1). Commission Internationale de l'Eclairage, Paris, France.

- Dadi H, Lee SH, Lee SS, Park C, Kim KS. 2014. Inter- and intra-population genetic divergence of East Asian cattle populations: Focusing on Korean cattle. *Genes Genom* 36:261-265.
- Ertbjerg P, Puolanne E. 2017. Muscle structure, sarcomere length and influences on meat quality: A review. *Meat Sci* 132:139-152.
- Folch J, Lees M, Sloane-Stanley GH. 1957. A simple method for the isolation and purification of total lipids from animal tissues. *J Biol Chem* 226:497-509.
- Gajaweera C, Chung KY, Lee SH, Wijayananda HI, Kwon EG, Kim HJ, Cho SH, Lee SH. 2020. Assessment of carcass and meat quality of *longissimus thoracis* and *semimembranosus* muscles of Hanwoo with Korean beef grading standards. *Meat Sci* 160:107944.
- Gotoh T, Nishimura T, Kuchida K, Mannen H. 2018. The Japanese Wagyu beef industry: Current situation and future prospects: A review. *Asian-Australas J Anim Sci* 31:933-950.
- Han SH, Cho IC, Kim JH, Ko MS, Kim YH, Kim EY, Park SP, Lee SS. 2011. Coat color patterns and genotypes of extension and agouti in Hanwoo and Jeju black cattle. *J Life Sci* 21:494-501.
- Hoa VB, Kim DG, Song DH, Ko JH, Kim HW, Bae IS, Kim YS, Cho SH. 2024. Quality properties and flavor-related components of beef *longissimus lumborum* muscle from four Korean native cattle breeds. *Food Sci Anim Resour* 44:832-848.
- Hopkins DL, Ponnampalam EN, van de Ven RJ, Warner RD. 2014. The effect of pH decline rate on the meat and eating quality of beef carcasses. *Anim Prod Sci* 54:407-413.
- Hwang YH, Kim GD, Jeong JY, Hur SJ, Joo ST. 2010. The relationship between muscle fiber characteristics and meat quality traits of highly marbled Hanwoo (Korean native cattle) steers. *Meat Sci* 86:456-461.
- Jeong JY, Hur SJ, Yang HS, Moon SH, Hwang YH, Park GB, Joo ST. 2009. Discoloration characteristics of 3 major muscles from cattle during cold storage. *J Food Sci* 74:C1-C5.
- Jo C, Cho SH, Chang J, Nam KC. 2012. Keys to production and processing of Hanwoo beef: A perspective of tradition and science. *Anim Front* 2:32-38.
- Johnston DM, Stewart DF, Mood WG, Boling J, Kemp JD. 1975. Effect of breed and time on feed on the size and distribution of beef muscle fiber types. *J Anim Sci* 40:613-620.
- Joo SH, Lee KW, Hwang YH, Joo ST. 2017. Histochemical characteristics in relation to meat quality traits of eight major muscles from Hanwoo steers. *Korean J Food Sci Anim Resour* 37:716-725.
- Kim CJ, Lee ES. 2003. Effects of quality grade on the chemical, physical and sensory characteristics of Hanwoo (Korean native cattle) beef. *Meat Sci* 63:397-405.
- Kim GD, Yang HS, Jeong JY. 2016. Comparison of characteristics of myosin heavy chain-based fiber and meat quality among four bovine skeletal muscles. *Korean J Food Sci Anim Resour* 36:819-828.
- Kim JY, Lee B, Kim DH, Lee K, Kim EJ, Choi YM. 2021. Sensory quality and histochemical characteristics of *longissimus thoracis* muscles between Hanwoo and Holstein steers from different quality grades. *Food Sci Anim Resour* 41:779-787.
- Lee C, Lee H, Baek J, Park J, Jung S, Jo C. 2015. Comparison of crude fat content and intramuscular fat score at different position of loin from Hanwoo. *Korean J Agric Sci* 42:47-51.
- Lee JM, Choe JH, Jin HJ, Kim TI, Park BY, Hwang DY, Koh KC, Kim CJ, Hwang KS. 2012. Effect of marbling score on carcass grade factors, physico-chemical and sensory traits of *M. longissimus dorsi* in Hanwoo. *Korean J Food Sci Anim Resour* 32:659-668.

- Lee SH, Kim CN, Ko KB, Park SP, Kim HK, Kim JM, Ryu YC. 2019. Comparisons of beef fatty acid and amino acid characteristics between Jeju black cattle, Hanwoo, and Wagyu breeds. *Food Sci Anim Resour* 39:402-409.
- Lee SH, Park BH, Sharma A, Dang CG, Lee SS, Choi TJ, Choy YH, Kim HC, Jeon KJ, Kim SD, Yeon SH, Park SB, Kang HS. 2014. Hanwoo cattle: Origin, domestication, breeding strategies and genomic selection. *J Anim Sci Technol* 56:2.
- Lee WS, Oh WY, Lee SS, Khan MA, Ko MS, Kim HS, Ha JK. 2007. Growth performance and carcass evaluation of Jeju native cattle and its crossbreds fed for long fattening period. *Asian-Australas J Anim Sci* 20:1909-1916.
- Lefaucheur L. 2010. A second look into fibre typing: Relation to meat quality. *Meat Sci* 84:257-270.
- Maltin CA, Sinclair KD, Warriss PD, Grant CM, Porter AD, Delday MI, Warkup CC. 1998. The effects of age at slaughter, genotype and finishing system on the biochemical properties, muscle fibre type characteristics and eating quality of bull beef from suckled calves. *Anim Sci* 66:341-348.
- Miller MF, Cross HR, Crouse JD. 1987. Effect of feeding regimen, breed and sex condition on carcass composition and feed efficiency. *Meat Sci* 20:39-50.
- Ministry for Food, Agriculture, Forestry and Fisheries. 2004. National report on the state of animal genetic resources. Ministry for Food, Agriculture, Forestry and Fisheries, Sejong, Korea.
- Moon YH. 2012. Comparison of quality characteristics among chilled loins obtained from Jeju black cattle, Hanwoo and imported Australian beef. *J East Asian Soc Diet Life* 22:497-505.
- National Institute of Animal Science [NIAS]. 2020. Korean feeding standard for Hanwoo. Rural Development Administration, Jeonju, Korea.
- Oh WY, Lee WS, Lee SS, Khan MA, Ko MS, Yang SH, Kim HS, Ha JK. 2008. Feed consumption, body weight gain and carcass characteristics of Jeju native cattle and its crossbreds fed for short fattening period. *Asian-Australas J Anim Sci* 21:1745-1752.
- Ozawa S, Mitsuhashi T, Mitsumoto M, Matsumoto S, Itoh N, Itagaki K, Kohno Y, Dohgo T. 2000. The characteristics of muscle fiber types of *longissimus thoracis* muscle and their influences on the quantity and quality of meat from Japanese black steers. *Meat Sci* 54:65-70.
- Park J, Song S, Cheng H, Im C, Jung EY, Moon SS, Choi J, Hur SJ, Joo ST, Kim GD. 2022. Comparison of meat quality and muscle fiber characteristics between porcine skeletal muscles with different architectures. *Food Sci Anim Resour* 42:874-888.
- Park SJ, Beak SH, Jung DJS, Kim SY, Jeong IH, Piao MY, Kang HJ, Fassah DM, Na SW, Yoo SP, Baik M. 2018. Genetic, management, and nutritional factors affecting intramuscular fat deposition in beef cattle: A review. *Asian-Australas J Anim Sci* 31:1043-1061.
- Picard B, Gagaoua M. 2020. Muscle fiber properties in cattle and their relationships with meat qualities: An overview. *J Agric Food Chem* 68:6021-6039.
- Ryu YC, Kim BC. 2006. Comparison of histochemical characteristics in various pork groups categorized by postmortem metabolic rate and pork quality. *J Anim Sci* 84:894-901.
- Schiaffino S, Reggiani C. 1996. Molecular diversity of myofibrillar proteins: Gene regulation and functional significance. *Physiol Rev* 76:371-423.
- Smith AM, Harris KB, Haneklaus AN, Savell JW. 2011. Proximate composition and energy content of beef steaks as influenced by USDA quality grade and degree of doneness. *Meat Sci* 89:228-232.
- Song S, Ahn CH, Kim GD. 2020. Muscle fiber typing in bovine and porcine skeletal muscles using immunofluorescence with

- monoclonal antibodies specific to myosin heavy chain isoforms. *Food Sci Anim Resour* 40:132-144.
- Song Z, Hwang I. 2023. Objective meat quality from quality grade and backfat thickness of Hanwoo steers. *Food Sci Anim Resour* 43:531-539.
- Spindler AA, Mathias MM, Cramer DA. 1980. Growth changes in bovine muscle fiber types as influenced by breed and sex. *J Food Sci* 45:29-31.
- Tuell JR, Nondorf MJ, Kim YHB. 2022. Post-harvest strategies to improve tenderness of underutilized mature beef: A review. *Food Sci Anim Resour* 42:723-743.
- Węglarz A. 2010. Meat quality defined based on pH and colour depending on cattle category and slaughter season. *Czech J Anim Sci* 55:548-556.
- Wegner J, Albrecht E, Fiedler I, Teuscher F, Papstein HJ, Ender K. 2000. Growth- and breed-related changes of muscle fiber characteristics in cattle. *J Anim Sci* 78:1485-1496.
- Wheeler TL, Cundiff LV, Koch RM. 1994. Effect of marbling degree on beef palatability in *Bos taurus* and *Bos indicus* cattle. *J Anim Sci* 72:3145-3151.
- Wulf DM, Wise JW. 1999. Measuring muscle color on beef carcasses using the L*a*b* color space. *J Anim Sci* 77:2418-2427.
- Xie X, Meng Q, Cui Z, Ren L. 2012. Effect of cattle breed on meat quality, muscle fiber characteristics, lipid oxidation and fatty acids in China. *Asian-Australas J Anim Sci* 25:824-831.
- Yu Q, Tian X, Shao L, Li X, Dai R. 2019. Targeted metabolomics to reveal muscle-specific energy metabolism between bovine *longissimus lumborum* and *psaos major* during early postmortem periods. *Meat Sci* 156:166-173.