

1  
2  
3  
4

**TITLE PAGE**  
**- Korean Journal for Food Science of Animal Resources -**  
**Upload this completed form to website with submission**

<b>ARTICLE INFORMATION</b>	<b>Fill in information in each box below</b>
<b>Article Type</b>	Research article
<b>Article Title</b>	Physicochemical analysis of yogurt produced by <i>Leuconostoc mesenteroides</i> H40 and its effects on oxidative stress in neuronal cells
<b>Running Title (within 10 words)</b>	Neuroprotective effects of <i>Leuconostoc mesenteroides</i> and its probiotic yogurt
<b>Author</b>	Na-Kyoung Lee, Sung-Min Lima, Min-Jeong Cheon, Hyun-Dong Paik
<b>Affiliation</b>	Department of Food Science and Biotechnology of Animal Resources, Konkuk University, Seoul 05029, Korea
<b>Special remarks – if authors have additional information to inform the editorial office</b>	Not applicable
<b>ORCID (All authors must have ORCID) <a href="https://orcid.org">https://orcid.org</a></b>	Na-Kyoung Lee ( <a href="https://orcid.org/0000-0002-2395-550X">https://orcid.org/0000-0002-2395-550X</a> ) Sung-Min Lim ( <a href="https://orcid.org/0000-0001-5384-3210">https://orcid.org/0000-0001-5384-3210</a> ) Min-Jeong Cheon ( <a href="https://orcid.org/0000-0003-4688-2750">https://orcid.org/0000-0003-4688-2750</a> ) Hyun-Dong Paik ( <a href="https://orcid.org/0000-0001-9891-7703">https://orcid.org/0000-0001-9891-7703</a> )
<b>Conflicts of interest</b> List any present or potential conflicts of interest for all authors. (This field may be published.)	The authors declare no potential conflict of interest.
<b>Acknowledgements</b> State funding sources (grants, funding sources, equipment, and supplies). Include name and number of grant if available. (This field may be published.)	Not applicable
<b>Author contributions</b> (This field may be published.)	Conceptualization: Lee NK, Lim SM, Paik HD. Data curation: Lee NK, Lim SM, Cheon MJ, Paik HD. Formal analysis: Lim SM, Cheon MJ Methodology: Lim SM, Cheon MJ Software: Lee NK, Lim SM. Validation: Lee NK, Lim SM, Paik HD. Investigation: Lee NK, Lim SM. Writing - original draft: Lee NK, Lim SM. Writing - review & editing: Lee NK, Lim SM, Cheon MJ, Paik HD.
<b>Ethics approval (IRB/IACUC)</b> (This field may be published.)	This manuscript does not require IRB/IACUC approval because there are no human and animal participants.

5  
6

**CORRESPONDING AUTHOR CONTACT INFORMATION**

<b>For the corresponding author (responsible for correspondence, proofreading, and reprints)</b>	<b>Fill in information in each box below</b>
First name, middle initial, last name	Hyun-Dong Paik
Email address – this is where your proofs will be sent	hdpaik@konkuk.ac.kr
Secondary Email address	
Postal address	Department of Food Science and Biotechnology of Animal Resources, Konkuk University, Seoul 05029, Korea
Cell phone number	+82-10-4586-6189
Office phone number	+82-2-2049-6011
Fax number	

7  
8

9 **Abstract**

10 *Leuconostoc mesenteroides* H40 (H40) was isolated from kimchi, and its probiotic  
11 properties and neuroprotective effect was evaluated in oxidatively stressed SH-SY5Y cells.  
12 H40 was stable in artificial gastric conditions and can be attached in HT-29 cells. In addition,  
13 H40 did not produce  $\beta$ -glucuronidase and showed resistant to several antibiotics. The  
14 conditioned medium (CM) was made using HT-29 cells refined with heat-killed probiotics  
15 (probiotic-CM) and heated yogurts (Y-CM) to investigate the neuroprotective effect.  
16 Treatment with H40-CM not only increased cell viability but also significantly improved  
17 brain derived neurotropic factor (*BDNF*) expression and reduced the *Bax/Bcl-2* ratio in  
18 oxidatively stress-induced SH-SY5Y cells. Besides, probiotic Y-CM significantly increased  
19 *BDNF* mRNA expression and decreased *Bax/Bcl-2* ratio. The physicochemical properties of  
20 probiotic yogurt with H40 was not significantly different from the control yogurt. The viable  
21 cell counts of lactic acid bacteria in control and probiotic yogurt with H40 was 8.66 Log  
22 CFU/mL and 8.96 Log CFU/mL, respectively. Therefore, these results indicate that H40 can  
23 be used as prophylactic functional dairy food having neuroprotective effects.

24  
25 **Keywords** probiotics, *Leuconostoc mesenteroides*, neuroprotective effect, probiotic yogurt,  
26 oxidative stress

27  
28

## 29 **Introduction**

30 Probiotics are living microbes that deliver health benefits to the host when ingested in  
31 adequate amounts by the FAO/WHO (Chamber et al., 2019). The common probiotic strains  
32 mainly belong to the *Lactobacillus*, *Leuconostoc*, *Pediococcus*, and *Bifidobacterium* species  
33 and are widely used in many probiotic products (O'Toole et al., 2017). Probiotics have  
34 many reported health benefits such as improvement of cognitive function (Ton et al., 2020),  
35 antioxidant (Jang et al., 2018; Yu et al., 2019b), anti-inflammatory (Yu et al., 2019c),  
36 antihypertensive (Klippel et al., 2016), or cholesterol lowering (Ishimwe et al., 2015)  
37 activities. To utilize this functionality, probiotics are also used as medical or food additives.

38 The brain, which is rich in phospholipids, is an organ with high oxygen demand and is  
39 vulnerable to the effects of reactive oxygen species (ROS) (Dussert et al., 2006). ROS is an  
40 essential byproduct of aerobic metabolism (Wang and Michaelis, 2010). However, excessive  
41 ROS levels cause cell damage by oxidizing cellular biomolecules, including nucleic acids,  
42 proteins, and lipids (Lobo et al., 2010). ROS can contribute to pathologies, such as cancer  
43 (Lee et al., 2014), cardiovascular disease (Elahi et al., 2009), diabetes, and aging (Pamplona  
44 and Barja, 2006).

45 The bidirectional signaling connecting the brain and the gastrointestinal tract is crucial  
46 for maintaining homeostasis and is regulated the neural, hormonal, and immunological levels  
47 (Ghaisas et al., 2016; Wang and Kasper, 2014). Probiotics have recently become a target as  
48 live bacterial cell biotherapies for neurodegenerative disease (Quigley, 2017; Wang et al.,  
49 2016). *Clostridium butyricum* can exert neuroprotective effects against ischemia/reperfusion  
50 injury mice through antioxidant and anti-apoptosis mechanisms (Sun et al., 2016).  
51 *Lactobacillus buchneri* KU200793 showed neuroprotective effect using SH-SY5Y cells  
52 induced with 1-methyl-4-phenylpyridinium (MPP<sup>+</sup>) (Cheon et al., 2020).

53 Brain derived neurotropic factor (BDNF) expression occurs in the brain, and low  
54 secretion of BDNF influences human memory and hippocampal functions (Egan et al., 2003).  
55 BDNF is mediated by extracellular signal-regulated kinase (ERK) 1/2, ERK5, and  
56 phosphatidylinositol-3 kinase (PI3k) pathways in cortical neurons to promote neuronal  
57 survival (Liu et al., 2003). Oxidative stress may induce mitochondrial dysfunction and  
58 deficiency in protein aggregation and ultimately cause nerve cell death (Lobo et al., 2010).  
59 The mitochondrial apoptotic pathways are mediated through the Bcl-2 family proteins, which  
60 include Bax that promotes pro-apoptotic mitochondrial permeability and anti-apoptotic Bcl-2  
61 that inhibits apoptotic effects (Azmi et al., 2013). The *Bax/Bcl-2* ratio is a determining factor  
62 in the regulation of apoptotic cell death.

63 *Leuconostoc mesenteroides* is bacteria sometimes related to fermentation under salinity  
64 and low temperature in fermented foods (Yoon et al., 2018). *L. mesenteroides* is an obligate  
65 heterofermentative lactic acid bacterium that is mostly used in dairy fermentation. *L.*  
66 *mesenteroides* has been studied as a probiotic strain that facilitates the removal of Pb (II)  
67 toxicity (Yi et al., 2017) and inhibits biofilm formation against *Listeria monocytogenes* (Shao  
68 et al., 2019). However, the neuroprotective effects of *L. mesenteroides* have not been studied.  
69 Therefore, the aims of this study were to demonstrate the probiotic properties and  
70 neuroprotective effect of *L. mesenteroides* H40 isolated from kimchi and confirm this effect  
71 in yogurt fermented using *L. mesenteroides* H40.

72

73

## 74 **Materials and Methods**

75

### 76 **Bacterial strains and culture condition**

77 *Lactobacillus fermentum* KU200060, *Lactobacillus brevis* KU200080, and *Leuconostoc*  
78 *mesenteroides* H40 were isolated from kimchi with salted water, mustard leaf (*Brassica*  
79 *juncea*) kimchi, and Chinese cabbage kimchi using lactobacilli MRS medium (MRS; BD  
80 Biosciences, Franklin Lakes, USA) and identified by 16S rRNA analysis (Bionics, Seoul,  
81 Korea). *Lactobacillus rhamnosus* GG (Cell Biotech., Ltd., Gimpo, Korea) was used as a  
82 control strain. Bacteria were propagated and maintained in MRS medium at 37°C for 24 h.

83

#### 84 **Cell culture condition**

85 The HT-29 (human colon adenocarcinoma, KCLB 30038) and SH-SY5Y (human  
86 neuroblastoma, KCLB 22266) cells were used for this study. The cells were grown in  
87 Roswell Park Memorial Institute (RPMI) 1640 medium (Gibco, Grand Island, USA) and  
88 Dulbecco's Modified Eagle's Medium (HyClone Laboratories Inc., Logan, USA),  
89 respectively. All media were accompanied with 10% (v/v) fetal bovine serum (Gibco) and 1%  
90 (v/v) penicillin/streptomycin (Gibco). The cells were maintained at 37°C in 5% CO<sub>2</sub>. The  
91 cultured cells were maintained to monolayer.

92

#### 93 **Tolerance to artificial gastric conditions**

94 To measure the stability against gastric conditions, artificial gastric juice and bile salts  
95 were followed the methods by Yang et al. (2019). The tested strains were incubated in MRS  
96 broth at 37°C for 18 h. Initial cells were inoculated at the concentration of 1×10<sup>7</sup> CFU/mL.  
97 Artificial gastric conditions were dealt on 0.3% pepsin (Sigma-Aldrich, St. Louis, USA)  
98 adjusted to pH 2.5 at 37°C for 3 h. Artificial bile conditions were used 0.3% oxgall (BD  
99 Biosciences) at 37°C for 24 h. After incubation, the survival rate was determined by  
100 calculating viable cells on MRS plates.

101

102        **Adhesion ability to HT-29 cells**

103        The adhesion ability of isolated strains was examined using HT-29. HT-29 cells ( $1 \times 10^5$   
104 cells/mL) was planted in a 24-well cell culture plate and incubated at 37°C (Lee et al., 2015).  
105 After 24 h, isolated strains ( $1 \times 10^7$  CFU/mL) were inoculated and incubated in HT-29 cells at  
106 37°C for 2 h. Non-adherent bacteria were washed three times using PBS buffer (Gibco), 1%  
107 Triton X-100 (Sigma-Aldrich) solution was used for separate the adherent bacteria. The  
108 number of adherent bacteria was determined by dilution and plating on MRS plates.

109

110        **Enzyme production**

111        To measure of enzyme production, the API ZYM kit (BioMerieux, Lyon, France) were  
112 used as manufacture's guideline. Each strain at  $10^6$  CFU/mL was put in each cupule and  
113 incubated at 37°C for 4 h. After incubation, zym A and B reagents put in each cupule, and  
114 represented as production concentration (between 0 and  $\geq 40$  nM).

115

116        **Antibiotic resistance**

117        Antibiotic resistance was followed Clinical and Laboratory Standards Institute guideline  
118 (CLSI, 2012). One hundred microliters of each lactic acid bacteria (LAB) strains ( $1 \times 10^7$   
119 CFU/mL) was inoculated onto MRS agar and paper disc were put on agar plate. Used  
120 antibiotics were ampicillin (10 µg), gentamycin (10 µg), kanamycin (30 µg), ciprofloxacin (5  
121 µg), chloramphenicol (30 µg), streptomycin (10 µg), tetracycline (30 µg), and doxycycline  
122 (30 µg). After incubation at 37°C for 24 h, the inhibitory diameter zone was calculated and  
123 compared to the cut-off value ( $>20$  mm, susceptible; 15-19 mm, intermediate;  $\leq 14$ , resistant)  
124 by represented in CLSI.

125

126        **Conditioned medium (CM) from HT-29 cells**

127 The CM was prepared using HT-29 cells following the method of Park et al. (2017) with  
128 minor modifications. For CM preparation, each sample of LAB strains and yogurt was heated  
129 at 121°C for 15 min and stored at -80°C upto use. HT-29 cells were inoculated into 6-well  
130 plates to  $1.0 \times 10^6$  cells/well and incubated to a confluent monolayer. After incubation, cells  
131 were handled with heat-killed LAB (8 Log CFU/mL) or heated yogurt for 24 h. CM treated  
132 PBS (Gibco) instead to samples were used as control. The mixture was centrifuged ( $12,000 \times g$ ,  
133 10 min) and the supernatant was assembled using a syringe filter (0.45  $\mu$ m pore size,  
134 Millipore Sigma, Burlington, MA, USA).

135

### 136 **Protective effect on oxidative stress-induced apoptosis**

137 To confirm the protective effect on oxidative stress-induced apoptosis, oxidative stress  
138 was induced utilizing  $H_2O_2$  (Junsei Chemical, Tokyo, Japan) or  $NaAsO_2$  (Sigma-Aldrich).  
139 The SH-SY5Y cells (100  $\mu$ L,  $1.0 \times 10^5$  cells/well) were inoculated in 96-well plate with of 50  
140  $\mu$ M  $H_2O_2$  (20  $\mu$ L) or 10  $\mu$ M  $NaAsO_2$  (20  $\mu$ L) for 20 h after pretreatment with 80  $\mu$ L of sample  
141 (CM) for 4 h. After incubation, the media were eliminated, and the cells were incubated with  
142 5 mg/mL MTT solutions (100  $\mu$ L) for 1 h. After incubation, the liquid was removed and  
143 DMSO (100  $\mu$ L) was added to each well. Absorbance was gauged at 540 nm utilizing  
144 microplate reader. The cell viability (%) was calculated as follows:

$$145 \quad \text{Cell viability (\%)} = \frac{\text{Absorbance of sample}}{\text{Absorbance of control}} \times 100$$

146

### 147 **Yogurt production, physicochemical composition, and viable cell counts of LAB**

148 Yogurt was prepare from whole milk (Seoul Milk Co., Ltd., Seoul, Korea) purchased  
149 from a local market. The milk was heated at 90°C for 10 min and cooled to 40°C using water  
150 bath. An overnight culture of *L. mesenteroides* H40 was centrifuged ( $14,000 \times g$ , 10 min, 4°C)  
151 and the cells were washed twice with PBS (Gibco). Then, the pasteurized milk was

152 inoculated with ABT-B commercial yogurt starter culture containing *Lactobacillus*  
153 *acidophilus*, *Lactobacillus delbrueckii* subsp. *bulgaricus*, *Bifidobacterium longum*, and  
154 *Streptococcus thermophilus* (Samik Dairy Co., Ltd., Gimje, Korea) or a mixed culture (1:1)  
155 of *L. mesenteroides* H40 and ABT-B commercial yogurt starter culture. The inoculated  
156 mixture was incubated at 40°C to pH 4.5. Then, the yogurt samples fortified with *L.*  
157 *mesenteroides* H40 were ripened for 24 h in the refrigerator, and its physicochemical  
158 properties were analyzed. Composition and pH of yogurt was analyzed using Milko Scan  
159 Minor (Foss, Hillerod, Denmark) and a pH-meter (WTW inoLab 7110, Weilheim, Germany),  
160 respectively. Titratable acidity was assessed according to AOAC International (1999) by  
161 titration with sodium hydroxide using phenolphthalein. Measurements of viscosity were  
162 performed with Brookfield DV-E Viscometer (Brookfield Eng. Lab. Inc., Middleboro, USA)  
163 using spindle No. 3 at 50 rpm. Viable cell counts of LAB in yogurt samples was confirmed  
164 using decimal dilutions, spread-plated on MRS medium, and incubation at 37°C for 48 h.

### 166 ***BDNF*, *Bax*, and *Bcl-2* expression on oxidative stress-induced apoptosis in SH-SY5Y** 167 **cells**

168 SH-SY5Y cells ( $1.0 \times 10^6$  cells/well) were seeded on 6-well plate and incubated to form a  
169 confluent monolayer. After incubation, the cells were treated with 800  $\mu$ L of CM for 4 h. To  
170 induce oxidative stress, 200  $\mu$ L of H<sub>2</sub>O<sub>2</sub> (50  $\mu$ M) or NaAsO<sub>2</sub> (10  $\mu$ M) was added for 20 h.  
171 Total RNA was isolated using the RNeasy Mini total RNA isolation kit (Cheon et al., 2020;  
172 Park et al., 2017).

### 174 **Real-time polymerase chain reaction**

175 The RNA quality was quantified using microplate reader (Multiscan™ Go, Thermo  
176 Fisher Scientific, Waltham, USA). cDNA was manufactured using cDNA synthesis kit



177 (Thermo Fisher Scientific). Semi-quantitative real-time PCR was performed according to the  
178 PikoReal 96 system (Thermo Fisher Scientific). The reactants contained SYBR Green master  
179 mix, primer (Table 1), cDNA, and RNase free water. Further, 20  $\mu$ L of the mixture was  
180 amplified as 95°C for 2 min as initial denaturation; 40 cycles of 95°C for 5 s as denaturation;  
181 60°C for 15 s as annealing and extension. The results were analyzed by  $\Delta\Delta$ Ct method using  
182 the melt curve analysis method.

183

### 184 **Statistical analysis**

185 All tested data are represented as mean $\pm$ standard deviation by three replicates. One-way  
186 analysis of variance (ANOVA) was utilized to verify significant differences. The mean values  
187 were used for the Duncan's multiple range test to perform post-hoc verification ( $p<0.05$ ).

188

189

## 190 **Results and Discussion**

191

### 192 **Tolerance to artificial gastric conditions and Adhesion to HT-29 cells**

193 *L. fermentum* KU200060, *L. brevis* KU200080, and *L. mesenteroides* H40 was isolated  
194 from various kimchi for probiotic use. *L. rhamnosus* GG, *L. fermentum* KU200060, *L. brevis*  
195 KU200080, and *L. mesenteroides* H40 was confirmed probiotic properties (Table 2;  $p<0.05$ ).  
196 These strains showed high tolerance to artificial gastric conditions. *L. rhamnosus* GG and *L.*  
197 *mesenteroides* H40 decreased to 8.51 Log CFU/mL and 7.17 Log CFU/mL in acidic  
198 conditions, however increased to 8.58 Log CFU/mL and 8.26 Log CFU/mL in bile conditions,  
199 respectively. *L. fermentum* KU200060 and *L. brevis* KU200080 showed strong acid tolerance  
200 having 8.29 Log CFU/mL and 7.91 Log CFU/mL, however decreased to 7.41 Log CFU/mL

201 and 7.75 Log CFU/mL in bile conditions, respectively. *L. plantarum* Ln1 and KCTC 3108  
202 showed similar trends having decrease in acidic conditions and remaining in bile conditions  
203 (Jang et al., 2018).

204 *L. rhamnosus* GG, *L. fermentum* KU200060, *L. brevis* KU200080, and *L. mesenteroides*  
205 H40 showed 2.34%, 1.18%, 3.42%, and 2.86% adhesion rate to HT-29 cells. Especially, *L.*  
206 *brevis* KU200080 and *L. mesenteroides* H40 showed a higher adhesion rate than *L.*  
207 *rhamnosus* GG. Jang et al. (2018) showed lower 2.19% adhesion rate of *L. plantarum* KCTC  
208 3108. Adhered probiotic strains may be temporary colonization and influence host health  
209 through adjustment of intestinal microflora (Jang et al., 2019; Yu et al., 2019a).

210

### 211 **Enzyme production**

212  $\beta$ -Glucuronidase can be produced by the human intestine microbiota and liberate toxin  
213 and mutagen in liver (Dabek et al., 2008). Therefore, isolated strains were confirmed  
214 nonproduction of  $\beta$ -glucuronidase using API ZYM kit (Table 2). *L. rhamnosus* GG produced  
215 30 nM of leucine arylamidase, 30 nM of valine arylamidase, 20 nM of naphthol-AS-BI-  
216 phosphohydrazase, 20 nM of  $\beta$ -galactosidase, and 30 nM of  $\beta$ -glucosidase. *L. fermentum*  
217 KU200060 produced 30 nM of  $\alpha$ -galactosidase and  $\geq 40$  nM of  $\beta$ -galactosidase. *L. brevis*  
218 KU200080 produced 20 nM of  $\beta$ -galactosidase, 30 nM of  $\beta$ -glucosidase, and 30 nM of  
219 leucine arylamidase. *L. mesenteroides* H40 produced 20 nM of  $\alpha$ -glucosidase and 30 nM of  
220  $\beta$ -glucosidase.  $\alpha$ -Galactosidase and  $\beta$ -galactosidase can act the use of indigestible  
221 carbohydrates of raffinose family oligosaccharides and milk products, respectively. In  
222 addition,  $\beta$ -glucosidase may influence bioavailability by the cleavage of glycosidic bonds in  
223 ginsenoside, isoflavone, and phenolic compounds (Son et al., 2018). Produced enzyme by  
224 these isolated strains may be useful for carbohydrate digestion.

225

226 **Antibiotic resistance**

227 *L. fermentum* KU200060 and *L. mesenteroides* H40 are resistant to gentamycin,  
228 kanamycin, and ciprofloxacin. *L. rhamnosus* GG and *L. brevis* KU200080 are resistant to  
229 gentamycin, kanamycin, streptomycin, and ciprofloxacin. Among tested antibiotics, most  
230 *Lactobacillus* sp. are intrinsically resistant to aminoglycoside (gentamycin, kanamycin, and  
231 streptomycin), inhibitors of nucleic acid synthesis (ciprofloxacin) (Campedelli et al., 2015).  
232 Therefore, isolated strains showed a potential of safe probiotic strains in a view of antibiotic  
233 resistance.

234

235 **Protective effects of probiotics-CM on oxidative stress-induced apoptosis in SH-**  
236 **SY5Y cells**

237  $H_2O_2$  and  $NaAsO_2$  converts to a highly reactive toxic hydroxyl radical (Pardillo-Díaz et  
238 al., 2016), causing damage by reducing antioxidant enzymes in brain (Herrera et al., 2013).  
239 Additionally, gut microbiota influence the neurophysicals at the base of the gut-brain axis  
240 (Park et al., 2017). The modulatory effect of probiotics in intestinal microbiota was  
241 demonstrated by increased a ratio of Firmicutes to Bacteriodes and it can relieve  
242 inflammation by cytokine expression (Martin et al., 2018). Therefore, the CM using HT-29  
243 cells with probiotics was used for neuroprotective effects.

244 Oxidative stress was induced in SH-SY5Y cells using  $H_2O_2$  or  $NaAsO_2$ , and cell viability  
245 was confirmed by MTT assay (Fig. 1). During the induction of oxidative stress by  $H_2O_2$ , the  
246 cell viability of SH-SY5Y cells was 53.5% (Fig. 1A;  $p < 0.05$ ). The cell viability of the  
247 probiotics-CM for *L. rhamnosus* GG, *L. fermentum* KU200060, *L. brevis* KU200080, and *L.*  
248 *mesenteroides* H40 was 70.7%, 49.7%, 65.0%, and 69.9%, respectively. *L. rhamnosus* GG, *L.*  
249 *brevis* KU200080, and *L. mesenteroides* H40 showed a protective effect compared to  $H_2O_2$   
250 treated cells (53.5%).

251 During induction of oxidative stress by NaAsO<sub>2</sub>, the cell viability of SH-SY5Y cells was  
252 55.8% (Fig. 1B; p<0.05). The cell viability of the probiotics-CM of *L. rhamnosus* GG, *L.*  
253 *fermentum* KU200060, *L. brevis* KU200080, and *L. mesenteroides* H40 was 55.3%, 49.2%,  
254 55.3%, and 70.7%, respectively. Only *L. mesenteroides* H40 showed a protective effect  
255 compared to NaAsO<sub>2</sub> treated cells (55.8%).

256 Among these strains, *L. mesenteroides* H40 has highest cell viability in SH-SY5Y cells  
257 using both H<sub>2</sub>O<sub>2</sub> and NaAsO<sub>2</sub>. Cheon et al. (2020) showed the cell viability of *L. rhamnosus*  
258 GG (72.0%), *L. fermentum* KU200060 (60.2%), *Lactobacillus delbrueckii* KU2000171  
259 (66.8%), and *L. buchneri* KU200793 (73.4%) with MPP<sup>+</sup> as Parkinson-inducing toxin having  
260 oxidative phosphorylation (Cheon et al., 2020). Therefore, *L. rhamnosus* GG and *L.*  
261 *mesenteroides* H40 was demonstrated neuroprotective effects against oxidative stress.

262

### 263 ***BDNF* mRNA expression and anti-apoptotic effects of probiotics-CM on oxidative** 264 **stress-induced apoptosis in SH-SY5Y cells**

265 The gut-brain axis (GBA) is bi-directional communication network encompassing the  
266 autonomic nervous system (ANS), the central nervous system (CNS), and the enteric nervous  
267 system (ENS). These complex network was influenced by gastrointestinal tract (Kennedy et  
268 al., 2016; Ranuh et al., 2019). Among serum response factor, BDNF have known as regulator  
269 of the synaptic protein and precursors for appropriated neuronal function, survival, and  
270 apoptosis (Numakawa et al., 2010). Decreased *BDNF* mRNA expression confirms brain  
271 related diseases such as Alzheimer's disease, Parkinson's disease, and depression. Increased  
272 ratio of *Bax/Bcl-2* induced apoptosis.

273 *BDNF* mRNA expression and *Bax/Bcl-2* ratio is shown in Fig. 2. Treatment with H<sub>2</sub>O<sub>2</sub>  
274 reduced *BDNF* mRNA expression by 0.73-fold compared with that in H<sub>2</sub>O<sub>2</sub> nontreated cells  
275 (Fig. 2A; p<0.05). *L. rhamnosus* GG and *L. mesenteroides* H40 showed 0.80- and 0.85-fold

276 *BDNF* mRNA expression, respectively. The *Bax/Bcl-2* ratio in H<sub>2</sub>O<sub>2</sub> nontreated cells was  
277 1.00-fold, whereas H<sub>2</sub>O<sub>2</sub> increased the ratio of 2.69-fold (Fig. 2B; p<0.05). Treatment with *L.*  
278 *rhamnosus* GG and *L. mesenteroides* H40 reduced the *Bax/Bcl-2* ratio to 2.24- and 2.03-fold,  
279 respectively.

280 Treatment with NaAsO<sub>2</sub> reduced 0.76-fold *BDNF* mRNA expression compared with that  
281 in the control without NaAsO<sub>2</sub> treatment (Fig. 2C; p<0.05). *L. rhamnosus* GG and *L.*  
282 *mesenteroides* H40 represented 0.95- and 1.08-fold *BDNF* mRNA expression, respectively.  
283 The *Bax/Bcl-2* ratio in NaAsO<sub>2</sub> nontreated cells was 1.00-fold, while NaAsO<sub>2</sub> increased 2.24-  
284 fold in NaAsO<sub>2</sub> treated cells. Treatment with *L. rhamnosus* GG increased 2.61-fold, while  
285 treatment with *L. mesenteroides* H40 reduced 1.46-fold (Fig. 2D; p<0.05).

286 *L. mesenteroides* H40 can increase *BDNF* mRNA expression and reduce apoptosis of  
287 SH-SY5Y cells oxidatively stressed using both H<sub>2</sub>O<sub>2</sub> and NaAsO<sub>2</sub>. The difference of  
288 neuroprotective effect of *L. rhamnosus* GG and *L. mesenteroides* H40 depends on strain and  
289 oxidant.

290

### 291 **Physicochemical property and LAB cell counts of control and probiotic yogurt**

292 Yogurt is a major probiotic carrier to consumers without side-effect. Each yogurt was  
293 manufactured using the following: 1) ABT-B commercial starter culture (control yogurt) and  
294 2) ABT-B commercial starter mixed with *L. mesenteroides* H40 (probiotic yogurt). The fat,  
295 protein, lactose, total solids, and acidity content are shown in Table 3 (p<0.05). Probiotic  
296 yogurt made using *L. mesenteroides* H40 had 2.96% fat, 3.23% protein, 6.16% lactose, and  
297 27.33% total solids. In addition, probiotic yogurt was not significantly different from control  
298 yogurt. However, probiotic yogurt exhibited significantly higher viscosity than control yogurt.  
299 Texture of stirred yogurt is the result of both acid aggregation of casein micelles by ropy  
300 strains during incubation (Zhao et al., 2016). The viable cell counts of lactic acid bacteria in

301 control and probiotic yogurt with H40 was  $8.66 \pm \text{Log CFU/mL}$  and  $8.96 \pm \text{Log CFU/mL}$ ,  
302 respectively (data not shown).

303

### 304 **Protective effects of Y-CM oxidative stress-induced apoptosis in SH-SY5Y** 305 **neuroblastoma cells**

306 Y-CM was manufactured with HT-29 cells and yogurt, and its neuroprotective effect  
307 was assessed in SH-SY5Y cells (Table 4;  $p < 0.05$ ). The treatment of  $\text{H}_2\text{O}_2$  reduced cell  
308 viability of SH-SY5Y cells to 55.5%. However, cell viability of control yogurt CM (CY-CM)  
309 and probiotic yogurt CM (PY-CM) was 72.2% and 114.8%, respectively. Under treatment  
310 with  $\text{NaAsO}_2$ , cell viability of positive control was 51.4% and that of CY-CM and PY-CM  
311 was 49.9% and 109.5%, respectively. The PY-CM using *L. mesenteroides* H40 showed high  
312 cell viability in oxidatively stressed SH-SY5Y cells in both  $\text{H}_2\text{O}_2$  and  $\text{NaAsO}_2$  treatment.  
313 When compare Fig. 1 and Table 4, PY-CM showed higher cell viability than *L. mesenteroides*  
314 H40. These results showed that the PY-CM effectively protected the cells from oxidative  
315 damage caused by  $\text{H}_2\text{O}_2$  and  $\text{NaAsO}_2$ .

316  $\text{H}_2\text{O}_2$  treatment resulted in a 0.78-fold increase in *BDNF* mRNA expression compared  
317 with that in the  $\text{H}_2\text{O}_2$  nontreated cells (Table 4;  $p < 0.05$ ). CY-CM and PY-CM increased  
318 *BDNF* mRNA expression by 0.83- and 1.12-fold, respectively. The ratio of *Bax/Bcl-2* ratio in  
319  $\text{H}_2\text{O}_2$  nontreated cells was 1.00, while  $\text{H}_2\text{O}_2$  increased to 2.69-fold. The treatment with CY-  
320 CM and PY-CM reduced the *Bax/Bcl-2* ratio to 2.05- and 1.24-fold, respectively.

321 The treatment with  $\text{NaAsO}_2$  reduced *BDNF* mRNA expression by 0.76-fold compared to  
322 that in  $\text{NaAsO}_2$  nontreated cells (Table 4;  $p < 0.05$ ). CY-CM and PY-CM treatment resulted in  
323 a 1.03- and 1.18-fold *BDNF* mRNA expression, respectively. The ratio of *Bax/Bcl-2* ratio in  
324  $\text{NaAsO}_2$  nontreated cells was 1.00, while  $\text{NaAsO}_2$  treatment increased this to 2.24-fold in

325 NaAsO<sub>2</sub> treated cells. The treatment with CY-CM and PY-CM reduced to 1.88- and 1.32-fold,  
326 respectively. Thus, PY-CM can reduce apoptosis of H<sub>2</sub>O<sub>2</sub> or NaAsO<sub>2</sub> stressed SH-SY5Y cells.

327 The treatment with PY-CM significantly increased *BDNF* mRNA expression and  
328 reduced apoptosis on SH-SY5Y mRNA. In addition, mRNA expression was markedly higher  
329 based on the yogurt type than with the strain alone or with the yogurt and probiotic strain.  
330 These synergistic effects originate from whey protein containing methionine, lysine, and  
331 proline, which are associated with apoptosis (Lee and Hur, 2019). Therefore, PY-CM using *L.*  
332 *mesenteroides* H40 has potent neuroprotective effect in preventing the oxidative stress  
333 induced by H<sub>2</sub>O<sub>2</sub> and NaAsO<sub>2</sub>.

334

335

## 336 **Conclusions**

337 *L. mesenteroides* H40 was isolated from kimchi and its probiotic property was  
338 demonstrated through stability in gastric conditions, adhesion to intestinal cells, enzyme  
339 production, and safe antibiotic resistance. For neuroprotective effect, *L. mesenteroides* H40-  
340 CM confirmed an increase of cell viability with increase of *BDNF* mRNA expression and  
341 decrease of the *Bax/Bcl-2* ratio in oxidatively stress-induced SH-SY5Y cells. Compared to  
342 probiotic strain and yogurt type, the probiotic yogurt showed a higher neuroprotective effect  
343 than the strain alone. Therefore, these results suggested a potential of prophylactic therapy as  
344 probiotics and functional dairy products. In addition, the cognitive function-related  
345 experiments will need to be performed in humans for efficacy verification.

346

## 347 **Conflict of Interest**

348 The authors declare no potential conflict of interest.

349

## 350 **References**

351 AOAC International. 1999. AOAC International association of caseins and casein micelle  
352 structure (16th), AOAC Int, Washington, DC.

353 Azmi NH, Ismail N, Imam MU, Ismail M. 2013. Ethyl acetate extract of germinated brown  
354 rice attenuates hydrogen peroxide-induced oxidative stress in human SH-SY5Y  
355 neuroblastoma cells: Role of anti-apoptotic, pro-survival and antioxidant genes. BMC  
356 Complem Altern M 13:177.

357 Campedelli I, Mathur H, Salvetti E, Clarke S, Rea MC, Torriani S, Ross RP, Hill C, O'Toole  
358 PW. 2019. Genus-wide assessment of antibiotic resistance in *Lactobacillus* spp. Appl  
359 Environ Microbiol 85:e01738.

360 Cheon MJ, Lim SM, Lee NK, Paik HD. 2020. Probiotic properties and neuroprotective  
361 effects of *Lactobacillus buchneri* KU200793 isolated from Korean fermented foods. Int J  
362 Mol Sci 21:1227.

363 Clinical and Laboratory Standards Institute. 2012. Performance standards for antimicrobial  
364 susceptibility testing; approved standard—11th ed CLSI document M02-A11. Clinical  
365 and Laboratory Standards Institute, Wayne, PA.

366 Dabek M, McCrae SJ, Stevens VJ, Duncan SH, Lous P. 2008. Distribution of  $\beta$ -glucosidase  
367 and  $\beta$ -glucuronidase activity and of  $\beta$ -glucuronidase gene *gus* in human colonic bacteria.  
368 FEMS Microbiol Ecol 66:487-495.

369 Dussert S, Davey MW, Laffargue A, Doulbeau S, Swennen R, Etienne H. 2006. Oxidative  
370 stress, phospholipid loss and lipid hydrolysis during drying and storage of intermediate  
371 seeds. Physiol Plant 127:192-204.

372 Elahi MM, Kong YX, Matata BM. 2009. Oxidative stress as a mediator of cardiovascular



373 disease. *Oxid Med Cell Longev* 2:259-269.

374 Egan MF, Kojima M, Callicott JH, Goldberg TE, Kolachana BS, Bertolino A, Zaitsev E, Gold  
375 B, Goldman D, Dean M, Lu B, Weinberger D. 2003. The BDNF val66met polymorphism  
376 affects activity-dependent secretion of BDNF and human memory and hippocampal  
377 function. *Cell* 112:257-269.

378 Ghaisas S, Maher J, Kanthasamy A. 2016. Gut microbiome in health and disease: Linking the  
379 microbiome–gut–brain axis and environmental factors in the pathogenesis of systemic and  
380 neurodegenerative diseases. *Pharmacol Ther* 158:52-62.

381 Herrera A, Pineda J, Antonio MT. 2013. Toxic effects of perinatal arsenic exposure on the  
382 brain of developing rats and the beneficial role of natural antioxidants. *Environ Toxicol*  
383 *Pharmacol* 36:73-79.

384 Ishimwe N, Daliri EB, Lee BH, Fang F, Du G. 2015. The perspective on cholesterol-lowering  
385 mechanisms of probiotics. *Mol Nutr Food Res* 59:94-105.

386 Jang HJ, Song MW, Lee NK, Paik HD. 2018. Antioxidant effects of live and heat-killed  
387 probiotic *Lactobacillus plantarum* Ln1 isolated from kimchi. *J Food Sci Technol*  
388 55:3174-3180.

389 Jang HJ, Lee NK, Paik HD. 2019. Probiotic characterization of *Lactobacillus brevis*  
390 KU15153 showing antimicrobial and antioxidant effect isolated from kimchi. *Food Sci*  
391 *Biotechnol* 28:1521-1528.

392 Kennedy PJ, Murphy AB, Cryan JF, Ross PR, Dinan TG, Stanton C. 2016. Microbiome in  
393 brain function and mental health. *Trends Food Sci Tech* 57:289-301.

394 Klippel BF, Duemke LB, Leal MA, Friques AGF, Dantas EM, Dalvi RF, Gava AL, Pereira  
395 TMC, Andrade TU, Meyrelles SS, Campagnaro BL, Vasquez EC. 2016. Effects of kefir  
396 on the cardiac autonomic tones and baroreflex sensitivity in spontaneously hypertensive  
397 rats. *Front Physiol* 7:211.

398 Lee NK, Kim SY, Han KJ, Eom SJ, Paik HD. 2014. Probiotic potential of *Lactobacillus*  
399 strains with anti-allergic effects from kimchi for yogurt starters. *LWT-Food Sci Technol*  
400 58:130-134.

401 Lee NK, Han KJ, Son SH, Eom SJ, Lee SK, Paik HD. 2015. Multifunctional effect of  
402 probiotic *Lactococcus lactis* KC24 isolated from kimchi. *LWT-Food Sci Technol* 64:1036-  
403 1041.

404 Lee SY, Hur SJ. 2019. Mechanisms of neuroprotective effects of peptides derived from  
405 natural materials and their production and assessment. *Compr Rev Food Sci F* 18:923-935.

406 Liu L, Cavanaugh JE, Wang Y, Sakagami H, Mao Z, Xia Z. 2003. ERK5 activation of MEF2-  
407 mediated gene expression plays a critical role in BDNF-promoted survival of developing  
408 but not mature cortical neurons. *Proc Natl Acad Sci U.S.A* 100:8532-8537.

409 Lobo V, Patil A, Phatak A, Chandra N. 2010. Free radicals, antioxidants and functional foods:  
410 Impact on human health. *Pharmacogn Rev* 4:118-126.

411 Martin CR, Osadchiy V, Kalani A, Mayer EA. 2018. The brain-gut-microbiome axis. *CMGH*  
412 6:133-148.

413 Numakawa T, Suzuki S, Kumamaru E, Adachi N, Richards M, Kunugi H. 2010. BDNF  
414 function and intracellular signaling in neurons. *Histol Histopathol* 25:237-258.

415 O'Toole PW, Marchesi JR, Hill C. 2017. Next-generation probiotics: the spectrum from  
416 probiotics to live biotherapeutics. *Nat Microbiol* 2:17057.

417 Pamplona R, Barja G. 2006. Mitochondrial oxidative stress, aging and caloric restriction: The  
418 protein and methionine connection. *Biochim Biophys Acta* 1757:496-508.

419 Park J, Lee J, Yeom Z, Heo D, Lim YH. 2017. Neuroprotective effect of *Ruminococcus albus*  
420 on oxidatively stressed SH-SY5Y cells and animals. *Sci Rep* 7:14520.

421 Pardillo-Díaz R, Carrascal L, Muñoz MF, Ayala A, Nunez-Abades P. 2016. Time and dose  
422 dependent effects of oxidative stress induced by cumene hydroperoxide in neuronal

423 excitability of rat motor cortex neurons. *Neurotoxicology* 53:201-214.

424 Quigley EM. 2017. Microbiota-brain-gut axis and neurodegenerative diseases. *Curr Neurol*  
425 *Neurosci Rep* 17:94.

426 Ranuh R, Athiyah AF, Darma A, Risky VP, Riawan W, Surono IS, Sudarmo SM. 2019.  
427 Effect of the probiotic *Lactobacillus plantarum* IS-10506 on BDNF and 5HT stimulation:  
428 role of intestinal microbiota on the gut-brain axis. *Iran J Microbiol* 11:145-150.

429 Shao X, Fang K, Medina D, Wan J, Lee JL, Hong SH. 2019. The probiotic, *Leuconostoc*  
430 *mesenteroides*, inhibits *Listeria monocytogenes* biofilm formation. *J Food Saf*  
431 2019:e12750.

432 Son SH, Jeon H., Yang SJ, Sim MH, Kim YJ, Lee NK, Paik HD. 2018. Probiotic lactic acid  
433 bacteria isolated from traditional Korean fermented foods based on  $\beta$ -glucosidase activity.  
434 *Food Sci Biotechnol* 27:123-129.

435 Sun J, Ling Z, Wang F, Chen W, Li H, Jin J, Zhang H, Pang M, Yu J, Liu J. 2016. *Clostridium*  
436 *butyricum* pretreatment attenuates cerebral ischemia/reperfusion injury in mice via anti-  
437 oxidation and anti-apoptosis. *Neurosci Lett* 613:30-35.

438 Ton AMM, Campagnaro BP, Alves GA, Aires R, Côco LZ, Arpini CM, Oliveira TG, Campos-  
439 Toimil M, Meyrelles SS, Pereira TMC, Vasquez EC. 2020. Oxidative stress and dementia  
440 in Alzheimer's patients: Effects of synbiotic supplementation. *Oxid Med Cell Longev*  
441 2020:2638703.

442 Wang H, Lee IS, Braun C, Enck P. 2016. Effect of probiotics on central nervous system  
443 functions in animals and humans: A systematic review. *J Neurogastroenterol Motil*  
444 22:589-605.

445 Wang X, Michaelis EK. 2010. Selective neuronal vulnerability to oxidative stress in the  
446 brain. *Front Aging Neurosci* 2:12.

447 Wang Y, Kasper LH. 2014. The role of microbiome in central nervous system

448 disorders. *Brain Behav Immun* 38:1-12.

449 Yang SJ, Lee JE, Lim SM, Kim YJ, Lee NK, Paik HD. 2019. Antioxidant and immune-  
450 enhancing effects of probiotic *Lactobacillus plantarum* 200655 isolated from kimchi.  
451 *Food Sci Biotechnol* 28:491-499

452 Yi YJ, Lim JM, Lim JM, Gu S, Lee WK, Oh E, Lee SM, Oh BT. 2017. Potential use of lactic  
453 acid bacteria *Leuconostoc mesenteroides* as a probiotic for the removal of Pb (II) toxicity.  
454 *J Microbiol* 55:296-303.

455 Yoon JY, Kim D, Kim EB, Lee SK, Lee M, Jang A. 2018. Quality and lactic acid bacteria  
456 diversity of pork salami containing kimchi powder. *Food Sci Anim Resour* 38:912-926.

457 Yu HS, Jang HJ, Lee NK, Paik HD. 2019a. Evaluation of the probiotic characteristics and  
458 prophylactic potential of *Weissella cibaria* strains isolated from kimchi. *LWT-Food Sci*  
459 *Technol* 112:108229.

460 Yu HS, Lee NK, Choi AJ, Choe JS, Bae CH, Paik HD. 2019b. Antagonistic and antioxidant  
461 effect of probiotic *Weissella cibaria* JW15. *Food Sci Biotechnol* 28:851-855.

462 Yu HS, Lee NK, Choi AJ, Choe JS, Bae CH, Paik HD. 2019c. Anti-inflammatory potential of  
463 probiotic strain *Weissella cibaria* JW15 isolated from kimchi through regulation of NF-  
464  $\kappa$ B and MAPKs pathways in LPS-induced RAW 264.7 cells. *J Microbiol Biotechnol*  
465 29:1022-1032.

466 Zhao LL, Wang XL, Tian Q, Mao XY. 2016. Effect of casein to whey protein ratios on the  
467 protein interactions and coagulation properties of low-fat yogurt. *J Dairy Sci* 99:7768-  
468 7775.

469

470 **Table 1. Primer sequence for neuroprotective effect used in semi-quantitative real-time**  
 471 **PCR**

Gene		Primer sequence	References
<i>GAPDH</i>	Forward	5' GAGTCAACGGATTTGGTCGT 3'	
	Reverse	5' GACAAGCTTCCCGTTCTCAG 3'	
<i>BDNF</i>	Forward	5' CAAACATCCGAGGACAAGGTGG 3'	
	Reverse	5' CTCATGGACATGTTTGCAGCATCT 3'	
<i>Bax</i>	Forward	5' GTGGTTGCCCTCTTCTACTTTGC 3'	Park et al., 2017
	Reverse	5' GAGGACTCCAGCCACAAAGATG 3'	
<i>Bcl-2</i>	Forward	5' CGGCTGAAGTCTCCATTAGC 3'	
	Reverse	5' CGGCTGAAGTCTCCATTAGC 3'	

472 *BDNF*, brain-derived neurotrophic factor; *Bax*, *Bcl-2*-associated X protein; *Bcl-2*, B-cell  
 473 lymphoma.

474

475

476 **Table 2. Probiotic properties of isolated strains**

Treatment	LGG <sup>a</sup>	200060	200080	H40
Tolerance to artificial gastric conditions (Viable cell number (Log CFU/mL))				
Initial cell number	8.55±0.01 <sup>a</sup>	8.26±0.01 <sup>b</sup>	7.79±0.01 <sup>c</sup>	8.17±0.03 <sup>b</sup>
0.3% (w/v) pepsin, pH 2.5, 3 h	8.51±0.05 <sup>a</sup>	8.29±0.01 <sup>ab</sup>	7.91±0.00 <sup>b</sup>	7.17±0.01 <sup>c</sup>
0.3% (w/v) oxgall, 24 h	8.58±0.03 <sup>a</sup>	7.41±0.01 <sup>bc</sup>	7.75±0.02 <sup>b</sup>	8.26±0.01 <sup>a</sup>
Adhesion rate (%) <sup>b</sup>	2.34±0.26 <sup>c</sup>	1.18±0.08 <sup>d</sup>	3.42±0.49 <sup>a</sup>	2.86±0.16 <sup>b</sup>
β-Glucuronidase (nM)	0	0	0	0
Antibiotic resistance	Gentamycin, kanamycin, streptomycin, ciprofloxacin	Gentamycin, kanamycin, ciprofloxacin	Gentamycin, kanamycin, streptomycin, ciprofloxacin	Gentamycin, kanamycin, ciprofloxacin

477 <sup>1</sup>LGG, *L. rhamnosus* GG; 200060, *L. fermentum* KU200060; 200080, *L. brevis* KU200080;  
478 H40, *L. mesenteroides* H40

479 <sup>2</sup>Adhesion rate = (adhered bacteria to HT-29 cells after 2 h)/(initial bacteria)×100

480 Data are represented as the mean±standard deviation of triplicate experiments. Means within  
481 a row with same superscript differ (p<0.05).

482

483 **Table 3. Physicochemical properties of control and probiotic yogurt**

Physicochemical properties	Yogurt type	
	Control yogurt <sup>1</sup>	Probiotic yogurt <sup>2</sup>
Fat (%)	2.96±0.05 <sup>a</sup>	2.96±0.20 <sup>a</sup>
Protein (%)	3.16±0.20 <sup>a</sup>	3.23±0.11 <sup>a</sup>
Lactose (%)	6.23±0.11 <sup>a</sup>	6.16±0.15 <sup>a</sup>
Total solid (%)	27.66±0.15 <sup>a</sup>	27.33±0.28 <sup>a</sup>
Titrateable acidity	0.82±0.01 <sup>a</sup>	0.81±0.01 <sup>a</sup>
pH	4.33±0.04 <sup>a</sup>	4.26±0.03 <sup>a</sup>
Viscosity (cP)	1,724.20±15.60 <sup>a</sup>	2,048.30±7.30 <sup>b</sup>

484 <sup>1</sup>Control yogurt, yogurt manufactured by ABT-B starter.

485 <sup>2</sup>Probiotic yogurt, yogurt manufactured by ABT-B starter and *L. mesenteroides* H40.

486 Data are represented as the mean±standard deviation of triplicate experiments. Means within  
 487 a row with same superscript differ (p<0.05).

488

489 **Table 4. Neuroprotective effect of control and probiotic yogurt**

Characteristics	Oxidant	
	H <sub>2</sub> O <sub>2</sub> (50 μM)	NaAsO <sub>2</sub> (10 μM)
Cell viability (%)		
Non-treatment	100±5.43 <sup>c</sup>	100±11 <sup>b</sup>
Control	55.45±0.78 <sup>a</sup>	51.37±3.5 <sup>a</sup>
CY-CM <sup>1</sup>	72.21±1.55 <sup>b</sup>	49.92±1.50 <sup>a</sup>
PY-CM <sup>2</sup>	114.76±9.30 <sup>d</sup>	109.5±11.5 <sup>b</sup>
BDNF expression (fold of control)		
Non-treatment	1.00±0.06 <sup>b</sup>	1.00±0.07 <sup>b</sup>
Control	0.73±0.02 <sup>a</sup>	0.76±0.03 <sup>a</sup>
CY-CM <sup>1</sup>	0.78±0.12 <sup>a</sup>	1.03±0.02 <sup>b</sup>
PY-CM <sup>2</sup>	1.05±0.05 <sup>b</sup>	1.18±0.02 <sup>b</sup>
<i>Bax/Bcl-2</i> ratio		
Non-treatment	1.00±0.08 <sup>a</sup>	1.00±0.07 <sup>a</sup>
Control	2.69±0.08 <sup>c</sup>	2.25±0.04 <sup>b</sup>
CY-CM <sup>1</sup>	2.05±0.07 <sup>b</sup>	1.88±0.21 <sup>b</sup>
PY-CM <sup>2</sup>	1.24±0.13 <sup>a</sup>	1.32±0.17 <sup>a</sup>

490 <sup>1</sup>CY-CM, conditioned medium using yogurt manufactured by ABT-B starter.

491 <sup>2</sup>PY-CM, conditioned medium using yogurt manufactured by ABT-B starter and *L.*  
 492 *mesenteroides* H40.

493 Data are represented as the mean±standard deviation of triplicate experiments.

494 Means within same characteristics a row with same superscript differ (p<0.05).

495



## Figure legends

496

497

498 **Fig. 1.** Cell viability of conditioned medium using lactic acid bacteria (LAB-CM) in SH-SY5Y  
499 cells with oxidative stress induced by (A) H<sub>2</sub>O<sub>2</sub> (50 μM) and (B) NaAsO<sub>2</sub> (10 μM). LGG, *L.*  
500 *rhamnosus* GG; 200060, *L. fermentum* KU200060; 200080, *L. brevis* KU200080; H40, *L.*  
501 *mesenteroides* H40. Error bars indicate standard deviation from three independent experiments.  
502 Different letters on each bar represent significantly different (p<0.05).

503

504 **Fig. 2.** mRNA expression levels of *BDNF* and apoptosis-related genes on oxidatively stressed  
505 SH-SY5Y cells treated with conditioned medium using lactic acid bacteria (LAB-CM). (A)  
506 *BDNF* mRNA expression and (B) *Bax/Bcl-2* ratio in oxidative stress-induced SH-SY5Y cells  
507 induced by H<sub>2</sub>O<sub>2</sub> (50 μM). (C) *BDNF* mRNA expression and (D) *Bax/Bcl-2* ratio in oxidatively  
508 stress-induced SH-SY5Y cells induced by NaAsO<sub>2</sub> (10 μM). LGG, *L. rhamnosus* GG; H40, *L.*  
509 *mesenteroides* H40. Error bars indicate standard deviation from three independent experiments.  
510 Different letters on each bar represent significantly different (p<0.05).

511


512

513 **Fig. 1.**


514

515

(A)




(B)




516


(A)



(B)



(C)



(D)

