

TITLE PAGE
- Food Science of Animal Resources -
 Upload this completed form to website with submission

ARTICLE INFORMATION	Fill in information in each box below
Article Type	Research article
Article Title	Probiotics increase intramuscular fat and improve the composition of fatty acids in Sunit sheep through the AMPK signaling pathway
Running Title (within 10 words)	Probiotics regulate intramuscular fat and fatty acid deposition in sheep
Author	Yue Zhang ^{1,2} , Duo Yao ³ , Huan Huang ¹ , Min Zhang ^{1,2} , Lina Sun ^{1,2} , Lin Su ^{1,2} , LiHua Zhao ^{1,2} , Yueying Guo ^{1,2} , Ye Jin ^{1,2*}
Affiliation	1 Inner Mongolia Agricultural University, Hohhot 010018, China 2 Integrative Research Base of Beef and Lamb Processing Technology, Ministry of Agriculture and Rural Affairs of the People's Republic of China, Hohhot 010018, China 3 Inner Mongolia Institute of Quality and Standardization, Hohhot, 010070, China
Special remarks – if authors have additional information to inform the editorial office	
ORCID (All authors must have ORCID) https://orcid.org	Yue Zhang (https://orcid.org/0000-0002-0189-9287) Duo Yao (https://orcid.org/0000-0002-9937-9182) Huan Huang (https://orcid.org/0000-0002-0804-9541) Min Zhang (https://orcid.org/0000-0002-2662-1142) Lin Su (https://orcid.org/0000-0002-2094-2906) LiHua Zhao (https://orcid.org/0000-0002-3334-1755) Lina Sun (https://orcid.org/0000-0003-4450-5038) Yueying Guo (https://orcid.org/0000-0002-0158-1850) Ye Jin (https://orcid.org/0000-0001-8960-879X)
Conflicts of interest List any present or potential conflicts of interest for all authors. (This field may be published.)	The authors declare no potential conflicts of interest.
Acknowledgements State funding sources (grants, funding sources, equipment, and supplies). Include name and number of grant if available. (This field may be published.)	This work was supported by the National Natural Science Foundation of China (32160589); the Major Science and Technology Projects of Ordos (2022EEDSKJZDZX029); the Major R&D and Achievements Transformation Projects of Inner Mongolia (2022YFXZ0017); the Natural Science Foundation of Inner Mongolia (2020ZD11); the Natural Science Foundation of Inner Mongolia (2022MS03055); the Natural Science Foundation of Inner Mongolia (2021MS03010) and the Scientific Research Projects of Universities Directly under Inner Mongolia (BR22-13-08).
Author contributions (This field may be published.)	Conceptualization: Zhang Y, Jin Y Data curation: Zhang Y, Yao D Formal analysis: Huang H, Zhang M Methodology: Zhang Y, Su L, Sun L Software: Zhang Y Validation: Zhao L, Guo Y, Jin Y Investigation: Zhang Y, Yao D, Huang H Writing - original draft: Zhang Y Writing - review & editing: Zhang Y, Yao D, Huang H, Zhang M, Su L, Zhao L, Sun L, Guo Y, Jin Y
Ethics approval (IRB/IACUC) (This field may be published.)	The study was conducted in accordance with the Declaration of Helsinki, and all animals handling procedures were performed in accordance with the Ethics Committee of Inner Mongolia Agricultural University (permit number: NND2021072).

CORRESPONDING AUTHOR CONTACT INFORMATION

For the <u>corresponding</u> author (responsible for correspondence, proofreading, and reprints)	Fill in information in each box below
First name, middle initial, last name	Ye, Jin
Email address – this is where your proofs will be sent	jinyeyc@sohu.com.
Secondary Email address	zhangyueimau@163.com
Postal address	010018
Cell phone number	13948111209
Office phone number	86 0471 436 2690
Fax number	86 0471 436 2690

ACCEPTED

Probiotics increase intramuscular fat and improve the composition of fatty acids in Sunit sheep through the AMPK signaling pathway

Abstract

This experiment aims to investigate the impact of probiotic feed on growth performance, carcass traits, plasma lipid biochemical parameters, intramuscular fat and triglyceride content, fatty acid composition, as well as mRNA expression levels of genes related to lipid metabolism and corresponds to the activity of the enzyme in Sunit sheep. In this experiment, 12 of 96 randomly selected Sunit sheep were assigned to receive the basic diet or the basic diet supplemented with probiotics. The results showed that supplementation with probiotics significantly increased the loin eye area, and decreased plasma triglycerides, low-density lipoproteins, and free fatty acids, increasing the content of intramuscular fat and triglycerides in the muscle and improving the composition of the fatty acids including arachidonic acid, γ -linolenic acid, eicosapentaenoic acid, and so on. Furthermore, the inclusion of probiotics in the diet reduced the expression of AMPK α 2 mRNA and CPT1B mRNA, while increasing the expression of ACC α mRNA, SREBP-1c mRNA, FASN mRNA, and SCD1 mRNA. The results of this study indicate that the addition of probiotics to the diet regulates fat deposition and improves the composition of fatty acids in Sunit sheep through the signaling pathways AMPK-ACC-CPT1B and AMPK-SREBP-1c. This regulatory mechanism leads to an increase in intramuscular fat content, a restructuring of muscle composition of the fatty acids, and an enhancement of the

nutritional value of meat. These findings contribute to a better understanding of the food science of animal resources and provide valuable references for the production of meat of higher nutritional value.

Keywords: probiotics, intramuscular fat, fatty acid composition, AMPK signaling pathway.

Introduction

Intramuscular fat (IMF) exists around muscle fibers embedded in connective tissue sheaths and is considered the final product of the fat deposition process. It is one of the primary indicators to evaluate muscle quality and positively influences quality traits of meat, including flavor, juiciness, and tenderness. (Chikwanha et al., 2018; Pećina et al., 2018; Wang et al., 2020). In addition to the content of IMF in meat, the composition of those fatty acids has also received increasing attention due to its possible implications for human health. In most studies, there was a significant positive correlation between monounsaturated fatty acids (MUFAs) and IMF content, while a negative relationship existed between IMF and most polyunsaturated fatty acids (PUFAs) (Razmaitė et al., 2020). There are different sources of fatty acids in the IMF, which are de novo synthesis by the animal body and direct intake of fatty acids from the diet (Bessa et al., 2015). As a meat-type sheep, the contribution of de novo synthesis to fat deposition is relatively small (Lalotiotis et al., 2010). Therefore, optimizing the deposition and composition of the IMF through sheep production and

feeding strategies will provide new opportunities for mutton producers to enhance meat quality characteristics (Juárez et al., 2021).

The probiotics feed improves the digestive and absorption functions of the host animals and enhances their immunity by altering their intestinal microbiota, which has a positive impact on the host animals. *Lactobacillus casei* has been proven to possess cholesterol-lowering and immune-modulating effects, with beneficial effects on host health (Ayala-Monter et al., 2019; Wang et al., 2021). *Lactobacillus plantarum* has been confirmed to reduce lipid level, and improve the functionality of the gut microbiota (Bao et al., 2012; Li et al., 2020). The probiotic preparation (*Lactobacillus casei* and *Lactobacillus plantarum*) possess the ability to simultaneously reduce serum cholesterol levels and improve the fatty acid composition of meat and milk (Rabelo et al., 2016; Toral et al., 2018). Dev concluded that a compound *Lactobacillus casei* preparation fed to sheep can increase body weight gain and strengthen immunity (Dev et al., 2021). After the mixed culture of *Lactobacillus* was fed to creole goats, tissues of the animals in the probiotics group exhibited an increase in MUFAs and PUFAs, including linoleic acid (C18:2), linolenic acid (C18:3), and cis-9, trans-11 conjugated linoleic acid (CLA) (Taboada et al., 2022). Lamb in the prebiotic group had better antioxidant status and water-holding capacity, and the meat was a tenderer (Nie et al., 2022). From these perspectives, dietary probiotics seem to be a technical method to improve animal performance, fatty acid composition, and meat quality. However, the molecular regulatory mechanism by which probiotics feed affects the IMF content and fatty acid composition in Sunit sheep is unclear.

The AMP-activated protein kinase (AMPK) signaling pathway is a major metabolic pathway that regulates lipid decomposition and synthesis. On the one hand, when AMPK is activated, the cascade effect of the AMPK signaling pathway that regulates fat metabolism is activated. AMPK is phosphorylated to rapidly inhibit the activity of acetyl-CoA carboxylase (ACC), the expression of ACC is reduced, and the catalyzed malonyl-CoA by ACC is reduced (malonyl-CoA is the main component of de novo lipogenesis, and it is an inhibitor of Carnitine palmitoyltransferase 1 (CPT1)) (Ke et al., 2018; Fang et al., 2019). This cascade reaction controls the rate at which free fatty acids in the organism enter mitochondria for β -oxidation, suggesting a central role for the AMPK-ACC signaling pathway in the regulation of lipolysis and synthesis (Thomson and Winder, 2009; Zhang et al., 2021). On the other hand, as a downstream target gene for AMPK, the sterol regulatory element-binding protein-1c (SREBP-1c) is a transcription factor that regulates lipid homeostasis by controlling the synthesis of cholesterol, fatty acids, triglycerides, and phospholipids (Ferré et al., 2021). AMPK can inhibit the activity of SREBP-1c by phosphorylation, and SREBP-1c can release the active fragment to bind to the SREBP-1c response element on its downstream target gene, fatty acid synthase (FASN), which theoretically regulates fatty acid metabolism (Tong et al., 2016; Yang et al., 2017; Fang et al., 2019). Based on this, we hypothesized that the AMPK signaling pathway is involved in the regulation of IMF and fatty acid composition in Sunit sheep by probiotics feed.

This study assumed that long-term supplementation with probiotics could affect the content of IMF and the composition of fatty acids of Sunit sheep through the

AMPK signaling pathway is contribute to improving the nutritional value of meat. This study aimed to investigate whether probiotics feed can be an effective means of promoting IMF deposition in Sunit sheep and to further evaluate the role of the AMPK signaling pathway in this process, thus revealing possible potential mechanisms by which probiotics feed regulates IMF and fatty acid composition in Sunit sheep. This study provides new ideas and strategies for the application of probiotics in animal husbandry.

Materials and Methods

Ethics Statement

The study was approved by the Institutional Animal Care and Use Committee of Inner Mongolia Agricultural University (permit number: NND2021072).

Experimental design and diets

Twelve male sheep of ninety-six three-month-old Sunit sheep of similar live weight were randomly selected and divided into two groups for two different dietary treatments, one without probiotics (CON) and the other with probiotics (PRO). The foundation diet provided by the CON group consisted of a mixture of hay and concentrate composed primarily of cornmeal, fed at a rate of 1.5% of the sheep's body weight. The PRO group was provided with the same basic diet but with 1% probiotics added to the concentrate. All experimental diets met the nutritional requirements of the sheep (Table 1). All sheep were individually housed in separate pens equipped

with feeders. They were fed twice daily at 8:00 and 17:00. All sheep had ad libitum access to feed. Throughout the entire experimental procedure, no dietary selectivity occurred, and no instances of refusal to consume forage and/or concentrate were observed. The probiotics used in this study were a mixture of equal parts *Lactobacillus casei* HM-09 and *Lactobacillus plantarum* HM-10, which were provided by Inner Mongolia Agricultural University (China), the Key Laboratory of Dairy Biotechnology and Engineering of the Ministry of Education. Both of these probiotic strains were identified by screening fermented mare's milk from Inner Mongolia.

All sheep were raised in a well-ventilated farmhouse with the same environmental conditions in Bayan Nur City, Inner Mongolia Autonomous Region, China. The light was provided naturally. The sheep were born on the farm and reared with their ewes until stopping breastfeeding about 90 days of age when they were raised alone. Sheep were allowed access to drinking water and salt blocks throughout the experiment. After 90 days of experimental feeding, all animals fasted for 12 hours and then were transported to a commercial slaughterhouse, where they were immediately slaughtered according to standard commercial 'halal' procedures by professional personnel (Abdullah et al., 2019).

Blood collection and analysis

On the last day of the experiment, all sheep were subjected to venous blood collection using a 10 ml vacuum blood collection tube containing anticoagulant in the

morning. Blood collected was separated by centrifugation at 4°C at 3200×g for 15 min, and serum was stored at -20°C for determination of triglycerides (TG), total cholesterol (TC), free fatty acids (FFA), high-density lipoprotein (HDL), and low-density lipoprotein (LDL) (Childs et al., 2008). Serum indicators were detected using the Hitachi-3110 fully automated biochemistry analyzer.

Carcass characteristics measurements and sample collection

The initial and final weight (the weight of the sheep at the end of the experiment) of the sheep were recorded to evaluate the average daily gain (ADG). The daily feed intake of each sheep was recorded to evaluate the average daily feed intake (ADFI). After slaughter, the carcass weight of each sheep was used to calculate the dressing percentage of the animals. After carcass segmentation, the characteristics of the relevant carcasses were measured, including the thickness of the back fat at the 12th rib, and the area of the *longissimus thoracis* (LT) muscle between the 12th and 13th ribs on the left side. The bones were separated from the muscle and fat tissues and weighed to calculate the net weight of meat and carcass yield.

Approximately 5 g of the LT muscle sample was placed in sterile tubes, frozen in liquid nitrogen, and stored at -80 °C for further analysis. In addition, the remaining sample of LT muscle was placed in a polyethylene sample bag and stored at -20 °C until the analysis of IMF, TG, and the composition of fatty acid.

Intramuscular fat, triglycerides, and fatty acid composition

The IMF content in the LT muscle was detected by the Soxhlet petroleum-ether extraction method based on the Chinese National Standards GB/T 5009.6.2016. The IMF contents in LT muscle were detected by the Soxhlet petroleum-ether extraction method based on Chinese National Standards GB/T 5009.6.2016. Homogenized tissue samples were prepared in physiological saline solution (tissue weight: physiological saline solution = 1:9), centrifuged for 10 minutes, and the TG content was determined using the GPO-PAP method reagent kit (A110-1-1 triglyceride assay kit, Nanjing, China).

The composition and contents of fatty acids were detected in the way described by Folch et al. (Folch et al., 1957). Samples of fatty acids were extracted from 5 g of meat using chloroform: methanol (2:1, v/v), filtered, and mixed with 20% sodium chloride, after shaking and standing, the chloroform lipid fraction was retained. After rotary evaporation, the lipid extracts were saponified and methylated. The upper solution was absorbed into the injection bottle and ready for gas chromatography analysis. Gas chromatography (Agilent/GC8860, California, USA), flame ionization detector (FID), automatic sample injection, and capillary column were used for detection. The initial temperature setting of the column oven is 120°C, kept for 5 min, then increased to 250°C at a rate of 10°C / min and maintained for 28 min. The carrier gas used is nitrogen. The fatty acid analysis was performed with a split ratio of 1:20, constant flow mode, with an average speed of 28 cm/s. The relative proportions of fatty acids were identified by comparing the peak areas with the reference standard

37-component FAME mix from Supelco Analytical. (Sigma-Aldrich, Darmstadt, Germany).

RNA extraction and real-time quantitative analysis

Total ribonucleic acid was extracted from LT muscle samples using the RNAiso Plus reagent (No. 9109, Takara, Japan). Reverse transcription was performed using the PrimeScript™ RT reagent kit (No. RR047, Takara, Japan). The Roche LightCycler 96 QPCR system and TB Green™ Premix Ex Taq™II (No. RR820A, Takara, Japan). were used for real-time PCR. The β -actin gene was used as a reference gene. The PCR data analysis program was used to quantify the Ct values, and the mRNA expressions were calculated using the $2^{-\Delta\Delta Ct}$ method. The primers for quantitative real-time PCR are shown in Table 2.

Determination of the key enzyme activity

Following the homogenization of the longissimus dorsi muscle tissue, the supernatant was collected, and the activity of p-AMPK α 2, p-ACC α , and CPT1B in the LT muscle tissue was measured using ELISA kits from Wuhan Xinqidi Biological Technology Co., Ltd (China). The experimental procedures were conducted according to the instructions provided with the kits.

Statistical analysis

Using IBM SPSS Statistics 22.0 to perform normality tests on the data, independent sample t-test analysis is used to analyze differences in group statistics. Partial least squares discriminant analysis (PLS-DA) was used to identify the main factors that affect the composition of the fatty acids. Data are presented as the mean \pm standard error of the mean (SEM). Extremely significant was defined as $p < 0.01$, significance was defined as $p < 0.05$, and trends were defined as $0.05 \leq p < 0.10$.

Results

Growth and slaughter performance

As shown in Table 3 The loin eye area (LEA) of the LT muscle in the PRO was significantly higher than that of the CON ($p < 0.05$). Compared to CON, PRO also showed an increasing trend in final weight and carcass weight, while carcass length showed a decreasing trend ($0.05 \leq p < 0.10$). There were no significant differences in other traits of growth and slaughter performance ($p > 0.05$).

Plasma lipid biochemical indicators

From Table 4 it can be concluded that the PRO showed significant decreases in plasma levels of TG, HDL, LDL, and FFA compared to the CON ($p < 0.01$). On the contrary, there were no significant differences in TC levels between CON and PRO ($p > 0.05$).

Intramuscular fat content and triglyceride content

The effect of probiotics feed on IMF and TG levels in Sunit sheep is reflected in Table 5.

It can be seen that compared to the CON, the PRO showed a significant increase in IMF content ($p < 0.05$). It should be noted that the significant trend in TG and IMF is the same ($p < 0.01$).

Fatty acid profiles

The probiotics feed to a certain extent ameliorated the fatty acid profiles in the LT of the Sunit sheep, as shown in Fig. 1 and Table 6. Compared to the CON, the proportion of MUFA and n-3 polyunsaturated fatty acids (n-3PUFAs) in the PRO increased significantly ($p < 0.05$). Further analysis found that probiotics supplementation with feed significantly decreased the proportion of capric acid (C10:0) and (C14:0) ($p < 0.05$), greatly increased the percentage of myristoleic acid (C14:1) ($p < 0.01$), and dramatically raised the proportion of oleic acid (C18:1n9 cis) ($p < 0.05$). At the same time, the effect of probiotics on the profile of PUFAs was more pronounced, with a substantial increase in the proportion of arachidonic acid (C20:4 n6) ($p < 0.01$), and a considerable increase in the percentage of γ -linolenic acid (C18:3n6), α -linolenic acid (C18:3n3), cis-8,11,14-eicosatrienoic acid (C20:3n6), cis-5,8,11,14,17-eicosapentaenoic acid (EPA, C20:5n3) and cis-15-tetracosenoic acid (C24:1n3) ($p < 0.05$), while the proportion of linoleic acid (C18:2n6) showed slowly increasing trend ($0.05 \leq p < 0.10$).

Specific description of fatty acids composition

In this study, PLS-DA was applied to elucidate the clustering and overall differences among different groups of fatty acids. There was some overlap between the CON and the PRO groups of fatty acids, but not complete consistency. The scores of the first two factors of PLS-DA are shown in (Fig. 2A). The first principal component factor (PC1) explained 35.7% of the variance, and the second principal component factor (PC2) explained 21.5% of the variance. Based on the PLS-DA loading plot shown in (Fig. 2B), C16:1, C14:1, C24:1n3, 8,11,14-eicosatrienoic acid (C20:3n6), C18:3n-3, and C18:2n6 had higher loading coefficients on PC1, indicating a strong association with PC1. At the same time, C14:1, C24:1n3, and C20:3n6 showed a positive correlation with PC1, while C16:1, C18:3n-3, and C18:2n6 showed a negative correlation with PC1. Meanwhile, PC2 had a close relationship with C18:1n9 cis, arachidic acid (C20:0), C18:3n6, lauric acid (C12:0), C14:0, and hexadecanoic acid (C16:0), among which only C18:1n9 cis showed a negative correlation with PC2, while the other indicators showed a positive correlation with PC2. It is worth noting that PC1 mainly described unsaturated fatty acids (UFAs), while PC2 was described by both saturated fatty acids (SFAs) and UFAs.

The levels of mRNA expression of genes related to lipid metabolism

The levels of mRNA expression of genes associated with fatty acid metabolism, including the synthesis and oxidation of fatty acid, are shown in Fig 3. The expression of AMPK α 2 mRNA, a key transcription factor involved in energy metabolism, was

significantly lower in the PRO compared to the CON ($p < 0.05$), while the expression of SREBP-1c mRNA, a key transcription factor involved in lipid metabolism, was significantly higher in the PRO ($p < 0.01$). Furthermore, compared to the CON, the addition of probiotics feed significantly upregulated the expression of ACC α mRNA, an intermediate metabolite that plays an important role in fatty acids synthesis and metabolism ($p < 0.01$). Conversely, the expression of CPT1B mRNA, a rate-limiting enzyme in the oxidation of fatty acids, was significantly downregulated by probiotics feed ($p < 0.01$). Similarly, the expression of FASN mRNA, a fatty acid synthesis enzyme, was significantly upregulated in the PRO ($p < 0.05$). Moreover, the expression of SCD1 mRNA, the enzyme responsible for desaturating SFAs into MUFAs, was also significantly upregulated in the PRO ($p < 0.01$).

The activity of key enzymes involved in lipid metabolism

The impact of probiotics feed on the key molecular activity of enzymes related to lipid metabolism in Sunit sheep is illustrated in Fig 4. Based on the figure, it can be observed that the activity of p-AMPK α 2 and CPT1B in the PRO was significantly reduced compared to the CON ($p < 0.05$), whereas the activities of p-ACC α , SREBP-1c, FASN, and SCD1 were significantly increased ($p < 0.01$). These findings demonstrate a similar trend to the mRNA expression results mentioned earlier.

Correlation between lipid composition parameters and gene expression

Pearson's correlation analysis was conducted to determine the correlation between lipid composition parameters and gene expression. As shown in Table 7, AMPK α 2 showed significant negative correlations with IMF, C14:1, C18:3 n6, C20:3n6, and C20:4n6, while ACC α exhibited significant positive correlations with C14:1, C18:2 n6, C18:3n6, C20:3n6, C20:4n6, and C24:1n3. CPT1B showed significant negative correlations with C14:1, C18:2n6, C18:3n6, C20:3n6, C20:4n6, and C24:1n3. Additionally, SREBP-1c exhibited significant positive correlations with C14:1, C20:3 n6, and C20:4n6. As predicted, FASN showed a strong positive correlation with IMF, while SCD1 demonstrated a significant negative correlation with C10:0, and significant positive correlations with C18:3n6, C20:3n6, and C20:4n6.

Discussion

The nutritional and dietary value of sheep is largely determined by its feed, which also affects consumer health by influencing the IMF content and the composition of fatty acids. Probiotics, as beneficial feed additives, are widely used in livestock production. Therefore, it is necessary to evaluate the effects of probiotics feed on growth performance, carcass characteristics, plasma biochemical parameters, and muscle lipid indicators, including IMF and TG content, fatty acid composition, and the expression of the lipid-related genes and the activity of the enzyme of sheep, in order to provide reasonable recommendations for intensive livestock production.

The present study revealed that probiotics supplementation in Sunit sheep resulted in an increasing trend in final weight and carcass weight, along with a significant improvement in the LEA, which is consistent with previous research findings (Lu et al., 2021). LEA, as a growth indicator, is highly positively correlated with meat production performance and lean meat percentage and is an important indicator for evaluating the carcass quality of domestic animals. The larger area of the loin eye and the potential higher fat content in the LT may positively influence the taste of the meat. It is noteworthy that the results of this study confirm the higher IMF content in the LT of Sunit sheep in the PRO, supporting the aforementioned interpretation. Similar observations have been made by Nie et al. and Ramzi et al., who reported that adding probiotics to the diet can increase IMF levels in mutton and alter the overall fat distribution in sheep, which is in line with our research results (Amran et al., 2021; Nie et al., 2022). The most plausible explanation for these findings is that probiotics modulate rumen fermentation patterns and alter indicators of lipid metabolism, including the composition of fatty acids, further promoting the synthesis and distribution of IMF. Therefore, it was necessary to explore the differences in the plasma biochemical indicators of sheep influenced by probiotics.

The results of this study on the effects of probiotics feed on plasma lipid biochemical indicators in Sunit sheep are consistent with the findings of Abas, Chiofalo, and Baiomy, who reported that probiotics supplementation can decrease the concentrations of plasma TG and LDL in lambs (Chiofalo et al., 2004; Abas et al., 2007; Baiomy, 2010). The reduction of plasma TG may be attributed to the improved

digestive and absorptive functions of ruminant animals by probiotics feed, which can reduce the absorption and synthesis of fats in the intestine, thereby lowering the synthesis and accumulation of plasma TG (Sheikh, 2019). The current research findings on the impact of probiotics on cholesterol metabolism in sheep plasma are inconclusive. The research conducted by Mahesh and Jia suggests that probiotics can influence cholesterol metabolism through various metabolic pathways, such as reducing cholesterol absorption and inhibiting the activity of cholesterol synthesis enzymes (Mahesh et al., 2021; Jia et al., 2022). This leads to a decrease in the levels of LDL in the plasma, which has a positive effect on plasma cholesterol metabolism. In this study, the addition of probiotics didn't significantly alter the levels of TC in the serum of Sunit sheep, but it did reduce the concentration of LDL in the plasma. This suggests that different types of probiotics may have varying effects on cholesterol metabolism in sheep. Therefore, more research is needed to elucidate the specific mechanisms by which probiotics regulate cholesterol lipoprotein levels in sheep plasma. In this study, the PRO showed a reduction in the levels of free fatty acid in the plasma of Sunit sheep. Probiotics were able to decrease the rate of release of free fatty acid, regulate fatty acid metabolism in plasma, and ultimately affect fat metabolism in muscle tissue. The decrease in free fatty acid levels in plasma may lead to a reduction in fatty acid oxidation in muscle tissue. Fatty acid oxidation is the process by which fatty acids are converted into energy within cells. When fatty acid oxidation is reduced, muscle tissue can tend to store fatty acids as fat. (Domaradzki et al., 2019; Roque-Jiménez et al., 2021). This could be a possible reason for the

increase in IMF content in the muscle of feed with probiotics, as observed by Mackey, who found that supplementation of probiotics components to cattle in feed resulted in an increased marbling score in LT, indicating that probiotics may enhance IMF deposition in cattle (Mackey et al., 2023).

In meat, IMF is abundant in fatty acids, which can be absorbed by the human body and enter the bloodstream for transportation to various tissues. Changes in IMF content and TG levels in meat lead to corresponding alterations in the relative content of fatty acids. Increasing the IMF content is associated with a decrease in the concentration of PUFAs in meat, while the concentration of SFAs and MUFAs increases (Purchas et al., 2005; Lukic et al., 2019; Razmaitè et al., 2020; Realini et al., 2021). Joo suggests that the C18:1n9 cis levels in beef increase with the marbling pattern, and this elevation of C18:1n9 cis content is beneficial to the human body (Joo et al., 2017). Kazala indicates that the relative content of C14:0 in beef cattle increases with the increase in total lipids and TG. Meanwhile, the relative content of C18:3 in beef is negatively correlated with the total lipid content (Kazala et al., 1999). In this study, the model of PLS-DA was constructed. PC1 was defined as describing UFAs, while PC2 was defined as being able to simultaneously describe both SFAs and UFAs. The variable loadings plot showed that the fatty acids that contribute significantly to PC1 and PC2 were positively correlated in their relative positions in the two-dimensional space, mainly concentrated along the positive axes of PC1 and PC2.

As stated above, the addition of probiotics to the feed has been found to increase the IMF content in Sunit sheep. Furthermore, it has been observed that the use of probiotic feed has led to an increasing trend in the relative proportion of UFAs in the LT of Sunit sheep. Specifically, the proportions of C14:1 and C18:1 n9 cis were significantly increased in the PRO. Among them, C18:1 n9 cis is the most abundant UFA, which is considered to be positively correlated with the tenderness of fat and the sensory quality of meat. The higher C18:1 n9 cis content can enhance the palatability of the meat (Tang et al., 2021). Furthermore, probiotics can modulate the synthesis and oxidation of MUFAs by regulating metabolic enzymes such as fatty acyl-CoA synthetase and enzymes limiting the rate of fatty acid oxidation, thereby altering the composition of UFAs in meat (Gheorghe et al., 2022). In accordance with the findings mentioned earlier, an increase in the relative proportion of MUFAs may result in a decrease in LDL levels in the body. In this study, the relative proportions of C18:3n6, C18:3n3, C20:3n6, C20:4n6, C20:5n3, C24:1n-3, and n-3 PUFA were found to be higher in the LT of Sunit sheep in the PRO. Among them, C18:3n6 and C18:3n3 are important substrates for the synthesis of long-chain polyunsaturated fatty acids of n6 and n3 origin (Ganesh et al., 2016). C20:4, as an important essential fatty acid, plays a role in changing cell membrane stability and participating in cell-mediated lipid metabolism (Calder, 2008). The content of C20:4 is related to the activity of fatty acid desaturase. The addition of probiotics to the diet benefits the activity of fatty acid desaturase. Generally speaking, higher expression levels of fatty acid desaturase are associated with a higher synthesis rate and content of C20:4 (Dev

et al., 2021). The composition and relative proportion of fatty acids in meat are key factors that influence its nutritional value. This study suggests that the addition of probiotics to the diet improves the proportion of PUFAs in the LT of Sunit sheep, indicating that probiotics may have a role in improving the nutritional value of fatty acids in meat.

In order to further elucidate the mechanisms underlying intramuscular fat (IMF) accumulation and fatty acid metabolism, we conducted a comprehensive analysis and assessment of gene expression levels and corresponding enzyme activities involved in the regulation of muscle fat deposition and fatty acid metabolic pathways. As widely acknowledged, the signaling of AMPK serves as a critical sensor of cellular energy, capable of participating in and regulating lipid homeostasis and fatty acid metabolism in muscle, including β -oxidation of fatty acid, synthesis of de novo fatty acid, and transport of fatty acid (Wang et al., 2013; Kim et al., 2014). According to research, AMPK is believed to play a crucial role in the regulation of lipid metabolism by modulating downstream enzymes, ACC α and CPT1B, which are involved in the fatty acid β -oxidation pathways (Ronnett et al., 2006). ACC α , as a central regulator of fatty acid metabolism, catalyzes the production of malonyl-CoA, an inhibitor of CPT1B, thus modulating CPT1B activity and regulating β -oxidation of fatty acid, therefore, the AMPK α 2-ACC α -CPT1B signaling pathway plays a crucial role in fat deposition and fatty acid metabolism (Kim et al., 2012). The results of this study revealed that supplementation with probiotic feed inhibited the expression of AMPK α 2 mRNA and CPT1 mRNA, while promoting ACC α mRNA expression in the LT of Sunit sheep.

This suggests that probiotics feed may inhibit the expression of AMPK α 2 signaling factor in Sunit sheep, impeding the AMPK α 2-ACC α pathway, inhibiting CPT1B mRNA, and weakening the β -oxidation process of fatty acids, leading to increased fatty acid synthesis. The precise mechanism underlying the modulation of CPT1B mRNA by probiotics and its ultimate regulation of fat deposition in Sunit sheep remains unclear. Some studies have proposed that lactate generated as a result of probiotic administration can promote the generation of acetyl-CoA, which directly inhibits the expression of CPT1B mRNA. Subsequent reactions in this pathway ultimately lead to intracellular lipid accumulation (Araújo et al., 2020; Brooks, 2020).

In addition, the transcription factor SREBP-1c is a key regulatory factor in the synthesis of fatty acids, capable of activating the expression of FASN mRNA, ACC α mRNA, and SCD1 mRNA, while its expression is also controlled by AMPK. Consistent with ACC α , the activity of SREBP-1c is negatively regulated by AMPK. Research has shown that AMPK inhibits the expression of SREBP-1c mRNA, thereby reducing the expression of FASN mRNA and subsequently inhibiting lipogenesis (Deng et al., 2015; An et al., 2019). SCD1, a crucial factor expressed in adipose tissue, is a target gene for SREBP-1c and plays a selective role in the regulation of de novo lipogenesis and fatty acid conversion, mediating adipogenesis and fatty acid metabolism. Deficiency of SCD in sheep is associated with decreased TG levels, as well as reduced expression of SREBP-1c mRNA and FASN mRNA (Tian et al., 2018; Liu et al., 2020; He et al., 2021). In this study, probiotics were found to inhibit the expression of AMPK α 2 mRNA while promoting the expression of SREBP-1c mRNA

and FASN mRNA, which may result in a tendency to fat synthesis in LT, leading to IMF deposition. Furthermore, probiotics significantly increased SCD mRNA expression, indicating that probiotics have the ability to gradually convert SFAs to UFAs, which may be one of the reasons why the probiotic group had a relatively higher proportion of unsaturated fatty acids as mentioned earlier (Tian et al., 2021). Further research unveiled a similarity between the enzyme activity outcomes and the mRNA expression level findings. The results indicate that probiotics feed can regulate the ACC α , CPT1B, SREBP-1c, FASN, and SCD1 by inhibiting the AMPK α 2, resulting in enhanced IMF synthesis and reduced oxidation of fatty acids in the LT of Sunit sheep. This ultimately leads to an increase in IMF content and improved fatty acid composition in meat.

The correlation analysis between gene expression and IMF, TG, and fatty acid composition has been used to investigate the relationship between gene signaling pathways and lipid composition parameters. Even minor changes in chemical structure can significantly alter lipid signaling activity, so research on gene-regulated fatty acids composition is limited (Tánori-Lozano et al., 2022). The AMPK signaling factor can regulate the cellular uptake of fatty acids through translocation of fatty acids. Studies suggest that inhibiting AMPK α 2 may increase key enzyme activities in the fatty acid synthesis pathway, leading to an increase in PUFAs synthesis, while simultaneously reducing cellular fatty acid β -oxidation, resulting in the accumulation of fatty acids (Dzamko et al., 2008; Kohan et al., 2009). In addition, downstream genes of the AMPK signaling pathway, including ACC α , CPT1B, SREBP-1c, FASN,

and SCD1, are closely associated with the composition of fatty acids. Studies suggest that elevated expression of ACC α mRNA promotes the deposition of UFAs, and the accumulation of UFAs inhibits the expression of CPT1B mRNA (Zhu et al., 2022). Meanwhile, the main function of SREBP-1c is to promote the synthesis of fatty acids and triglycerides, exerting a positive regulatory effect on adipose deposition and fatty acid metabolism (Guo et al., 2023). In this study, the expression levels of AMPK α 2 mRNA and CPT1B mRNA were negatively correlated with IMF, C18:3 n6, C20:3 n6, and C20:4 n6, while the expression levels of ACC α mRNA were positively correlated with their contents. Furthermore, it was further confirmed that inhibition of the AMPK α 2 signaling pathway could promote IMF deposition in Sunit sheep by enhancing ACC α and SREBP-1c factors and improving the composition of fatty acids (Fig 5).

Conclusion

In conclusion, the results of this study indicate that the addition of probiotics to the diet can regulate fat deposition and improve fatty acid composition in the LT of Sunit sheep through the signaling pathways AMPK-ACC-CPT1 and AMPK-SREBP-1c. This regulatory mechanism leads to an increase in the loin eye area in Sunit sheep, improved growth performance, and altered plasma lipid biochemical indicators. The findings of this experiment highlight the regulatory effects of probiotics added to the diet on fat deposition and fatty acid composition in the LT of Sunit sheep and provide new insights into the nutritional interactions of ruminant animals mediated by the

AMPK pathway. The research results contribute to a better understanding of the potential application of probiotics and provide new insights into the future of food science and animal resources.

Conflicts of interest

The authors declare no potential conflicts of interest.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (32160589); the Major Science and Technology Projects of Ordos (2022EEDSKJZDZX029); the Major R&D and Achievements Transformation Projects of Inner Mongolia (2022YFXZ0017); the Natural Science Foundation of Inner Mongolia (2020ZD11); the Natural Science Foundation of Inner Mongolia (2022MS03055); the Natural Science Foundation of Inner Mongolia (2021MS03010) and the Scientific Research Projects of Universities Directly under Inner Mongolia (BR22-13-08).

Author Contributions

Conceptualization: Ye Jin and Yue Zhang ; Data curation: Yue Zhang and Duo Yao; Formal analysis: Huan Huang and Min Zhang; Methodology: Yue Zhang, Lin Su, and Lina Sun; Software: Yue Zhang; Validation: LiHua Zhao; Ye Jin, and Yueying Guo; Investigation: Yue Zhang, Huan Huang, and Duo Yao; Writing -

original draft: Yue Zhang; Writing - review & editing: Yue Zhang, Ye Jin, and Yueying Guo.

Ethics Approval

The study was conducted in accordance with the Declaration of Helsinki, and all animals handling procedures were performed in accordance with the Ethics Committee of Inner Mongolia Agricultural University (permit number: NND2021072).

References

- Abas, I., H. C. Kutay, R. Kahraman, N. Y. Toker, D. Ozcelik, F. Ates, and A. Kaçakci. 2007. Effects of organic acid and bacterial direct-fed microbial on fattening performance of Kivircik-Male yearling lambs. *Pak J Nutr* 6:149-154.
- Abdullah, F. A. A., G. Borilova, and I. Steinhäuserova. 2019. Halal criteria versus conventional slaughter technology. *Animals* 9(8):530.
- Amran, R. A., A. R. Alhimaidi, M. Q. Al-Ghadi, A. A. Ammari, and A.-R. Z. J. O. C. Gaafar. 2021. Carryover effect of direct-fed microbial supplementation and early weaning on the growth performance and carcass characteristics of growing Najdi lambs. *Open Chemistry* 19(1):1265-1278.
- An, J.-P., J. H. Choi, J. Huh, H. J. Lee, S. Han, J.-R. Noh, Y.-H. Kim, C.-H. Lee, and W.-K. Oh. 2019. Anti-hepatic steatosis activity of *Sicyos angulatus* extract in high-fat diet-fed mice and chemical profiling study using UHPLC-qTOF-MS/MS spectrometry. *Phytomedicine* 63:152999.
- Araújo, J. R., A. Tazi, O. Burlen-Defranoux, S. Vichier-Guerre, G. Nigro, H. Licandro, S. Demignot, P. Sansonetti, and microbe. 2020. Fermentation products of commensal bacteria alter enterocyte lipid metabolism. *Cell host* 27(3):358-375.
- Ayala-Monter, M. A., D. Hernández-Sánchez, S. González-Muoz, R. Pinto-Ruiz, J. A. Martínez-Aispuro, N. Torres-Salado, J. Herrera-Pérez, and A. Gloria-Trujillo. 2019. Growth performance and health of nursing lambs supplemented with inulin and *Lactobacillus casei*. *Asian-Australasian Journal of Animal Sciences* 32(8):1137-1144.

- Baiomy, A. 2010. Influence of live yeast culture on milk production, composition and some blood metabolites of Ossimi ewes during the milking period. *Journal of Animal and Poultry Production* 1(10):469-480.
- Bao, Y., Y. Zhang, H. Li, Y. Liu, S. Wang, X. Dong, F. Su, G. Yao, T. Sun, and H. Zhang. 2012. In vitro screen of *Lactobacillus plantarum* as probiotic bacteria and their fermented characteristics in soymilk. *Annals of Microbiology* 62(3):1311-1320.
- Bessa, R. J., S. P. Alves, and J. Santos-Silva. 2015. Constraints and potentials for the nutritional modulation of the fatty acid composition of ruminant meat. *European Journal of Lipid Science and Technology* 117(9):1325-1344.
- Brooks, G. A. 2020. Lactate as a fulcrum of metabolism. *Redox biology* 35:101454.
- Calder, P. C. 2008. The relationship between the fatty acid composition of immune cells and their function. *Prostaglandins, Leukotrienes Essential Fatty Acids* 79(3-5):101-108.
- Chikwanha, O. C., P. Vahmani, V. Muchenje, M. E. Dugan, and C. Mapiye. 2018. Nutritional enhancement of sheep meat fatty acid profile for human health and wellbeing. *Food Research International* 104:25-38.
- Childs, S., A. Hennessy, J. Sreenan, D. Wathes, Z. Cheng, C. Stanton, M. Diskin, and D. Kenny. 2008. Effect of level of dietary n-3 polyunsaturated fatty acid supplementation on systemic and tissue fatty acid concentrations and on selected reproductive variables in cattle. *Theriogenology* 70(4):595-611.
- Chiofalo, V., L. Liotta, and B. Chiofalo. 2004. Effects of the administration of *Lactobacilli* on body growth and on the metabolic profile in growing Maltese goat kids. *Reproduction Nutrition Development* 44(5):449-457.
- Deng, X., Q. Dong, D. Bridges, R. Raghov, E. A. Park, and M. B. Elam. 2015. Docosahexaenoic acid inhibits proteolytic processing of sterol regulatory element-binding protein-1c (SREBP-1c) via activation of AMP-activated kinase. *Biochimica et Biophysica Acta -Molecular Cell Biology of Lipids* 1851(12):1521-1529.
- Dev, K., J. Begum, A. Biswas, J. Kannoujia, N. A. Mir, J. Sonowal, R. Kant, and T. Narender. 2021. Dietary *Lactobacillus acidophilus* and mannan-oligosaccharides alter the lipid metabolism and health indices in broiler chickens. *Probiotics Antimicrobial Proteins* 13:633-646.
- Domaradzki, P., M. Florek, P. Skąlecki, A. Litwińczuk, M. Kędzierska-Matysek, A. Wolanciuk, and K. Tajchman. 2019. Fatty acid composition, cholesterol content and lipid oxidation indices of intramuscular fat from skeletal muscles of beaver (*Castor fiber L.*). *Meat science* 150:131-140.
- Dzamko, N., J. D. Schertzer, J. G. Ryall, R. Steel, S. L. Macaulay, S. Wee, Z. P. Chen, B. J. Michell, J. S. Oakhill, and M. J. Watt. 2008. AMPK-independent pathways regulate skeletal muscle fatty acid oxidation. *The Journal of physiology* 586(23):5819-5831.
- Fang, K., F. Wu, G. Chen, H. Dong, J. Li, Y. Zhao, L. Xu, X. Zou, and F. Lu. 2019. Diosgenin ameliorates palmitic acid-induced lipid accumulation via AMPK/ACC/CPT-1A and SREBP-1c/FAS signaling pathways in LO2 cells. *BMC complementary and alternative medicine* 19(225):1-12.

- Ferré, P., F. Phan, and F. Foufelle. 2021. SREBP-1c and lipogenesis in the liver: An update. *Biochemical Journal* 478(20):3723-3739.
- Folch, J., M. Lees, and G. H. Sloane Stanley. 1957. A simple method for the isolation and purification of total lipids from animal tissues. *JOURNAL OF BIOLOGICAL CHEMISTRY* 226(1):497-509.
- Ganesh, V., N. S. Hettiarachchy, and nutrition. 2016. A review: supplementation of Foods with essential fatty acids—can it Turn a Breeze without further Ado? *Critical reviews in food science* 56(9):1417-1427.
- Gheorghe, A., M. Habeanu, G. Ciurescu, N. A. Lefter, M. Ropota, I. Custura, and M. Tudorache. 2022. Effects of dietary mixture enriched in polyunsaturated fatty acids and probiotic on performance, biochemical response, breast meat fatty acids, and lipid indices in broiler chickens. *Agriculture* 12(8):1120.
- Guo, R., L. Chen, J. Zhu, J. Li, Q. Ding, K. Chang, Q. Han, and S. Li. 2023. Monounsaturated fatty acid-enriched olive oil exacerbates chronic alcohol-induced hepatic steatosis and liver injury in C57BL/6J mice. *Food Function* 14(3):1573-1583.
- He, L., Y. Qin, Y. Wang, D. Li, W. Chen, and J. Ye. 2021. Effects of dietary replacement of fish oil with soybean oil on the growth performance, plasma components, fatty acid composition and lipid metabolism of groupers *Epinephelus coioides*. *Aquaculture Nutrition* 27(5):1494-1511.
- Jia, B., Y. Zou, X. Han, J.-W. Bae, and C. O. Jeon. 2022. Gut microbiome-mediated mechanisms for reducing cholesterol levels: implications for ameliorating cardiovascular disease. *Trends in Microbiology* 31(1):76-91.
- Joo, S.-T., Y.-H. Hwang, and D. Frank. 2017. Characteristics of Hanwoo cattle and health implications of consuming highly marbled Hanwoo beef. *Meat science* 132:45-51.
- Juárez, M., S. Lam, B. M. Bohrer, M. E. Dugan, P. Vahmani, J. Aalhus, A. Juárez, O. López-Campos, N. Prieto, and J. Segura. 2021. Enhancing the nutritional value of red meat through genetic and feeding strategies. *Foods* 10(4):872.
- Kazala, E. C., F. J. Lozeman, P. S. Mir, A. Laroche, D. R. Bailey, and R. Weselake. 1999. Relationship of fatty acid composition to intramuscular fat content in beef from crossbred Wagyu cattle. *Journal of Animal Science* 77(7):1717-1725.
- Ke, R., Q. Xu, C. Li, L. Luo, and D. Huang. 2018. Mechanisms of AMPK in the maintenance of ATP balance during energy metabolism. *Cell Biology International* 42(4):384-392.
- Kim, J., S. Ingale, S. Lee, Y. Choi, E. Kim, D. Lee, Y. Kim, and B. Chae. 2014. Impact of dietary fat sources and feeding level on adipose tissue fatty acids composition and lipid metabolism related gene expression in finisher pigs. *Animal Feed Science Technology* 196:60-67.
- Kim, M. K., S. H. Kim, H. S. Yu, H. G. Park, U. G. Kang, Y. M. Ahn, and Y. S. Kim. 2012. The effect of clozapine on the AMPK-ACC-CPT1 pathway in the rat frontal cortex. *International Journal of Neuropsychopharmacology* 15(7):907-917.

- Kohan, A. B., I. Talukdar, C. M. Walsh, and L. M. Salati. 2009. A role for AMPK in the inhibition of glucose-6-phosphate dehydrogenase by polyunsaturated fatty acids. *Biochemical biophysical research communications* 388(1):117-121.
- Laliotis, G., I. Bizelis, and E. Rogdakis. 2010. Comparative approach of the de novo fatty acid synthesis (lipogenesis) between ruminant and non ruminant mammalian species: from biochemical level to the main regulatory lipogenic genes. *Current genomics* 11(3):168-183.
- Li, H., F. Liu, J. Lu, J. Shi, and G. Huo. 2020. Probiotic Mixture of *Lactobacillus plantarum* Strains Improves Lipid Metabolism and Gut Microbiota Structure in High Fat Diet-Fed Mice. *Frontiers in Microbiology* 11:512.
- Liu, L., Y. Wang, X. Liang, X. Wu, J. Liu, S. Yang, C. Tao, J. Zhang, J. Tian, and J. Zhao. 2020. Stearoyl-CoA desaturase is essential for porcine adipocyte differentiation. *International Journal of Molecular Sciences* 21(7):2446.
- Lu, J., Z. Chen, Q. Gao, P. Li, J. Wang, Y. Cai, Z. Wang, D. Li, H. Li, and F. Wang. 2021. Combined supplementation of probiotics and enzymes improves performance and regulates rumen microbiota in fattening goats. *Research Square* 1:1-26.
- Lukic, M., D. Trbovic, R. Petronijevic, V. Djordjevic, D. Karan, J. B. Milijasevic, and Z. Petrovic. 2019. Chemical composition and fatty acid profile of the longissimus dorsi muscle in Simmental bulls. *IOP Conference Series: Earth and Environmental Science* 333:012075.
- Mackey, S., R. Cooke, E. Colombo, A. Pickett, L. Batista, E. Block, and A. Brandão. 2023. Supplementing pre-and probiotic ingredients to feedlot steers: effects on health, growth performance, and physiological responses. *animal*:100770.
- Mahesh, M., R. K. Mohanta, and A. K. Patra. 2021. Probiotics in livestock and poultry nutrition and health. *Advances in Probiotics for Sustainable Food* 21:149-179.
- Nie, C., Y. Hu, R. Chen, B. Guo, L. Li, H. Chen, H. Chen, and X. Song. 2022. Effect of probiotics and Chinese medicine polysaccharides on meat quality, muscle fibre type and intramuscular fat deposition in lambs. *Italian Journal of Animal Science* 21(1):811-820.
- Pećina, M., Z. Luković, D. Škorput, M. Bolčić, K. Salajpal, and D. Karolyi. 2018. Relationship between intramuscular fat content and fatty acid composition of pork. *Journal of Central European Agriculture* 19(4):840-845.
- Purchas, R., T. Knight, and J. Busboom. 2005. The effect of production system and age on concentrations of fatty acids in intramuscular fat of the longissimus and triceps brachii muscles of Angus-cross heifers. *Meat Science* 70(4):597-603.
- Rabelo, C. H. S., F. C. Basso, T. A. Mcallister, J. F. Lage, G. S. Goncalves, E. C. Lara, A. A. Oliveira, T. T. Berchielli, and R. A. Reis. 2016. Influence of *Lactobacillus buchneri* as silage additive and forage:concentrate ratio on the growth performance, fatty acid profile in longissimus muscle, and meat quality of beef cattle. *Canadian Journal of Animal Science* 96(4):1-13.
- Razmaitė, V., A. Šiukščius, R. Šveistienė, S. Bliznikas, and V. Jatkauskienė. 2020. Relationships between fat and cholesterol contents and fatty acid composition in different meat-producing animal species. *Acta Veterinaria* 70(3):374-385.

- Realini, C., E. Pavan, R. Purchas, M. Agnew, P. Johnson, E. Bermingham, and C. Moon. 2021. Relationships between intramuscular fat percentage and fatty acid composition in *M. longissimus lumborum* of pasture-finished lambs in New Zealand. *Meat Science* 181:108618.
- Ronnett, G. V., A. M. Kleman, E. K. Kim, L. E. Landree, and Y. Tu. 2006. Fatty acid metabolism, the central nervous system, and feeding. *Obesity* 14(S8):201S-207S.
- Roque-Jiménez, J. A., M. Rosa-Velázquez, J. M. Pinos-Rodríguez, J. G. Vicente-Martínez, G. Mendoza-Cervantes, A. Flores-Primo, H. A. Lee-Rangel, and A. E. Relling. 2021. Role of long chain fatty acids in developmental programming in ruminants. *Animals* 11(3):762.
- Sheikh, G. G. M., Danish Ganai, A. M. Muzamil, S. Afzal, Y. Ahmad, H. A. . 2019. Effect of probiotics mix supplementation on haemato-biochemical parameters and bacterial faecal shedding in Corriedale lambs fed paddy straw based complete feed. *Indian Journal of Animal Research* 53(8):1049-1053.
- Taboada, N., M. F. Salom, A. Cordoba, S. N. González, S. L. Alzogaray, and C. Van Nieuwenhove. 2022. Administration of selected probiotic mixture improves body weight gain and meat fatty acid composition of creole goats. *Food Bioscience* 49:101836.
- Tang, X., X. Liu, and H. Liu. 2021. Effects of dietary probiotic (*Bacillus subtilis*) supplementation on carcass traits, meat quality, amino acid, and fatty acid profile of broiler chickens. *Frontiers in Veterinary Science* 8:767802.
- Tánori-Lozano, A., A. I. Quintana-Romandía, M. Montalvo-Corral, A. Pinelli-Saavedra, M. Valenzuela-Melendres, J. L. Dávila-Ramírez, T. Y. Islava-Lagarda, and H. González-Ríos. 2022. Influence of ferulic acid and clinoptilolite supplementation on growth performance, carcass, meat quality, and fatty acid profile of finished lambs. *Journal of Animal Science Technology* 64(2):274-290.
- Thomson, D. M., and W. W. Winder. 2009. AMP-activated protein kinase control of fat metabolism in skeletal muscle. *Acta physiologica* 196(1):147-154.
- Tian, H., J. Luo, Z. Zhang, J. Wu, T. Zhang, S. Busato, L. Huang, N. Song, and M. Bionaz. 2018. CRISPR/Cas9-mediated stearoyl-CoA desaturase 1 (SCD1) deficiency affects fatty acid metabolism in goat mammary epithelial cells. *Journal of agricultural food chemistry* 66(38):10041-10052.
- Tian, Z., Y. Cui, H. Lu, G. Wang, and X. Ma. 2021. Effect of long-term dietary probiotic *Lactobacillus reuteri* 1 or antibiotics on meat quality, muscular amino acids and fatty acids in pigs. *Meat Science* 171:108234.
- Tong, X., P. Li, D. Zhang, K. VanDommelen, N. Gupta, L. Rui, M. B. Omary, and L. Yin. 2016. E4BP4 is an insulin-induced stabilizer of nuclear SREBP-1c and promotes SREBP-1c-mediated lipogenesis [S]. *Journal of lipid research* 57(7):1219-1230.
- Toral, P. G., F. J. Monahan, G. Hervás, P. Frutos, and A. P. Moloney. 2018. Review: Modulating ruminal lipid metabolism to improve the fatty acid composition of meat and milk. Challenges and opportunities. *animal* 12(s2):1-10.

- Wang, H., J. Zhong, C. Zhang, Z. Chai, H. Cao, J. Wang, J. Zhu, J. Wang, and Q. Ji. 2020. The whole-transcriptome landscape of muscle and adipose tissues reveals the ceRNA regulation network related to intramuscular fat deposition in yak. *BMC Genomics* 21(1):1-15.
- Wang, J., H. Sun, S. Guo, Y. Sun, L. Y. Kwok, H. Zhang, and C. Peng. 2021. Comparison of the effects of single probiotic strains *Lactobacillus casei* Zhang and *Bifidobacterium animalis* ssp. *lactis* Probio-M8 and their combination on volatile and nonvolatile metabolomic profiles of yogurt. *Journal of Dairy Science* 104(7):7509-7521.
- Wang, W., W. Xue, B. Jin, X. Zhang, F. Ma, and X. Xu. 2013. Candidate gene expression affects intramuscular fat content and fatty acid composition in pigs. *Journal of applied genetics* 54:113-118.
- Yang, C., J. Liu, X. Wu, P. Bao, R. Long, X. Guo, X. Ding, and P. Yan. 2017. The response of gene expression associated with lipid metabolism, fat deposition and fatty acid profile in the longissimus dorsi muscle of Gannan yaks to different energy levels of diets. *PLoS one* 12(11):e0187604.
- Zhang, Z., X. Liang, L. Tong, Y. Lv, H. Yi, P. Gong, X. Tian, Q. Cui, T. Liu, and L. Zhang. 2021. Effect of *Inonotus obliquus* (Fr.) Pilat extract on the regulation of glycolipid metabolism via PI3K/Akt and AMPK/ACC pathways in mice. *Journal of ethnopharmacology*, 273:113963.
- Zhu, T., Y. Shen, X. Li, T. Pan, J. Luo, J. Lu, Y. Bao, Z. Wu, L. Jiao, and D. R. Tocher. 2022. Effects of an alternating linseed oil-fish oil feeding strategy on growth, fatty acid restoration and expression of lipid related genes in black seabream (*A. schlegelii*). *Aquaculture* 547:737456.

Table 1. Ingredients and chemical composition of the base diet (% , as-fed basis)

Items	Diet ¹	
	CON	POR
Ingredient composition		
Corn	60.29	58.63
Sunflower meal	17.86	18.42
Soybean meal	13.45	13.40
Rapeseed	4.65	4.80
Limestone powder	1.00	1.00
Calcium phosphate	0.50	0.50
Probiotic	0.00	1.00
Vitamin and mineral premix ²	1.50	1.50
NaCl	0.75	0.75
Total	100.00	100.00
Chemical composition		
Crude protein	17.86	17.85
Crude fat	3.66	3.65
Crude fiber	10.60	10.54
Crude ash	7.80	7.79
Calcium	1.21	1.23
Total Phosphorus	0.63	0.65
Lysine	0.35	0.33

¹ CON, basic diet; POR, basic diet+1% probiotic.

² Vitamin and mineral premix provided the following per kg of diet:20,000 IU of vitamin A; 4,000 IU of vitamin D₃; 80 IU of vitamin E; 16 mg of vitamin K; 2.5 mg

of vitamin B₁; 2.5 mg of vitamin B₂; 8 mg of vitamin B₆; 3.4 mg of vitamin B₁₂; 54 mg of Ca-pantothenate; 25 mg of Cu ; 36 mg of Zn ; 58 mg of Mn; 0.5 mg of I ; 0.3 mg of Co; 0.4 mg of Se.

ACCEPTED

Table 2. Primer sequence for cloning and quantitative RNA assay

Gene Name ¹	GenBank	Primer sequence (5'–3')	Product size (bp)
β -actin	NM_001009784.3	F:CTCACGGAGCGTGGCTACA R:GCCATCTCCTGCTCGAGGTC	107
AMPK α 2	XM_042245133.1	F:ATGAGGTGGTGGAGCAGAGAAG R:CGTGAGAGAGCCAGAGAGTGAA	131
ACC α	XM_027973952.1	F:GACACATCACATCCGTCCTCT R:GTCCATCACACAGCCTTCAT	188
CPT1B	XM_027965744.1	F:TGTTCAACACCACTCGCATC R:CTCGTAGAGCCACAGCTTGA	116
SREBP-1c	XM_027974786.1	F:CGCAAAGCCATCGACTACATC R:TGAGCTTCTGGTTGCTGTGCT	52
FASN	XM_027974304.1	F:GTGTGGTACAGCCCCTCAAG R:ACGCACCTGAATGACCACTT	110
SCD1	NM_001009254.1	F:GAGTACCGCTGGCACATCAA R:CTAAGACGGCAGCCTTGGAT	103

¹ β -actin, actin beta; AMPK α 2, adenosine 5'-monophosphate (AMP)-activated protein kinase alpha 2; ACC α , acetyl-CoA carboxylase alpha; CPT1B, carnitine palmitoyltransferase 1B; SREBP-1c, sterol regulatory element binding protein 1; FASN, fatty acid synthase; SCD1, stearoyl-CoA desaturase.

Table 3. Effects of probiotic feed on growth and slaughter performance of Sunit sheep

Item ¹	Treatment ²		SEM	p-value
	CON	POR		
Initial weight, kg	17.65	18.12	1.88	0.814
Final weight, kg	31.77	33.33	2.05	0.082
ADG, kg/d	0.16	0.15	0.01	0.715
ADFI, kg/d	1.64	1.59	0.06	0.860
Carcass weight, kg	14.73	15.40	1.60	0.084
Dressing percentage, %	45.95	46.58	0.01	0.287
Carcass length, cm	73.80	70.00	1.73	0.065
Net meat weight, kg	12.04	12.51	1.16	0.665
Carcass yield, %	80.62	81.20	0.78	0.499
Back fat thickness, mm	4.93	4.03	0.58	0.173
Loin eye area, cm ²	14.00 ^a	16.25 ^b	0.85	0.038

¹ ADG, average daily gain; ADFI, average daily feed intake.

² CON, basic diet; POR, basic diet +1% probiotic.

a, b within a row, values with different superscripts differ significantly at $p < 0.05$ or $p < 0.01$. Data are expressed as means \pm SEM.

Table 4. Effects of probiotic feed on plasma lipid biochemical indicators of Sunit sheep

Item ¹	Treatment ²		SEM	p-value
	CON	POR		
TG, mmol/L	3.59 ^a	1.72 ^b	0.08	<0.001
TC, mmol/L	7.23	7.70	0.47	0.333
HDL, mmol/L	1.52 ^a	0.91 ^b	0.03	<0.001
LDL, mmol/L	0.27 ^a	0.06 ^b	0.13	<0.001
FFA, mmol/L	1.09 ^a	0.86 ^b	0.04	<0.001

¹ TG, triglycerides; TC, total cholesterol; HDL, high-density lipoprotein; LDL, low-density lipoprotein; FFA, free fatty acids.

² CON= basic diet; POR= basic diet+1% probiotic.

a, b within a row, values with different superscripts differ significantly at $p < 0.05$ or $p < 0.01$. Data are expressed as means \pm SEM.

Table 5. Effect of probiotics feed on intramuscular fat content and triglyceride content of the LT of Sunit sheep

Item¹	Treatment²		SEM	p-value
	CON	POR		
IMF, g/100g	0.46 ^b	0.57 ^a	0.04	0.024
TG, mmol/g prot	2.98 ^b	3.74 ^a	0.25	0.006

¹ IMF, intramuscular fat; TG, triglycerides.

² CON= basic diet; POR= basic diet+1% probiotic.

a, b within a row, values with different superscripts differ significantly at p<0.05 or p<0.01. Data are expressed as means ± SEM.

Table 6. Effects of probiotics feed on the fatty acid profiles of Sunit sheep

Items	Treatment ¹		SEM	p-value
	CON	PRO		
C10:0	0.11 ^a	0.08 ^b	0.02	0.048
C12:0	0.11	0.10	0.01	0.253
C14:0	2.13 ^a	1.67 ^b	0.17	0.020
C16:0	22.79	23.62	0.47	0.109
C18:0	15.36	15.26	0.39	0.798
C20:0	0.14	0.16	0.02	0.256
SFA ²	41.30	39.67	1.01	0.206
C14:1	0.47 ^b	1.01 ^a	0.09	<0.001
C16:1	2.56 ^a	1.89 ^b	0.16	0.010
C18:1n9 cis	44.22 ^b	45.61 ^a	0.54	0.026
MUFA ³	45.36 ^b	48.10 ^a	0.89	0.028
C18:2n6	7.32	8.74	0.69	0.097
C18:3n6	0.24 ^b	0.46 ^a	0.05	0.002
C18:3n3	0.45 ^b	0.57 ^a	0.05	0.036
C20:3n6	0.13 ^b	0.38 ^a	0.06	0.041
C20:4n6	0.21 ^b	0.40 ^a	0.02	<0.001
C20:5n3	0.38 ^b	0.48 ^a	0.04	0.017
C24:1n3	0.12 ^b	0.16 ^a	0.02	0.007
PUFA ⁴	10.78	11.88	1.17	0.371
n-3 PUFA ⁵	0.94 ^b	1.12 ^a	0.07	0.023
n-6 PUFA ⁶	10.28	9.43	1.07	0.445

¹ CON= basic diet; POR= basic diet +1% probiotic.

a, b within a row, values with different superscripts differ significantly at $p < 0.05$ or $p < 0.01$. Data are expressed as means \pm SEM.

² SFA=C10:0+C12:0+C14:0+C16:0+C18:0+C20:0.

³ MUFA=C14:1+C16:1+C18:1n9 cis.

⁴ PUFA=C18:2n6+C18:3n6+C18:3n3+C20:3n6+C20:4n6+C20:5n3+C24:1n3.

⁵ n-3 PUFA=C18:3n3+C20:5n3+C22:6n3.

⁶ n-6 PUFA=C18:2n6+C18:3n6+C20:3n6+C20:4n6.

ACCEPTED

Table 7. Correlation between lipid composition parameters and gene expression of Sunit sheep

	AMPKα2	ACCa	CPT1B	SREBP-1c	FASN	SCD1
	Pearsonr					
IMF	-0.585*	0.256	-0.429	0.437	0.734**	0.111
TG	-0.233	-0.032	-0.200	0.035	0.217	0.465
C10:0	0.558*	-0.434	0.538*	-0.329	0.090	-0.509*
C12:0	0.307	0.050	0.151	0.095	0.118	0.145
C14:0	0.286	-0.216	0.332	-0.118	-0.077	-0.007
C16:0	-0.097	0.459	-0.320	0.344	0.068	0.066
C18:0	0.029	-0.095	0.015	-0.170	0.115	-0.361
C20:0	0.062	0.449	-0.195	0.195	0.280	0.377
C14:1	-0.536*	0.706**	-0.633**	0.550**	-0.130	0.479
C16:1	0.556*	-0.730**	0.718**	-0.454	-0.052	-0.408
C18:1cis-9	0.487*	-0.603*	0.563*	0.260	0.127	-0.386
C18:2n6	-0.384	0.635**	-0.559*	0.213	-0.227	0.362
C18:3n6	-0.493*	0.701**	-0.578*	0.410	-0.185	0.544*
C18:3n3	-0.268	0.349	-0.367	0.092	-0.069	0.437
C20:3n6	-0.662**	0.799**	-0.790**	0.599**	-0.058	0.534*
C20:4n6	-0.656**	0.810**	-0.797**	0.572**	-0.105	0.539*
C20:5n3	-0.157	0.169	-0.096	0.077	-0.108	0.205
C24:1n3	-0.384	0.530*	-0.502*	0.283	0.038	0.247

* and ** indicated the significance of the correlations.

ACCEPTED










Fig 1. Effects of probiotics on composition of the fatty acids in the LT of Sunit Sheep. (A) saturated fatty acid; **(B)** monounsaturated fatty acid; **(C)** polyunsaturated fatty acid; **(D)** total fatty acid. CON= basic diet; POR= basic diet +1% probiotic.*p < 0.05, **p < 0.01.

coefficients of the linear combination of principal component 1 and principal component 2.

CON= basic diet; POR= basic diet +1% probiotic.

ACCEPTED






Fig 3. Effects of probiotics feed on the relative expression of mRNA of lipid metabolism genes from Sunit sheep. (A), (B), (C), (D), (E), (F) represents the relative of AMPKα2, ACCα, CPT1B, SREBP-1c, FASN, and SCD1 in the CON group and the PRO group, respectively. (G) represents the value of log₂ fold change for each gene.

CON= basic diet; POR= basic diet +1% probiotic. * p< 0.05, **p<0.01.




Fig 4. Effect of probiotics feeds on the activity of enzymatic of key molecules of lipid metabolism in Sunit sheep. (A), (B), (C), (D), (E), (F) represents the activity of enzymatic of AMPK α 2, ACC α , CPT1B, SREBP-1c, FASN, and SCD1 in the CON group and the PRO group, respectively. CON= basic diet; POR= basic diet +1% probiotic. * p < 0.05, **p < 0.01.

ACCEPTED




Fig 5. Possible pathways by which probiotics feed affect the AMPK signaling pathway to promote IMF deposition and improve fatty acid composition.

ACCEPTED