



Evaluation on Antimicrobial Activity of *Psoraleae semen* Extract Controlling the Growth of Gram-Positive Bacteria

Sejeong Kim^{1,2}, Soomin Lee^{1,2}, Heeyoung Lee^{1,2}, Jimyeong Ha^{1,2}, Jeeyeon Lee^{1,2},
Yukyung Choi^{1,2}, Hyemin Oh^{1,2}, Joonbae Hong³, Yohan Yoon^{1,2*}, and Kyoung-Hee
Choi^{4,5*}

¹Department of Food and Nutrition, Sookmyung Women's University

²Risk Analysis Research Center, Sookmyung Women's University

³Test & Research Department, Consumer Safety Center, Korea Consumer Agency

⁴Department of Oral Microbiology, College of Dentistry, Wonkwang University

⁵Institute of Biomaterials-Implant, Wonkwang University

Abstract

This study investigated bacterial growth-inhibitory effect of 69 therapeutic herbal plants extracts on 9 bacterial strains using a disc diffusion assay. Especially, the antimicrobial activity of *Psoraleae semen*, which showed different activity on pathogenic Gram-positive and Gram-negative bacteria, was evaluated by MIC (minimal inhibition concentration) and biofilm formation assay. The effect of *Psoraleae semen* extract on bacterial cell membranes was examined by measurement of protein leakage (optical density at 280 nm) and scanning electron microscope (SEM). No clear zone was formed on discs containing Gram-negative bacteria, but Gram-positive bacteria exhibited clear zones. The MICs of *Psoraleae semen* extract were 8 µg/mL for *Streptococcus mutans*, and 16 µg/mL for Enterococci and *Staphylococcus aureus*. In addition, biofilm formation was inhibited at concentration 8-16 µg/mL. Protein leakage values and SEM images revealed that cell membranes of Gram-positive bacteria were impaired following exposure to the extract. Further, the extract inhibited the growth of *Listeria monocytogenes* in sausages. These results indicate that *Psoraleae semen* extract could be utilized as a natural antimicrobial agent against Gram-positive bacteria.

Keywords antibacterial activity, natural antimicrobials, therapeutic herbal plants

Received February 8, 2017

Revised June 21, 2017

Accepted July 4, 2017

*Corresponding authors

Kyoung-Hee Choi
College of Dentistry, Wonkwang University, Iksan 54538, Korea
Tel: +82-63-850-6911
Fax: +82-63-850-6911
E-mail: kheechoi@wonkwang.ac.kr

Yohan Yoon
Department of Food and Nutrition,
Sookmyung Women's University, Seoul
04310, Korea
Tel: +82-2-2077-7585
Fax: +82-2-710-9479
E-mail: yoon@sookmyung.ac.kr

Introduction

For last decades, abuse of antibiotics has increased the prevalence of antibiotic-resistant bacteria, resulting in significant problems to human health (Fani *et al.*, 2007). Therefore, investigation into new antibacterial agents is required. Currently, there are negative perceptions regarding synthetic additives, and natural materials are attractive as alternative antimicrobial agents because of their limited effects on the environment and their low toxicity to human cells (Kaya *et al.*, 2008). Various lifeforms, including microorganisms, plants, and animals, have been investigated to identify novel antimicrobial agents that possess these favored characteristics (Guliani *et al.*, 2007; Yoon and Choi, 2010).

Plant-derived extracts, which are biodegradable and readily available, are known to possess antimicrobial activity against bacteria, yeast, and mold. Furthermore, they contain various antimicrobial components and phytochemicals, including alk-

© This is an open access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

aloids, terpenes, essential oils, and phenols. Therefore, plant-derived compounds are considered to be valuable sources of potential novel antimicrobial agents (Liu *et al.*, 2007). For instance, a mixture of lemon and cherry powder was effective at inhibiting *Clostridium perfringens* growth in frankfurters and ham (Jackson *et al.*, 2011), and an extract from the stem bark of *Iringia gabonensis*, known as African mango, also showed antimicrobial activity on various bacteria such as *Enterococcus faecalis*, *Staphylococcus aureus*, and *Bacillus cereus* at concentrations of 75-160 µg/mL (Kuete *et al.*, 2007).

Psoraleae semen is the seed of natural herb *Psoralea corylifolia* L. (Shim *et al.*, 2009). It has been used in traditional Asian medicine, and it has been proved that they have many pharmacological effects including anti-angiogenesis, anti-cancer, and promoting bone formation (Hwang *et al.*, 2013; Xiong *et al.*, 2003). Extract of the plant contains many useful compounds such as bakuchiol, bavachinin, corylin, corylifolin, isopsorlen, psoracorylifols, psoralen, and psoralidin (Wong and Rabiem, 2010). In particular, bakuchiol was found to inhibit growth of *Helicobacter pylori* (Zaide *et al.*, 2009), and Yoon and Choi (2012) reported that among various therapeutic herbal extracts, *Psoraleae semen* had a considerable antilisterial effect. Nevertheless, the antimicrobial effect of this extract on other important pathogenic bacteria remains to be determined.

In the present study, various therapeutic herbal plant extracts were screened for antimicrobial activity, and one of them, *Psoraleae semen* was evaluated as an antimicrobial agent in *in vitro* and in sausages for application in food, and its effect on bacterial cells was observed by measuring protein leakages and observing a scanning electron microscope (SEM).

Materials and Methods

Preparation of therapeutic herbal plant extracts

Sixty-nine of herbal plant extract powder were acquired from New Natural Material Bank of Korea National Research Resources (Korea) (Table 1). The powder was produced by heating 70% ethanol extraction, followed by vacuum evaporation of the filtrate. A liquid stock was made by dissolving the extract powder in 100% dimethyl sulfoxide (DMSO; Sigma, USA) to give a final concentration of 50 mg/mL.

Bacterial strains and media

The following bacterial strains were used in this study;

Gram-negative bacteria (*Acinetobacter baumannii* ATCC 19606, *Burkholderia thailandensis* E264, *Klebsiella pneumoniae* ATCC25306, *Pseudomonas aeruginosa* PAO1, *Salmonella enterica* serovar Derby ATCC6960) and Gram-positive bacteria (*E. faecalis* ATCC35038, *Enterococcus faecium* ATCC19434, *S. aureus* ATCC25923, *Streptococcus mutans* ATCC25175), and brain heart infusion (BHI, Becton, Dickinson and Company, USA) broth was used as the bacterial growth medium.

Disc diffusion assay

Each bacteria was activated and subcultured in BHI broth, and incubated at 37°C until they were grown to mid-logarithmic phase. The bacteria were mixed with 3 mL of BHI medium containing 0.8% agar to obtain an optical density of 0.1 at 600 nm (OD₆₀₀), and the mixtures were overlaid on BHI agar plates. After solidifying the agar under room temperature, the paper discs (6 mm in diameter) containing 3 µL of the extract (50 mg/mL) were placed on the agar plates. After 24 h incubation at 37°C, the diameters of inhibition zones were measured.

Determination of MICs

The minimum inhibitory concentration (MIC) of *Psoraleae semen* extract on bacterial growth was determined using the two-fold broth micro-dilution method (NCCLS, 1997). Briefly, each bacterial strain was subcultured and was grown to mid-logarithmic growth phase in BHI broth. The bacterial cells were diluted in BHI broth to an optical density of 0.01 at 600 nm. An equal volume of the bacterial diluent was added to each well on the 96-well plate containing 100 µL of the serially-diluted *Psoraleae semen* extract (0-256 µg/mL). The 96-well plates were incubated at 37°C for 24 h, and the MICs were determined by measuring OD₆₀₀ using a spectrophotometer (Sunrise, Tecan, Austria).

Biofilm assay

Biofilm formation experiment was performed by referring to the protocol provided by Loo *et al.* (2000). Bacterial cells were prepared and treated with *Psoraleae semen* extract following the method used to determine the MIC. Each bacterial culture was incubated in a 96-well plate containing *Psoraleae semen* extract (0-32 µg/mL) at 37°C for 24 h. The suspensions were carefully discarded, and the biofilm formed on the surface of the well plate was stained using 200 µL of crystal violet (1%; v/v). The stained cells were rinsed twice with 300 µL of PBS, fol-

lowed by solubilizing the cells in 200 μ L of 95% ethanol, and the degree of biofilm formation was determined by measuring the OD₅₇₅ with a spectrophotometer.

Protein leakage measurement

Subcultured bacterial cells were suspended in phosphate buffered solution (PBS, pH 7.4; 0.2 g of KH₂PO₄, 1.5 g of Na₂HPO₄, 8.0 g of NaCl, and 0.2 g of KCl in 1 L of distilled water) for adjusting OD₆₀₀ to 0.7. The bacterial cells were exposed to 2×MIC of *Psoraleae semen* extract or equal volume of DMSO for 1 h. After centrifugation, protein leakage was determined by measuring optical density of the cell supernatants using a spectrophotometer (Bio-photometer, Eppendorf, Germany) at OD₂₈₀. Values were calibrated with control group treated in PBS instead of cultured cells, and relative ratio (OD₂₈₀ of bacterial cell exposed to the extract/OD₂₈₀ of bacterial cell exposed to DMSO) was calculated.

Scanning electron microscopy

Bacteria were incubated in BHI broth containing a glass slide to allow bacterial cells to adhere to the glass slide. Adherent cells were exposed to 2×MIC of *Psoraleae semen* extract in PBS, and the bacterial cells that had been treated with DMSO at the same volume as *Psoraleae semen* extract, were used as a negative control (Bereksi *et al.*, 2002). After pre-fixing the treated cells in 1.8% glutaraldehyde solution (Sigma, USA), they were washed using distilled water. The bacterial cells were post-fixed with 2% osmium tetra oxide (Sigma, USA), followed by washing and dehydrating sequentially in 25, 50, 75, 90, and 100% ethanol. Bacterial cells were platinum-coated using a sputter coater 108 auto (Cressington Scientific Instruments Ltd., England) and then observed under field emission SEM (JEOL Ltd., JSM-7600F, Japan).

Statistical analysis

The experiments were replicated twice with six samples in each replication. The results were analyzed by the mixed model procedure of SAS[®] version 9.2 (SAS Institute, USA). Mean comparisons for the fixed effect were performed with the pairwise *t*-test, and significance was determined at alpha = 0.05.

Evaluation of antimicrobial activity in sausages

Inoculum for sausage contamination was prepared as follows. Two strains of *Listeria monocytogenes* (sausage isolate) cells identified and provided by Korea Consumer

Agency (Korea) were incubated in BHI broth. The cultures were centrifuged at 1,912 g and 4°C for 15 min. The cell pellet was washed with PBS and resuspended in PBS. Sausage formulation was prepared by mixing lean pork (58.5%), pork fat (19.5%), NaCl (1.2%), sodium phosphate (0.3%), isolated soy protein (0.5%), spice (0.4%), and ice water (19.5%) using a blender (HR1372, Phillips, China). Ten grams of homogenized formulations were transferred into a 6-well plate, and were boiled at 80°C in a waterbath for 40 min, followed by cooling. *Psoraleae semen* extract (final concentration: 2×MIC) or same volume of DMSO was spread on the sausage samples, and the samples were dried. The sausages were inoculated with *L. monocytogenes* inoculum to obtain 2 Log CFU/g. During storage at 4°C, *L. monocytogenes* growth was determined by plating sausages samples on Palcam agar on day 0, 5, and 10.

Results and Discussion

In a previous study, *Psoraleae semen* exhibited antimicrobial activity against *L. monocytogenes*, and it was suggested that the herb has potential to be used as an antibacterial agent (Yoon and Choi, 2012). This preliminary evaluation was further progressed and the effect was assessed against an extended range of pathogens such as Enterococci, *S. aureus* and *Salmonella* Derby in the present study. In addition, antimicrobial activities of 69 herbal plant extracts on various pathogens including Gram-positive and Gram-negative bacteria were tested using a disc diffusion assay. Among them, only five extracts (*Rubi fructus*, *Galla rhois*, *Moutan xortex Radicis*, *Sanguisorba officinalis* Linne, and *Psoraleae semen*) were effective for inhibiting more than three different bacteria (Table 1). *Rubi fructus* inhibited growth of all bacterial strains tested in this study. Interestingly, there were no clear zones formed on the disc containing *Psoraleae semen* extract against any Gram-negative bacteria, but obvious clear zones (7-13 mm) were observed against Gram-positive bacteria such as *S. mutans*, *E. faecium*, *E. faecalis*, and *S. aureus* (Table 1). This contrary result observed against Gram-positive and Gram-negative bacteria in the presence of *Psoraleae semen* might be caused by the structural dissimilarity of the cell wall between them (Henie *et al.*, 2009). Low permeability against lipophilic solutes and good adaptability of outer membrane is the key of resistance in Gram-negative bacteria (Plesiat and Nikaido, 1992). For example, Gram-negative bacteria have outer membranes, which may prevent the extract used in this study from

Table 1. Therapeutic herbal extracts (50 mg/mL) used in this study and their antimicrobial activities against Gram-positive and Gram-negative bacteria

Number of NNMBS	Name of extracts	Scientific name of extracts	Gram-negative bacteria					Gram-positive bacteria				
			AB ²	BT	KP	PA	SE	EFS	EFM	SA	SM	
NNMBS4	Gaehyuldeng	<i>Spatholobi caulis</i>	* ³	*	*	*	*	*	*	*	*	*
NNMBS7	Gosam	<i>Sophorae radix</i>	*	*	*	*	*	*	*	*	*	*
NNMBS10	Ildangui	<i>Angelicae radix</i>	*	*	*	*	*	*	*	*	*	*
NNMBS13	Mogua	<i>Chaenomelis fructus</i>	*	*	*	*	*	*	*	*	*	*
NNMBS16	Baekduong	<i>Pulsatillae radix</i>	*	-	*	*	*	*	*	*	*	*
NNMBS19	Baeksunpi	<i>Dictamni radices Cortex</i>	*	-	*	-	*	*	*	*	*	*
NNMBS22	Baekji	<i>Angelicae dahuricae Radix</i>	*	*	*	*	*	*	*	*	*	*
NNMBS25	Bokbunja	<i>Rubi fructus</i>	‡	§	‡	‡	‡	‡	‡	‡	‡	‡
NNMBS31	Sansuyu	<i>Corni fructus</i>	*	*	*	*	*	*	*	*	*	*
NNMBS34	Sanglyuk	<i>Phytolaccae radix</i>	*	*	*	*	*	*	*	*	*	*
NNMBS37	Sangbaekpi	<i>Mori radices Cortex</i>	*	*	*	*	*	*	†	†	*	*
NNMBS40	Sansimja	<i>Mori fructus</i>	*	*	*	*	*	*	*	*	*	*
NNMBS42	Aeyub	<i>Artemisiae argyi Herba</i>	*	*	*	*	*	*	*	*	*	*
NNMBS45	Youngsil	<i>Rosae fructus</i>	*	*	*	*	*	*	*	*	*	*
NNMBS48	Omija	<i>Schizandrae fructus</i>	*	-	*	*	*	*	*	*	*	*
NNMBS51	Obaeja	<i>Galla rhois</i>	‡	§	‡	‡	*	‡	‡	‡	‡	‡
NNMBS54	Oyak	<i>Linderae radix</i>	‡	*	*	*	*	*	*	*	*	*
NNMBS60	Wonji	<i>Polygalae radix</i>	*	*	*	*	*	*	*	*	*	*
NNMBS63	Yukyongjong	<i>Cistanchis gerba</i>	*	*	*	*	*	*	*	*	*	*
NNMBS66	Baekjakyak	<i>Paeoniae radix</i>	*	*	*	*	*	*	*	*	*	*
NNMBS69	Chunlyunja	<i>Meliae fuctus</i>	*	*	*	*	*	*	*	*	*	*
NNMBS72	Taeksa	<i>Alismatis rhizoma</i>	*	*	*	*	*	*	*	*	*	*
NNMBS75	Panlamgun	<i>Indigo rulverata Levis</i>	*	*	*	*	*	*	*	*	*	*
NNMBS78	Jeukhasuo	<i>Polygoni multiflori Radix</i>	*	*	*	*	*	*	*	*	*	*
NNMBS81	Dansam	<i>Salviae miltiorrhizae Radix</i>	*	*	*	*	*	*	*	†	*	*
NNMBS84	Yongdam	<i>Gentianae scabrae Radix</i>	*	*	*	*	*	*	*	*	*	*
NNMBS87	Jagun	<i>Lithospermi radix</i>	*	*	*	*	*	*	*	†	*	*
NNMBS90	Kunjihwang	<i>Rehmanniae radix</i>	*	*	*	*	*	*	*	*	*	*
NNMBS92	Hyunsam	<i>Scrophulariae radix</i>	*	*	*	*	*	*	*	*	*	*
NNMBS94	Hwanggum	<i>Scutellariae radix</i>	*	*	*	*	*	*	*	†	*	*
NNMBS97	Gunkang	<i>Zingiberis rhizoma</i>	*	*	*	*	*	*	*	*	*	*
NNMBS100	Daehwang	<i>Rhei rhizoma</i>	*	*	*	*	*	*	*	*	*	*
NNMBS103	Sungma	<i>Cimicifugae rhizoma</i>	*	*	*	*	*	*	*	*	*	*
NNMBS105	Jimo	<i>Anemarrhenae rhizoma</i>	*	-	*	*	*	*	*	*	*	*
NNMBS108	Chunkung	<i>Cnidii rhizoma</i>	*	*	*	*	*	*	*	*	*	*
NNMBS111	Chunma	<i>Gastrodiae rhizoma</i>	*	*	*	*	*	*	*	*	*	*
NNMBS114	Hyangbuja	<i>Cyperii rhizoma</i>	*	*	*	*	‡	*	*	*	*	*
NNMBS117	Hyunhosaeak	<i>Corydalis tuber</i>	*	*	*	*	*	*	*	*	*	*
NNMBS120	Hwanglyun	<i>Coptidis rhizoma</i>	*	*	*	*	*	*	*	†	*	*
NNMBS123	Kyulmyungja	<i>Cassiae semen</i>	‡	*	*	*	*	*	*	†	*	*
NNMBS126	Binlangja	<i>Arecae semen</i>	*	*	*	*	*	*	*	†	*	*
NNMBS129	Yukdugu (Nutmeg)	<i>Myristicae semen</i>	*	*	*	*	*	*	*	†	*	*
NNMBS132	Chajunja	<i>Plantaginis semen</i>	*	-	*	-	*	*	*	*	*	*
NNMBS135	Kugija	<i>Lycii fructus</i>	*	*	*	*	*	*	*	†	*	*
NNMBS141	Sancho	<i>Zanthoxyli fructus</i>	*	*	*	*	*	*	*	*	*	*
NNMBS144	Yunkyo	<i>Forsythiae fructus</i>	*	-	-	-	*	*	*	*	*	*

NNMBS: Standard sample from new natural materials bank.

²AB, *Acinetobacter baumannii* ATCC19606; BT, *Burkholderia thailandensis* E264; KP, *Klebsiella pneumoniae* ATCC25306; PA, *Pseudomonas aeruginosa* PAO1; SE, *Salmonella enterica* serovar Derby ATCC6960; EFS, *Enterococcus faecalis* ATCC35038; EFM, *Enterococcus faecium* ATCC19434; SA, *Staphylococcus aureus* ATCC25923; SM, *Streptococcus mutans* ATCC25175.³-, not determined; *, no clear zone; †, <10 mm; ‡, 10 mm-12 mm; §, >12 mm.

Table 1. Therapeutic herbal extracts (50 mg/mL) used in this study and their antimicrobial activities against Gram-positive and Gram-negative bacteria (Continued)

Number of NNMBS	Name of extracts	Scientific name of extracts	Gram-negative bacteria					Gram-positive bacteria			
			AB ²	BT	KP	PA	SE	EFS	EFM	SA	SM
NNMBS147	Osuyu	<i>Evodiae fructus</i>	*	-	*	*	*	*	*	†	*
NNMBS150	Ikjiin	<i>Alpiniae fructus</i>	*	*	*	*	*	*	*	*	*
NNMBS156	Jinpi	<i>Aurantii nobilis Pericarpium</i>	*	*	*	*	*	*	*	†	*
NNMBS159	Chija	<i>Gardeniae fructus</i>	*	*	*	*	*	*	*	*	*
NNMBS163	Umyanggwak	<i>Epimedii herba</i>	*	*	*	*	*	*	*	*	*
NNMBS166	Ikmocho	<i>Leonuri herba</i>	*	*	*	*	*	*	*	*	*
NNMBS172	Duchung	<i>Eucommiae xortex</i>	*	*	*	*	*	*	*	*	*
NNMBS175	Mokdanpi	<i>Moutan xortex Radicis</i>	†	-	†	*	†	†	*	†	†
NNMBS178	Ogapi	<i>Acanthopanax cortex</i>	*	*	*	*	*	*	*	*	*
NNMBS179	Hwangbaek	<i>Phellodendri cortex</i>	*	*	*	*	*	*	*	†	*
NNMBS182	Hubak	<i>Magnoliae cortex</i>	*	*	*	*	*	*	*	*	*
NNMBS188	Kumunwha	<i>Lonicerae flos</i>	*	*	*	*	*	*	*	*	*
NNMBS191	Hagocho	<i>Prunellae spica</i>	*	*	*	*	*	*	*	*	*
NNMBS203	Jiyu	<i>Sanguisorba officinalis Linne</i>	‡	-	‡	-	†	†	†	†	†
NNMBS209	Bogolji	<i>Psoraleae semen</i>	*	*	*	*	*	†	†	†	‡
NNMBS212	Sasangja	<i>Torilis fructus</i>	*	*	*	*	*	*	*	*	*
NNMBS215	Omae	<i>Mume fructus</i>	*	*	*	*	*	*	*	*	*
NNMBS220	Jiguja	<i>Hoveniae semen cum Fructus</i>	*	*	*	*	*	*	*	†	*
NNMBS222	Jilyuja	<i>Tribuli fructus</i>	*	*	*	*	*	*	*	†	*
NNMBS225	Chogwa	<i>Amomi tsao-ko Fructus</i>	*	*	*	*	*	*	*	†	*
NNMBS229	Chodugu	<i>Alpiniae kotsumadaai Semen</i>	*	*	*	*	*	*	*	*	*
NNMBS231	Sangkisaeng	<i>Loranthi rtamulus</i>	*	*	*	*	*	*	*	†	*
NNMBS234	Injinho	<i>Artemisiae capillaris Herba</i>	*	*	*	*	*	*	*	†	*

NNMBS: Standard sample from new natural materials bank.

²AB, *Acinetobacter baumannii* ATCC19606; BT, *Burkholderia thailandensis* E264; KP, *Klebsiella pneumoniae* ATCC25306; PA, *Pseudomonas aeruginosa* PAO1; SE, *Salmonella enterica* serovar Derby ATCC6960; EFS, *Enterococcus faecalis* ATCC35038; EFM, *Enterococcus faecium* ATCC19434; SA, *Staphylococcus aureus* ATCC25923; SM, *Streptococcus mutans* ATCC25175.

³-, not determined; *, no clear zone; †, <10 mm; ‡, 10 mm-12 mm; §, >12 mm.

exerting its antimicrobial activity. However, further study is necessary to elucidate the antimicrobial mechanisms of this extract against Gram-negative bacteria.

The antimicrobial effect of *Psoraleae semen* extract was further assessed by MIC determination for Gram-positive bacteria, which were susceptible to *Psoraleae semen* extract. As a result, *S. mutans* ATCC25175 exhibited no visible growth at 8 µg/mL, and *E. faecalis* ATCC35038, *E. faecium* ATCC19434, and *S. aureus* ATCC25923 exhibited no growth at 16 µg/mL (Fig. 1). Yim *et al.* (2013) also found that the MIC of *Psoraleae semen* extract was 625 µg/mL for *E. faecalis* and 156 µg/mL for *S. mutans*. This discrepancy may be due to use of different fractions of the extract for the experiment. Yim *et al.* (2013) used a water-soluble fraction of *Psoraleae semen*, whereas our group used total extract, which contains both water-soluble and water-insoluble fractions. Yoon and Choi (2012) suggested that the water-insoluble fraction of *Psoraleae semen* exhibits higher antimicrobial activity than the water-soluble

fraction. *Psoraleae semen* extract contains bakuchiol in water-insoluble fraction, which has been known to have antimicrobial activity on various oral bacteria (Katsura *et al.*, 2001), as Yoon and Choi (2012), and Yim *et al.* (2013) suggested that it may be a major compound responsible for the antimicrobial activity of *Psoraleae semen* against the bacteria.

Biofilm is a collection of bacterial cells that form on surfaces in reaction to stressful environmental conditions, and biofilm formation is of concern in various areas including food safety, dental health, and medicine (Loesche, 1986). In this study, the effect of *Psoraleae semen* extract on biofilm formation was investigated. Biofilm formation of *S. mutans* ATCC25175 and *S. aureus* ATCC25923 was inhibited when the extract was present at concentrations greater than 8 µg/mL, and it was significantly decreased when concentrations exceeded 16 µg/mL in *E. faecalis* ATCC35038 (Fig. 2). *S. mutans* biofilms, namely plaque, are responsible for dental diseases like caries (Loesche,




Fig. 1. Minimum inhibition concentration of *Psoraleae semen* extract on *Streptococcus mutans* ATCC25175 (a), *Enterococcus faecalis* ATCC35038 (b), *Enterococcus faecium* ATCC19434 (c), and *Staphylococcus aureus* ATCC25923 (d). Means with different letters are significantly different ($p < 0.05$).




Fig. 2. Biofilm formation of *Streptococcus mutans* ATCC25175 (a), *Enterococcus faecalis* ATCC35038 (b), and *Staphylococcus aureus* ATCC25923 (c) after exposure of *Psoraleae semen* extract. Means with different letters are significantly different ($p < 0.05$).

1986), and *S. aureus* and *E. faecalis* biofilms are a problem in the food industry and medical devices (Hood and Zottola, 1995). Thus, our results indicate that *Psoraleae semen* extract may be of potential use for the inhibition of biofilm formation by these bacterial strains.

The effects of antimicrobial agents on bacterial cell structures and membranes have been widely used to investigate the mode of action of natural antimicrobial agent (Leela and Satirapipathkul, 2011). In this study, protein leakage induced by cell membrane damage was measured, and the membranes were observed by SEM analysis following exposure to *Psoraleae semen*. In case of Gram-positive bacteria (*E. faecalis* and *E. faecium*), values of OD₂₈₀ were

increased in extract-treated cells, compared to non-treated cells (relative ratios = 3.17 and 3.07 in *E. faecalis* and *E. faecium*, respectively). This result suggests that exposure of the extract to the bacterial cells induce membrane damage, resulting in leakage of intracellular proteins. Whereas, relative ratio of OD₂₈₀ (= 1.57) was relatively low in *Salmonella* Typhimurium, a Gram-negative bacteria (Fig. 3). Further, SEM images showed that the membranes of Gram-positive bacteria, such as *E. faecalis*, *E. faecium*, and *S. aureus* cells have been damaged by the treatment of *Psoraleae semen* extract, whereas no damage was observed in *Sal. Typhimurium* (Fig. 4). Taken together, antibacterial activity of *Psoraleae semen* extract on Gram-positive




Fig. 3. Relative ratio of OD₂₈₀ in bacterial cells exposed to 2× MIC of *Psoraleae semen* extract to the one exposed to same volume of DMSO in ST (*Salmonella* Typhimurium NCCP 10812), EFS (*Enterococcus faecalis* ATCC35038), EFM (*Enterococcus faecium* ATCC19434), and SA (*Staphylococcus aureus* ATCC25923).




Fig. 5 Growth of *Listeria monocytogenes* in sausages with *Psoraleae semen* extract (opened circle) or DMSO (closed circle) after incubation at 4°C for 10 d.




Fig. 4. Bacterial cell membrane of *Salmonella* Typhimurium NCCP10812 (A), *Enterococcus faecalis* ATCC35038 (B), and *Enterococcus faecium* ATCC19434 (C), observed by scanning electron microscope following exposure to DMSO (a; negative control) or *Psoraleae semen* extract (b).

bacteria seems to be attributed to this membrane damage-mediated action, but *Sal.* Typhimurium possessing outer membrane, which contributes to homeostasis and adaptability, avoids membrane injury by the extract. Although

S. aureus is gram-positive bacteria, protein leakage was not observed in this species. This can be explained by two possible reasons as follows. First, longer incubation time may have been required for inhibition of the *S. aureus* growth. Briefly, MIC against *S. aureus* was determined after 24 h, whereas protein leakage by the extract was observed only after 1 h treatment, which can be too short to induce the effects on *S. aureus* compared to the other bacteria. Second, structural difference of the bacteria probably causes discrepant outcome. *S. aureus* cell wall is solubilized by lysostaphin, unlike the ones of other Gram-positive bacteria can be solubilized by lysozyme (Navarre and Schneewind, 1999).

A Gram-positive bacterial pathogen *L. monocytogenes* usually contaminates livestock products such as ham and sausages, and can grow at low storage temperature. Our research group previously demonstrated that *Psoraleae semen* extract (named bogolgi) had antibacterial effects on this pathogen in *in vitro* environment (Yoon and Choi, 2012). Subsequently, current study investigated this antimicrobial activity of *Psoraleae semen* extract on livestock products-originated *L. monocytogenes* in sausages at refrigeration temperature. As a result, the growth of *L. monocytogenes* was impeded in the sausages treated with the extract, compared to the ones treated with DMSO as a negative control on day 10 at 4°C (Fig. 5). This result may suggest that *Psoraleae semen* extract could be applied to control *L. monocytogenes* growth in processed meat products. Although *Psoraleae semen* extract exhibits these antimicrobial activities, it was also reported that high dosage

of psoralens, the component of *Psoraleae semen* could induce phototoxicity (Wu *et al.*, 2005). Thus, further study is necessary to find appropriate conditions and procedure to use *Psoralea semen* extract as food antimicrobial.

In conclusions, *Rubi fructus*, *Galla rhois*, *Moutan xortex Radicis*, *Sanguisorba officinalis* Linne, and *Psoraleae semen* have antimicrobial activity against various pathogenic bacteria. Among them, *Psoraleae semen* extract could be used as a natural antimicrobial to control the growth of Gram-positive bacteria including Enterococci, *S. aureus*, and *L. monocytogenes*.

Acknowledgements

This work was supported by the Wonkwang university in 2016.

References

- Bereksi, N., Gavini, F., Benezech, T., and Faille, C. (2002) Growth, morphology and surface properties of *Listeria monocytogenes* Scott A and LO28 under saline and acid environments. *J. Appl. Microbiol.* **92**, 556-565.
- Fani, M. M., Kohanteb, J., and Dayaghi, M. (2007) Inhibitory activity of garlic (*Allum sativum*) extract on multidrug-resistant *Streptococcus mutans*. *J. Indian Soc. Pedod. Prev. Dent.* **25**, 164-168.
- Guliani, A., Pirri, G., and Nicoletto, S. F. (2007) Antimicrobial peptides: An overview of a promising class of therapeutics. *Cent. Eur. J. Biol.* **2**, 1-33.
- Henie, E. F. P., Zaiton, H., and Suhaila, M. (2009) Bacterial membrane disruption in food pathogens by *Psidium guajava* leaf extracts. *Int. Food Res. J.* **16**, 297-311.
- Hood, S. K. and Zottola E. A. (1995) Biofilms in food processing. *Food Control* **6**, 9-18.
- Hwang, J. H., Rha, S. J., Cho, J. K., Shin, T. S., Kim, M. Y., and Kim, J. D. (2013) *Psoraleae semen* extract inhibits angiogenesis and adipogenesis. *Biotechnol. Bioproc. Eng.* **18**, 942-949.
- Jackson, A. L., Kulchaiyawat, C., Sullivan, G. A., Sebranek, J. G., and Dickson, J. S. (2011) Use of natural ingredients to control growth of *Clostridium perfringens* in naturally cured frankfurters and hams. *J. Food. Prot.* **74**, 417-424.
- Katsura, H., Tsukiyama, R. I., Suzuki, A., and Kobayashi, M. (2001) *In vitro* antimicrobial activities of bakuchiol against oral microorganisms. *Antimicrob. Agents Chemother.* **45**, 3009-3013.
- Kaya, I., Yigit, N., and Benli, M. (2008) Antimicrobial activity of various extracts of *Ocimum Basilicum* L. and observation of the inhibition effect on bacterial cells by use of scanning electron microscopy. *Aft. J. Trad. CAM.* **5**, 363-369.
- Kuete, V., Wabo, G. F., Ngameni, B., Mbaveng, A. T., Metuno, R., Etoa, F. X., Ngadjui, B. T., Beng, V. P., Meyer, J. J. M., and Lall, N. (2007) Antimicrobial activity of the methanolic extract, fractions and compounds from the stem bark of *Irvinia gabonensis* (Ixonanthaceae). *J. Ethnopharmacol.* **114**, 54-60.
- Leela, T. and Satirapipathkul, C. (2011) Studies on the antibacterial activity of quercus infectoria galls. *Int. Con. Biosci. Biochem. Bioinform. (IPCBE)* **5**, 410-414.
- Liu, C. S., Cham, T. M., Yan, C. H., Chang, H. W., Chen, C. H., and Chuan, L. Y. (2007) Antibacterial properties of Chinese herbal medicines against nosocomial antibiotic resistant strains of *Pseudomonas aeruginosa* in Taiwan. *Am. J. Chinese Med.* **35**, 1047-1060.
- Loesche, W. J. (1986) Role of *Streptococcus mutans* in human dental decay. *Microbiol. Rev.* **50**, 353-380.
- Loo, C. Y., Corliss, D. A., and Ganeshkumar, N. (2000) *Streptococcus gordonii* biofilm formation: identification of genes that code for biofilm phenotypes. *J. Bacteriol.* **182**, 1374-1382.
- Navarre, W. W. and Schneewind, O. (1999) Surface proteins of Gram-positive bacteria and mechanisms of their targeting to the cell wall envelope. *Microbiol. Mol. Biol. Rev.* **63**, 174-229.
- NCCLS (National Committee for Clinical Laboratory Standards). (1997) Performance standards for antimicrobial disk susceptibility tests: Approved standard M2-A7. Wayne: PA: USA.
- Plesiat, P. and Nikaido, H. (1992) Outer membranes of gram-negative bacteria are permeable to steroid probes. *Mol. Microbiol.* **6**, 1323-1333.
- Shim, S. H., Kim, Y., Lee, J. Y., and Song, D. G. (2009) Aldose reductase inhibitory activity of the compounds from the seed of *Psoralea corylifolia*. *J. Korean Soc. Appl. Biol. Chem.* **52**, 568-572.
- Wong, R. W. K. and Rabiem, A. B. M. (2010) Effect of Buguzhi (*Psoralea corylifolia* fruit) extract on bone formation. *Phytother. Res.* **24**, 155-160.
- Wu, F., Wang, A. F., Zhou, Y., Zhou, T. S., Jiang, H. L., and Fang, Y. Z. (2005) Determination of psoralen and isopsoralen in traditional Chinese medicines by capillary zone electrophoresis with amperometric detection. *Chromatographia* **61**, 157-160.
- Xiong, Z., Wang, D., Wu, Y., and Li, F. (2003) Osteoblastic differentiation bioassay and its application to investigating the activity of fractions and compounds from *Psoralea corylifolia* L. *Int. J. Pharmaceutic. Sci.* **58**, 925-928.
- Yim, N. H., Jung, Y. P., Cho, W. K., Kim, T., Kim, A., Im, M., and Ma, J. Y. (2013) Screening of aqueous extracts of medicinal herbs for antimicrobial activity against oral bacteria. *J. Integr. Med.* **2**, 18-24.
- Yoon, Y. and Choi, K. H. (2010) Identification of inhibitory effect on *Streptococcus mutans* by oleanolic acid. *J. Life Sci.* **3**, 321-325.
- Yoon, Y. and Choi, K. H. (2012) Antimicrobial activities of therapeutic herbal plants against *Listeria monocytogenes* and the herbal plant cytotoxicity on Caco-2 cell. *Lett. Appl. Microbiol.* **55**, 47-55.

25. Zaide, S. F., Yamada, K., Kadowaki, M., Usmanhani, K., and Sugiyama, T. (2009) Bactericidal activity of medicinal plants, employed for the treatment of gastrointestinal ailments, against *Helicobacter pylori*. *J. Ethnopharmacol.* **121**, 286-291.