



ARTICLE

## Effect of Aged and Fermented Garlic Extracts as Natural Antioxidants on Lipid Oxidation in Pork Patties

Hyun-Jin Lee, Dong-kyu Yoon, Na-yeon Lee, and Chi-ho Lee\*

Department of Food Science and Biotechnology of Animal Resources, College of Animal Bioscience and Technology, Konkuk University, Seoul 05029, Korea

OPEN ACCESS

Received May 3, 2019

Revised July 3, 2019

Accepted July 4, 2019

\*Corresponding author : Chi-Ho Lee  
Department of Food Science and  
Biotechnology of Animal Resources, College  
of Animal Bioscience and Technology,  
Konkuk University, Seoul 05029, Korea  
Tel: +82-2-450-3681  
Fax: +82-2-453-1948  
E-mail: [leech@konkuk.ac.kr](mailto:leech@konkuk.ac.kr)

\*ORCID  
Hyun-Jin Lee  
<https://orcid.org/0000-0001-6031-4966>  
Dong-kyu Yoon  
<https://orcid.org/0000-0003-1217-4502>  
Na-yeon Lee  
<https://orcid.org/0000-0002-3024-0043>  
Chi-ho Lee  
<https://orcid.org/0000-0001-9406-8310>

**Abstract** The aim of this study was to investigate the natural antioxidant activity of raw garlic (RG), aged black garlic (AG), and garlic fermented with *Bacillus subtilis* (FG) extracts on pork patty lipid oxidation throughout refrigerated storage. The total polyphenol, total flavonoid content, 1,1-diphenyl-2-picrylhydrazyl (DPPH) and 2,2'-azino-bis-3-ethylbenzothiazoline-6-sulfonic acid (ABTS) radical scavenging activity of three different types of garlic extracts were measured. The total phenolic and flavonoid content of AG was significantly higher than that of FG and RG; FG also showed a significantly higher total phenolic content than that of RG ( $p < 0.05$ ). The DPPH and ABTS radical scavenging activity of AG and FG was significantly higher than that of RG and that of AG was significantly higher than that of FG ( $p < 0.05$ ). To investigate the effect of processed garlic extracts on pork patty lipid oxidation, freeze-dried extracts of RG, FG, and AG were added to the patties at levels of 0.5% (w/w). Patties containing 0.01% (w/w) ascorbic acid (AA) and patties without treatment (CON) were compared with patties containing garlic extracts. The pH value, 2-thiobarbituric acid reactive substances value, and volatile basic nitrogen value of pork patties containing AG and FG extracts were significantly decreased compared to the other groups (CON, AA, and RG;  $p < 0.05$ ). Taken together, these results suggest that AG and FG extracts possess strong antioxidative activity and can serve as natural antioxidative additives to prevent pork patty lipid oxidation.

**Keywords** aged garlic, fermented garlic, *Bacillus subtilis*, pork patty, lipid oxidation

### Introduction

One of most important components in meat products is fat, which improves product flavor and texture when added at the appropriate amount (Kumar et al., 2016). However, oxidation of lipid is one of the main factors deteriorating the quality and shelf-life of meat products (Dave and Ghaly, 2011). Synthetic antioxidants, such as butylated hydroxytoluene (BHT) and butylated hydroxyanisole (BHA), have been used to inhibit the oxidation of various meat products (Chang et al., 1997). However,

carcinogenicity and safety issues have been suggested for synthetic antioxidants (Ito et al., 1986); thus, several studies have examined replacing synthetic antioxidants with natural additives such as rosemary extract (Sebranek et al., 2005), golden thread (*Coptis chinensis* Franch), clove extract (Zahid et al., 2018), and sapota powder (Kumar et al., 2018).

*Allium sativum*, commonly known as garlic, exhibits antioxidant, antimicrobial, and anticancer effects and reduces cardiovascular diseases and blood cholesterol (Harris et al., 2001). However, the spicy and irritating flavor of garlic limits its use as a common food additive; this flavor is caused by allicin and its derivatives (Jang et al., 2018). Alliin, a major component of garlic, is converted to allicin by allinase released from vacuoles when the cells are damaged. As allicin is unstable, it is degraded to lipophilic organic sulfur compounds, such as diallyl sulfide, diallyl disulfide, diallyl trisulfide, and thiosulfinate, which cause a distinct aroma (Rose et al., 2005).

Several studies have examined reducing the irritant flavor of garlic by processing. Aged garlic (also known as black garlic or red garlic) is processed by aging fresh garlic for several weeks at temperatures of 40°C to 90°C (Jang et al., 2008). Aged garlic has a less irritating flavor and increased functionality as a result of browning reactions (You et al., 2011). Processed garlic has greater antioxidant ability than raw garlic (RG) owing to the higher total phenolic and flavonoid content during processing (Imai et al., 1994).

Garlic can also be processed by fermentation. For example, *Bacillus subtilis*, a Gram-positive, catalase-positive bacterium common in soil environment (also known as grass bacillus or hay bacillus), can be used for the production of fermented soybean foods such as cheonggukjang (Korea), natto (Japan), thua nao (Thailand), kinema (India) and douchi (China) (Steinkraus, 1997). It is known that soybean and sword bean fermented with *B. subtilis* have increased antioxidant ability due to increased phenolic contents and flavonoids (Han et al., 2015; Juan and Chou, 2010). Zhang et al. (2014) identified antioxidant peptides from peanut meal fermented with *B. subtilis*. Moreover, recent studies have demonstrated that extracts from garlic fermented with *B. subtilis* alleviate cardiovascular diseases and blood cholesterol (Park et al., 2016; Park et al., 2017).

Processed garlic studies have mainly focused on the antioxidative (You et al., 2011) and physicochemical properties (Choi et al., 2008) of aged garlic or steamed garlic, as well as products supplemented with processed garlic such as bread and sausage (Shin et al., 2011; Wang et al., 2012). However, little is known regarding garlic processed by fermentation. Thus, our aim was to compare the antioxidant ability of garlic fermented with *B. subtilis* and aged black garlic (AG) prepared by conventional heat treatment and their potential use as natural antioxidants when added to pork patties by analyzing lipid oxidation during vacuum packaged, refrigerated storage.

## Materials and Methods

### Materials

Extracts of RG, aged garlic, and garlic fermented by *B. subtilis*, as described by Yun (2017), were purchased from OZL DNF Co., Ltd. (Dam Yang, Korea). The pork loin and pork back fat used to produce the pork patties were purchased from GUMDON Corp. (Won Ju, Korea).

### Sample preparation

RG, AG, and FG extracts were filtered using Whatman filter paper (No. 1), freeze-dried, and stored at -80°C in a deep freezer. Samples were diluted with distilled water to concentrations of 0.2%, 0.6%, and 1.0% (w/v).

### Total phenolic content assay

The Folin-Ciocalteu method was used to evaluate the total phenolic content of the three garlic extracts as described by Simoes et al. (2018). One milliliter of sample was mixed with 5 mL of 0.2 N Folin-Ciocalteu reagent for 10 min at 25°C. The mixture was supplemented with 4 mL of 7.5% (w/v) sodium carbonate and incubated at 25°C for 2 h. The absorbance was read at 765 nm using a spectrophotometer (Optizen 2120UV; Mecasys Co., Ltd., Daejeon, Korea). Distilled water was used as the blank control and flavonoid content was calculated with a standard curve by gallic acid.

### Total flavonoid assay

The total flavonoid content was measured using Dowd's method, as described by Adefegha et al. (2018). One milliliter of sample was mixed with the same amount of 2% (w/v) aluminum chloride then incubated for 10 min at 25°C. The absorbance of the mixture was read at 415 nm using an Optizen 2120UV spectrophotometer. Distilled water was used as the blank control and the contents were calculated based on a standard curve for quercetin.

### DPPH radical scavenging activity

1,1-diphenyl-2-picrylhydrazyl (DPPH) radical scavenging activity was measured using the methods of Blois (1958). One milliliter of sample was mixed with 5 mL of 100  $\mu$ M DPPH/95% (v/v) ethanol and incubated for 30 min at 25°C in a dark room. The absorbance was read at 517 nm using an Optizen 2120UV spectrophotometer and 1 mL of distilled water was mixed with 5 mL of DPPH solution for blank controls. DPPH radical scavenging activity was determined as:

$$\text{DPPH radical scavenging activity (\%)} = [1 - (A_1 / A_0)] \times 100$$

A<sub>1</sub>: sample absorbance

A<sub>0</sub>: blank control (i.e., DPPH solution with distilled water) absorbance

### ABTS radical scavenging activity

2'-azino-bis-3-ethylbenzothiazoline-6-sulfonic acid (ABTS) radical scavenging activity was measured using the methods of Re et al. (1999). An ABTS stock solution containing 7 mM ABTS and 2.45 mM potassium persulfate was prepared and incubated in a dark room for 12 h to form the ABTS<sup>•+</sup> radical. The ABTS stock solution was diluted with sodium phosphate buffer (pH 7.4) until the absorbance of the solution reaches 0.70±0.02 at 732 nm. Next, 2.9 mL of diluted ABTS solution were mixed with 0.1 mL of sample, incubated in a dark room for 10 min at 25°C, and the absorbance was read at 732 nm using an Optizen 2120UV spectrophotometer. Scavenging activity was calculated based on a standard curve for ascorbic acid.

### Preparation of pork patties

The basic composition of pork patty batter mix was consisted of 75% lean pork, 24% pork back fat, and 1% salt; they were ground through a 3 mm plate, divided into five groups, and each group was pre-mixed prior to additional treatment. The five groups included the CON (control), patties with no additional treatment; the AA group, patties with 0.1% (w/w) ascorbic acid; the RG group, patties with 0.5% (w/w) freeze-dried garlic extract; the FG group, patties with 0.5% (w/w) freeze-dried FG extract; and the AG group, patties with 0.5% (w/w) freeze-dried aged garlic extract. All groups were mixed using a bowl mixer (SM246, Poking Industrial Co., Ltd., Hong Kong, China) for 7 min, the paste was weighed, and 80±2 g patties were

formed using a rectangular burger press (Spikom Ltd., Nottingham, UK), in order to ensure the size of the patties was exactly the same. Once the patties were produced, a total of three batches of patties from each of the five groups were vacuum packaged and stored at 4°C for 21 d. Samples were taken at 0, 3, 7, 14, and 21 d of storage.

### pH analysis

The pH value of the pork patties was measured by homogenizing 2 g samples in 18 mL of distilled water by a homogenizer (AM-1, Nihon Seiki Kaisha Co., Ltd., Nagoya, Japan) for 1 min at 3,220×g. The pH of homogenate was then measured using a pH meter (LAQUA F-71, Horiba Co., Kyoto, Japan).

### TBARS value analysis

The 2-thiobarbituric acid reactive substances (TBARS) value was measured using the method of Buege and Aust (1978). Five grams of sample were mixed with 15 mL of distilled water plus 100 µL of 6% (w/v) BHT in ethanol solution and homogenized at 3,780×g for 1 min with an AM-1 homogenizer. Then, 2 mL of homogenate were mixed with 4 mL of TCA/TBA reagent (15% (w/v) trichloroacetic acid with 20 mM thiobarbituric acid solution) and heated in an 80°C water bath for 15 min. The samples were cooled in ice cold water, then vortexed and centrifuged for 10 min at 2,000×g, 25°C. After centrifugation, the supernatant of samples were filtered with Whatman filter paper No. 1, The absorbance of the filtrate was then measured at 531 nm using an Optizen 2120UV spectrophotometer. The malondialdehyde (MDA) concentration of the samples was determined using a standard curve for 1,1,3,3-tetraethoxypropane (TEP).

### Volatile basic nitrogen analysis

The volatile basic nitrogen (VBN) content of the samples was analyzed using Conway's microdiffusion method. Five grams of sample were mixed with 15 mL of distilled water and homogenized at 2,240×g for 1 min; the volume was then adjusted to 50 mL with distilled water in a mass cylinder. The homogenate was filtered with Whatman filter paper No. 1 and 1 mL of the filtrate was added to the outer chamber of the Conway unit. Next, 1 mL of 0.01 N boric acid and 100 µL of Conway reagent (0.066% (w/v) methyl red and 0.066% (w/v) bromocresol in ethanol) were placed in the inner chamber of the Conway unit; 1 mL of 50% (w/v) potassium carbonate was added to the other side of the outer chamber of the Conway unit. The unit was then sealed, slowly agitated in a horizontal direction to mix the reagents in the outer chamber, and incubated at 37°C for 120 min. Following incubation, the inner chamber of the Conway unit was titrated with 0.02 N sulfuric acid. The VBN concentration was determined as follows:

$$\text{VBN (mg\%)} = (A_1 - A_0) \times F \times 28.014 \times 100 / \text{amount of sample}$$

A<sub>1</sub>: sulfuric acid consumed for sample titration (mL)

A<sub>0</sub>: sulfuric acid consumed for blank titration (mL)

F: standardized index of 0.02 N sulfuric acid

28.014: amount required to consume 1 mL of 0.02 N sulfuric acid

### Statistical analysis


All results in this study are expressed as mean±SD. Analysis of variance was performed using one-way ANOVA and means

were compared using Tukey's multiple range test (significance,  $p < 0.05$ ) with the SPSS/PC Statistics 18.0 program (SPSS Inc., Chicago, USA).

## Results and Discussion

### Total phenolic and flavonoid content

As demonstrated in Fig. 1A, the total phenolic content of RG, AG, and FG significantly increased with increasing amounts



**Fig. 1. Total phenolic (A) and flavonoid (B) content of raw garlic, aged garlic, and fermented garlic extracts.** <sup>a-c</sup> Different letters in the same sample concentration are significantly different ( $p < 0.05$ ). <sup>A-C</sup> Different letters in the same sample group (RG, AG, and FG) are significantly different ( $p < 0.05$ ). Total phenolic content was measured as gallic acid (ppm). Total flavonoid content was measured as quercetin (ppm). RG, raw garlic; AG, aged garlic; FG, fermented garlic.

(2, 6, and 10 mg/mL) of sample ( $p < 0.05$ ). Additionally, the total phenolic content of AG and FG was significantly higher than that of RG ( $p < 0.05$ ). The total phenolic content decreased in the following order: AG > FG > RG ( $p < 0.05$ ).

As demonstrated in Fig. 1B, the total flavonoid content of RG, AG, and FG significantly increased with increasing amounts (2, 6, and 10 mg/mL) of sample ( $p < 0.05$ ). Additionally, the total flavonoid content of AG was significantly higher than that of FG and RG ( $p < 0.05$ ).

Phenolic compounds and flavonoids, such as myricetin, caffeic acid, quercetin, chlorogenic acid, and ferulic acid are representative antioxidants in garlic (Beato et al., 2011) and have strong antioxidant ability owing to their hydroxyl groups (Harborne, 1986). Choi et al. (2008) found that the total phenolic and flavonoid content was increased in aged garlic because of the conversion of compounds to phenolic compounds during high-temperature processing. Fermented black soybean prepared with *B. subtilis* exhibits elevated phenol and flavonoid content (Juan and Chou, 2010). On the other hand, Yang et al. (2012) have reported that *Allium cepa* (onion) fermented with *B. subtilis* did not show a significant increase in flavonoids. Similar results were reported by Lee et al. (2016); garlic fermented with various *Lactobacillus* spp. exhibited only a slight increase in the total phenolic content and flavonoids.

#### **DPPH and ABTS radical scavenging ability**


The DPPH and ABTS radical scavenging activity of RG, AG, and FG significantly increased with increasing amounts (2, 6, and 10 mg/mL) of sample ( $p < 0.05$ ). The DPPH and ABTS radical activities were significantly higher in AG and FG than in RG ( $p < 0.05$ ; Fig. 2).

DPPH and ABTS radical scavenging activities are commonly calculated by measuring the reduction of free radicals by electrons transferred from antioxidants (Blois, 1958). Phenolic compounds and flavonoids are known to have antioxidant abilities such as regenerating  $\alpha$ -tocopherol, scavenging free radicals, and chelating metal ions (Rice-Evans et al., 1996). This is due to aromatic features, conjugated structures with numerous different hydroxyl groups, which make phenolic compounds effective electron or hydrogen atom donors, scavenging free radicals and reactive oxygen species (Zhang and Tsao, 2016). The higher antioxidant power of AG and FG compared to RG might be related to the higher levels of total flavonoids and phenols (Fig. 1).

#### **pH analysis**

The change in pork patty pH values during storage are presented in Fig. 3. The initial pH value was 5.62–5.75 and AG and FG showed a significantly lower pH compared to CON at 0 d ( $p < 0.05$ ). The pH values in all groups significantly decreased with time until 7 d ( $p < 0.05$ ). After 7 d, the pH value of AG and FG continued to significantly decrease until 21 d ( $p < 0.05$ ).


Decreased pH during the storage period has been associated with the activity of lactic acid bacteria (Shin et al., 2017). Garlic is known to contain lactic acid bacteria including *Leuconostoc* spp., *Weissella* spp., and *Lactobacillus* spp. (Jung et al., 2012). These lactic acid bacteria may have also remained in the aged garlic, decreasing pH of AG. In addition, the decrease in the FG pH value might be due to *B. subtilis* fermentation, which uses sugar as a growth substrate and produces acid via pyruvate (Sini et al., 2007). The increase in pH might be due to protein decomposition and amine production by microorganisms in the meat (Biswas et al., 2004). It is assumed that the high antioxidant ability of AG and FG (Fig. 2) effectively inhibited microorganism growth, thus preventing the increase in pH value.




**Fig. 2. DPPH (A) and ABTS (B) radical scavenging activity of raw garlic, aged garlic, and fermented garlic extracts.** <sup>a-c</sup> Different letters in the same sample concentration are significantly different ( $p < 0.05$ ). <sup>A-C</sup> different letters in the same sample group (RG, AG, and FG) are significantly different ( $p < 0.05$ ). DPPH radical scavenging activity is presented as percentage of radical reduced (%). ABTS radical scavenging activity is presented as ascorbic acid ppm. DPPH, 1,1-diphenyl-2-picrylhydrazyl. ABTS, 2'-azino-bis-3-ethylbenzothiazoline-6-sulfonic acid; RG, raw garlic; AG, aged garlic; FG, fermented garlic.

### TBARS

Fig. 4 demonstrates the changes in the pork patty TBARS values during refrigerated storage. The TBARS values of all groups significantly increased with time ( $p < 0.05$ ). AG and FG showed significantly lower TBARS values compared to CON, AA, and RG ( $p < 0.05$ ). Additionally, the final TBARS values of the patties at 21 d were 0.25–0.77 and AG



**Fig. 3.** Changes in the pH value of pork patties with different additives during refrigerated storage at 4°C for 21 d. Error bars indicate the standard deviation. CON (○), patties without treatment; AA (●), patties with 0.01% (w/w) ascorbic acid; RG (▲), patties with 0.5% (w/w) freeze-dried garlic extract; FG (□), patties with 0.5% (w/w) freeze-dried fermented garlic extract; AG (■), patties with 0.5% (w/w) freeze-dried aged garlic extract.



**Fig. 4.** Changes in the TBARS value of pork patties with different additives during refrigerated storage at 4°C for 21 d. Error bars indicate the standard deviation. CON (○), patties without treatment; AA (●), patties with 0.01% (w/w) ascorbic acid; RG (▲), patties with 0.5% (w/w) freeze-dried garlic extract; FG (□), patties with 0.5% (w/w) freeze-dried fermented garlic extract; AG (■), patties with 0.5% (w/w) freeze-dried aged garlic extract. MDA, malondialdehyde.

showed the significantly lowest TBARS value compared to the other groups. The TBARS values increased in the following order: AG<FG<AA<RG<CON ( $p<0.05$ ).



The TBARS assay quantifies the MDA content of meat, a secondary product of lipid oxidation shown to deteriorate meat quality (Choe et al., 2017). It has been reported that a rancid flavor appears in meat products when the TBARS value is more than 0.5 mg MDA/kg meat and that the meat is non-edible if the value reaches 1 mg MDA/kg meat (Park et al., 1988). Ibrahim et al. (2010) reported that various phenolic compounds contained in food additives are strongly correlated with inhibition of lipid oxidation, via free radical neutralization and metal ion chelation. This suggests that the phenolic compounds of AG and FG (Fig. 1A) contributed to the decrease in pork patty TBARS values.


## VBN

The changes in pork patty VBN values during storage are shown in Fig. 5. The initial VBN value at 0 d did not show significant difference between the groups ( $p>0.05$ ). The VBN value of all groups significantly increased with time ( $p<0.05$ ). At 21 d, the VBN values of AG and FG were significantly lower than those of the CON and RG groups ( $p<0.05$ ).

VBN is related to spoilage microorganisms and enzymes in meat products and is an indicator of meat quality (Cai et al., 2015). The main cause of increased VBN values is amino acid decomposition by proteolytic Gram-negative bacteria during storage (Lefebvre et al., 1994). *Allium* species, such as garlic, are known to have an antimicrobial effect (Harris et al., 2001) and phenolic compounds are known to have antimicrobial abilities owing to hydroxyl groups, which reacts with sulfhydryl groups or other protein elements of bacteria (Cowan, 1999). This suggests that the AG and FG groups had a greater antioxidative and antimicrobial capacity because of their phenolic compounds, compared to the CON and RG groups.

## Conclusion

The effect of the antioxidant activity of RG, FG, and AG on pork patty lipid oxidation during refrigerated storage was



**Fig. 5.** Changes in the VBN value of pork patties with different additives during refrigerated storage at 4°C for 21 d. Error bars indicate the standard deviation. CON (○), patties without treatment; AA (●), patties with 0.01% (w/w) ascorbic acid; RG (▲), patties with 0.5% (w/w) freeze-dried garlic extract; FG (□), patties with 0.5% (w/w) freeze-dried fermented garlic extract; AG (■), patties with 0.5% (w/w) freeze-dried aged garlic extract.

investigated. The DPPH and ABTS radical scavenging ability of AG and FG was higher than that of RG. This is due to higher total phenolic content found in AG and FG, which is a representative antioxidant compound in garlic. Furthermore, the pH, TBARS, and VBN values of pork patties containing 0.5% (w/w) of AG or FG were lower than those of the CON and RG groups. As these assays are related to lipid oxidation and the growth of spoilage microorganisms in meat products, we hypothesize that the high antioxidant and antimicrobial power of AG and FG prevented pork patty spoilage, increasing shelf-life. Taken together, these results suggest that AG and FG extracts can be used as natural antioxidative additives for inhibiting pork patty lipid oxidation. As AG has already been used as an additives in food products, further studies on FG are needed to assess the effect of FG extracts on texture, color, and sensory evaluation including customer preference of the products, in order to determine the practicality of using FG as an antioxidant in meat products.

## Conflict of Interest

The authors declare no potential conflict of interest.

## Acknowledgements

The results presented in this study are part of the "Leaders in Industry-University Cooperation +" Project, supported by the Ministry of Education and National Research Foundation of Korea.

## Author Contributions

Conceptualization: Lee CH. Data curation: Lee HJ, Lee CH. Formal analysis: Lee HJ, Lee NY. Methodology: Lee HJ, Yoon DK. Validation: Yoon DK, Lee NY. Investigation: Lee HJ. Writing - original draft: Lee HJ, Lee CH. Writing - review & editing: Lee HJ, Yoon DK, Lee NY, Lee CH.

## Ethics Approval

This article does not require IRB/IACUC approval because there are no human and animal participants.

## References

- Adefegha SA, Oboh G, Olabiy AA. 2018. Nutritional, antioxidant and inhibitory properties of cocoa powder enriched wheat-plantain biscuits on key enzymes linked to type 2 diabetes. *Int Food Res J* 25:793-803.
- Beato VM, Orgaz F, Mansilla F, Montano A. 2011. Changes in phenolic compounds in garlic (*Allium sativum* L.) owing to the cultivar and location of growth. *Plant Foods Hum Nutr* 66:218-223.
- Biswas AK, Keshri RC, Bisht GS. 2004. Effect of enrobing and antioxidants on quality characteristics of precooked pork patties under chilled and frozen storage conditions. *Meat Sci* 66:733-741.
- Blois MS. 1958. Antioxidant determinations by the use of a stable free radical. *Nature* 181:1199-1200.
- Buege JA, Aust SD. 1978. Microsomal lipid peroxidation. In *Methods in enzymology*. Academic Press, New York, NY, USA. pp 302-310.

- Cai L, Cao A, Bai F, Li J. 2015. Effect of  $\epsilon$ -polylysine in combination with alginate coating treatment on physicochemical and microbial characteristics of Japanese sea bass (*Lateolabrax japonicus*) during refrigerated storage. *LWT-Food Sci Technol* 62:1053-1059.
- Chang SS, Ostric-Matijasevic B, Hsieh OAL, Huang CL. 1997. Natural antioxidants from rosemary and sage. *J Food Sci* 42:1102-1106.
- Choe JH, Kim HY, Kim CJ. 2017. Effect of persimmon peel (*Diospyros kaki* Thumb.) extracts on lipid and protein oxidation of raw ground pork during refrigerated storage. *Korean J Food Sci Anim Resour* 37:254-263.
- Choi DJ, Lee SJ, Kang MJ, Cho HS, Sung NJ, Shin JH. 2008. Physicochemical characteristics of black garlic (*Allium sativum* L.). *J Korean Soc Food Sci Nutr* 37:465-471.
- Cowan MM. 1999. Plant products as antimicrobial agents. *Clin Microbiol Rev* 12:564-582.
- Dave D, Ghaly AE. 2011. Meat spoilage mechanisms and preservation techniques: A critical review. *Am J Agric Biol Sci* 6:486-510.
- Han SS, Hur SJ, Lee SK. 2015. A comparison of antioxidative and anti-inflammatory activities of sword beans and soybeans fermented with *Bacillus subtilis*. *Food Funct* 6:2736-2748.
- Harborne JB. 1986. Nature, distribution and function of plant flavonoids. *Prog Clin Biol Res* 213:15-24.
- Harris JC, Cottrell SL, Plummer S, Lloyd D. 2001. Antimicrobial properties of *Allium sativum* (garlic). *Appl Microbiol Biotechnol* 57:282-286.
- Ibrahim HM, Abou-Arab AA, Salem FMA. 2010. Addition of some natural plant extracts and their effects on lamb patties quality. *J Food Technol* 8:134-142.
- Imai J, Ide N, Nagae S, Moriguchi T, Matsuura H, Itakura Y. 1994. Antioxidant and radical scavenging effects of aged garlic extract and its constituents. *Planta Med* 60:417-420.
- Ito N, Hirose M, Fukushima S, Tsuda H, Shirai T, Tatematsu M. 1986. Studies on antioxidants: Their carcinogenic and modifying effects on chemical carcinogenesis. *Food Chem Toxicol* 24:1071-1082.
- Jang EK, Seo JH, Lee SP. 2008. Physiological activity and antioxidative effects of aged black garlic (*Allium sativum* L.) extract. *Korean J Food Sci Technol* 40:443-448.
- Jang HJ, Lee HJ, Yoon DK, Ji DS, Kim JH, Lee CH. 2018. Antioxidant and antimicrobial activities of fresh garlic and aged garlic by-products extracted with different solvents. *Food Sci Biotechnol* 27:219-225.
- Juan MY, Chou CC. 2010. Enhancement of antioxidant activity, total phenolic and flavonoid content of black soybeans by solid state fermentation with *Bacillus subtilis* BCRC 14715. *Food Microbiol* 27:586-591.
- Jung HJ, Hong Y, Yang HS, Chang HC, Kim HY. 2012. Distribution of lactic acid bacteria in garlic (*Allium sativum*) and green onion (*Allium fistulosum*) using SDS-PAGE whole cell protein pattern comparison and 16S rRNA gene sequence analysis. *Food Sci Biotechnol* 21:1457-1462.
- Kumar P, Chatli MK, Mehta N, Malav OP, Verma AK, Kumar D, Rathour M. 2018. Antioxidant and antimicrobial efficacy of sapota powder in pork patties stored under different packaging conditions. *Korean J Food Sci Anim Resour* 38:593-605.
- Kumar Y, Kairam N, Ahmad T, Yadav DN. 2016. Physico chemical, microstructural and sensory characteristics of low-fat meat emulsion containing aloe gel as potential fat replacer. *Int J Food Sci Technol* 51:309-316.
- Lee JB, Joo WH, Kwon GS. 2016. Biological activities of solid-fermentation garlic with lactic acid bacteria. *J Life Sci* 26:446-452.
- Lefebvre N, Thibault C, Charbonneau R, Piette JP. 1994. Improvement of shelf-life and wholesomeness of ground beef by

- irradiation—2. Chemical analysis and sensory evaluation. *Meat Sci* 36:371-380.
- Park BM, Cha SA, Kim HY, Kang DK, Yuan K, Chun H, Chae SW, Kim SH. 2016. Fermented garlic extract decreases blood pressure through nitrite and sGC-cGMP-PKG pathway in spontaneously hypertensive rats. *J Funct Foods* 22:156-165.
- Park BM, Chun H, Chae SW, Kim SH. 2017. Fermented garlic extract ameliorates monocrotaline-induced pulmonary hypertension in rats. *J Funct Foods* 30:247-253.
- Park GB, Kim YJ, Lee HG, Kim JS, Kim YH. 1988. Changes in freshness of meats during postmortem storage, 1; change in freshness of pork. *Korean J Anim Sci* 30:561-566.
- Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. 1999. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic Biol Med* 26:1231-1237.
- Rice-Evans CA, Miller NJ, Paganga G. 1996. Structure-antioxidant activity relationships of flavonoids and phenolic acids. *Free Radic Biol Med* 20:933-956.
- Rose P, Whiteman M, Moore PK, Zhu YZ. 2005. Bioactive S-alk(en)yl cysteine sulfoxide metabolites in the genus *Allium*: The chemistry of potential therapeutic agents. *Nat Prod Rep* 22:351-368.
- Sebranek JG, Sewalt VJ, Robbins KL, Houser TA. 2005. Comparison of a natural rosemary extract and BHA/BHT for relative antioxidant effectiveness in pork sausage. *Meat Sci* 69:289-296.
- Shin DM, Hwang KE, Lee CW, Kim TK, Park YS, Han SG. 2017. Effect of Swiss chard (*Beta vulgaris* var. *cicla*) as nitrite replacement on color stability and shelf-life of cooked pork patties during refrigerated storage. *Korean J Food Sci Anim Resour* 37:418-428.
- Shin JH, Kang MJ, Kim RJ, Sung NJ. 2011. The quality characteristics of sausage with added black garlic extracts. *Korean J Food Cook Sci* 27:701-711.
- Simoës GD, Rychecki-Hecktheuer LH, Hubscher GH, Boligon AA. 2018. Quantification of bioactive compounds in crem (*Tropaeolum pentaphyllum* Lam) tubers: Fibers, phenolic compounds and evaluation of its antioxidant activity. *Int Food Res J* 25:1315-1321.
- Sini TK, Santhosh S, Mathew PT. 2007. Study on the production of chitin and chitosan from shrimp shell by using *Bacillus subtilis* fermentation. *Carbohydr Res* 342:2423-2429.
- Steinkraus KH. 1997. Classification of fermented foods: Worldwide review of household fermentation techniques. *Food Control* 8:311-317.
- Wang SJ, Lee JH, Choi MJ, Lee SK. 2012. Effects of aged black garlic extracts on the rheology of flour dough. *J Korean Soc Food Sci Nutr* 41:430-435.
- Yang EJ, Kim SI, Park SY, Bang HY, Jeong JH, So JH, Rhee IK, Song KS. 2012. Fermentation enhances the *in vitro* antioxidative effect of onion (*Allium cepa*) via an increase in quercetin content. *Food Chem Toxicol* 50:2042-2048.
- You BR, Kim HR, Kim MJ, Kim MR. 2011. Comparison of the quality characteristics and antioxidant activities of the commercial black garlic and lab-prepared fermented and aged black garlic. *J Korean Soc Food Sci Nutr* 40:366-371.
- Yun CH. 2017. Method for preparing food or pharmaceutical composition comprising fermented extract of garlic sprout as an effective ingredient. KR Patent 1020170099631.
- Zahid MA, Seo JK, Park JY, Jeong JY, Jin SK, Park TS, Yang HS. 2018. The effects of natural antioxidants on protein oxidation, lipid oxidation, color, and sensory attributes of beef patties during cold storage at 4°C. *Korean J Food Sci Anim Resour* 38:1029-1042.
- Zhang H, Tsao R. 2016. Dietary polyphenols, oxidative stress and antioxidant and anti-inflammatory effects. *Curr Opin Food*

Sci 8:33-42.

Zhang Y, Liu J, Lu X, Zhang H, Wang L, Guo X, Qi X, Qian H. 2014. Isolation and identification of an antioxidant peptide prepared from fermented peanut meal using *Bacillus subtilis* fermentation. Int J Food Prop 17:1237-1253.