



SHORT COMMUNICATION

Macroscopic, Histological, and Microbiological Characterization of Contact Lesions at the Tibiotarsal Region of Broilers

Ricardo Cavani^{1,*}, Marcela da Silva Rubio¹, Khauston Augusto Pereira Alves², Lucas José Luduverio Pizauro¹, Marita Vedovelli Cardozo³, Paulo Lourenço Silva⁴, Iran José Oliveira Silva⁵, and Fernando Antônio Avila¹

OPEN ACCESS

Received August 25, 2021

Revised November 12, 2021

Accepted December 9, 2021

*Corresponding author : Ricardo Cavani
Department of Veterinary Pathology, School
of Agricultural and Veterinary Sciences,
São Paulo State University (UNESP),
Jaboticabal, São Paulo 14884-900, Brazil
Tel: +55-16-3209-7100
E-mail: ricardocavani@hotmail.com

*ORCID

Ricardo Cavani
<https://orcid.org/0000-0002-8984-2372>
Marcela da Silva Rubio
<https://orcid.org/0000-0003-1389-4130>
Khauston Augusto Pereira Alves
<https://orcid.org/0000-0002-2142-7148>
Lucas José Luduverio Pizauro
<https://orcid.org/0000-0002-1071-8740>
Marita Vedovelli Cardozo
<https://orcid.org/0000-0003-3972-0198>
Paulo Lourenço Silva
<https://orcid.org/0000-0001-8186-8693>
Iran José Oliveira Silva
<https://orcid.org/0000-0002-4416-8433>
Fernando Antônio Avila
<https://orcid.org/0000-0002-9779-2213>

¹Department of Veterinary Pathology, School of Agricultural and Veterinary Sciences, São Paulo State University (UNESP), Jaboticabal, São Paulo 14884-900, Brazil

²Cesumar University, Maringa, Paraná, Brazil

³Minas Gerais State University, Passos, Minas Gerais, Brazil

⁴Uberlandia Federal University, Uberlandia, Brazil

⁵Department of Rural Engineering, São Paulo University, Piracicaba, São Paulo, Brazil

Abstract Brazil is considered as a great broiler feet exporter, especially for the Chinese trade. Contact lesions at the tibiotarsal region are responsible for economic losses and there is no model for its classification, thereby this study presents a fast and practical grade system to be used in the poultry industry and proposes these lesion characterizations into three different grades. For this, correlation was made between macroscopic, histological findings and microbiological quantification (*Escherichia coli*, *Staphylococcus* spp., *Streptococcus* spp. and sulphite-reducing clostridia) from contact lesions in the tibiotarsal region of 112 broiler carcasses, divided in four groups (n=28), accordingly to the lesion's intensity. There were no significant differences in microbiological quantification among the groups (p>0.05) except for the grade 3 group, as grade 1 and 2 lesions were in the early stages and histopathological changes such as ulceration were not observed. In grade 3 lesion group, it was observed bacterial cocci grume and ulceration at the articular region and significantly higher microbiological count (p<0.05) for *E. coli* and *Staphylococcus* spp. In conclusion, the visual standard proposed in this work, correlated and confirmed by the histopathologic, and microbiologic characterization, allows to precise and fast ascertainment of the contact lesion grade in the tibiotarsal regions of broiler carcasses. Moreover, it should be highlighted that grades 1 and 2 alterations are not caused by an inflammatory process caused by pathogenic agents and should not be considered a public health risk.

Keywords hock burn, dermatitis, slaughterhouse, carcass condemnation, inspection line

Introduction

Brazil is an important broiler feet exporter to the Asian trade, especially China and

Hong Kong, which consider this product as a delicacy in their gastronomy and as collagen supplement for the cosmetic and medicine industry (ApexBrasil, 2018; Munasinghe et al., 2014). Moreover, it is considered in many countries, as well as in Brazil, as the third cut of chicken with the highest economic value (Chen et al., 2016).

Among the main factors for the condemnation of broiler feet are the contact lesions and keratosis. The contact lesions in the tibiotarsal region, also known as hock burn, contact dermatitis and dermatitis are characterized by the presence of blackish or brownish coloration on the skin of the chickens' tibiotarsal region (Bessei, 2006). Besides the economic losses due to discards, it also impairs the broilers performance and violates the technical recommendation of animal welfare in poultry production, because it results in pain for the bird, evidenced by the slowly locomotion or even reluctance to move (Louton et al., 2020).

It is known that the weight is a risk factor that can be associated with the occurrence of lesions on broiler legs (Louton et al., 2018) and that it will simultaneously acts alongside the bad litter quality and the presence of infectious agents (Thøfner et al., 2019). The litter of the poultry house is used to provide a better life quality for the birds since it contributes to the thermal comfort and avoid direct contact with the floor or tread, preventing the formation of calluses in the animal's leg and breast. Besides the litter absorbs and incorporates the waste, such as excrete, desquamation, feathers and food and water residues that fall from the feeders and drinking fountains (Avila et al., 2008). According to Heidemann Olsen et al. (2018), the high humidity and high ammonia concentrations originated from the accumulated faecal matter result in ammonia gas liberation that causes chemical burns and weakening of the dermis. In addition, the humidity makes the external dermis softens, facilitating the entry and proliferation of microorganisms.

Although a visual classification system for contact lesion exists (Michel et al., 2012), a histopathological validation for this assessment scheme was just recently described (Louton et al., 2020). Besides, until now, only the correspondence between the macroscopic aspects of the lesions and their histopathological characteristic was demonstrated with no correlation between their microbiological content. Nevertheless, since no standard visual method was developed to compare this kind of lesion, this study aimed to present a fast and practical visual grade to be used in the industry and validated with the histological findings and microbiological quantification in the tibiotarsal region in broiler carcasses.

Materials and Methods

Experimental facilities

This study was conducted in a commercial slaughterhouse located in the Sao Paulo state, Brazil, with a slaughter capacity of 120.000 birds/day. The birds came from 14 producers of the Northern regions of the Sao Paulo state. The bird batches were of the Cobb and Ross linages, with a mean of 42 days of age and 2.8 kg of weight at the slaughter.

Experimental design

For this study, 112 broiler carcasses were obtained at the post-mortem inspection line (pre-inspection) with or without visual characteristics of contact lesion at the tibiotarsal region. Then, carcasses were removed from the inspection line and with the aid of a disinfect knife in alcohol 70%, the leg was removed by disarticulating the tibiofemoral region, followed by the disarticulation and removal of the metatarsophalangeal region. Each sample was constituted by a single joint, having collected 28 articulations for each evaluated grade (n=28), which were packed either in sterile bag under refrigeration or flasks containing 10% formaldehyde and, thus, sent to the laboratory for analysis.

Visual classification

For evaluation and visual classification of the tibiotarsal joint, to attribute a macroscopic score of the lesions, was carried out over a period of six months and the lesions were divided into four different groups: a control group (CG) and grades 1 (G1), 2 (G2) and 3 (G3). It was used a total of 942 (CG=234, G1=236, G2=235, and G3=237) pictures of tibiotarsal region observed in the post-mortem inspection line to perform macroscopic classification. The grading methods was based on the principals described by Welfare-Quality® (Welfare Quality Consortium, 2009) and Louton et al. (2020), utilizing fewer lesion grades (three instead of four) that were created based on characteristics such as intensity and color of the contact lesion in the tibiotarsal region and the presence or absence of scarification.

Histopathological evaluation

The samples harvested for histopathological examination were placed in flasks containing 10% formalin and sent to the “Centro de Diagnóstico de Sanidade Animal” (CEDISA, Concordia, SC, Brazil). The histological sections were stained with hematoxylin and eosin for evaluation of the epidermis, dermis and hypodermis (subcutaneous) of the tibiotarsal region.

Microbiological evaluation

The microbiological analysis were performed at the “Laudo Laboratório Avícola Uberlândia Ltda” (LAUDO, Uberlandia, MG, Brazil). From the refrigerated samples for microbiological count, sterile swabs were scraped on a 5 cm² area of the articular surface. After sampling, the swabs were placed in tubes containing 5 mL of 1% buffered peptone water solution (Neogen, Lansing, MI, USA). After sampling, *E. coli*, *Staphylococcus* spp. and sulphite-reducing clostridia were quantified using 3M™ Petrifilm™ Plates (3M do Brazil, Sumaré, Brazil), and *Streptococcus* spp. used Blood agar with 5% sheep blood plates (Laborclin, Pinhais, PR, Brazil) as described by Chadfield et al. (2004).

Statistical analysis

Microbiological data were subjected to statistical analysis using the R-Project for Statistical Computing (Wilson and Norden, 2015) and Python for Data Analysis software (Tobergte and Curtis, 2013). Microbiological counts were logarithmic transformed and after not passing the normality test (Shapiro-Wilk), the results were evaluated with a non-parametric Kruskal-Wallis test and statistical significance of the groups were evaluated by the paired sample Wilcoxon test and considered statistically significant when the p-value was lower than 5% ($p < 0.05$).

Results and Discussion

Considering the principals described by Welfare-Quality® (Welfare Quality Consortium, 2009) and Louton et al. (2020) and the characteristics about the intensity and coloration of the contact lesions at the tibiotarsal region, the lesions were allocated in four different grades: The CG the articulation without visual alteration; Grade 1 (G1), refers to the joint with mild scarification and pink or reddish color (erythema); Grade 2 (G2), refers to the joint with moderate scarification, crust formation and brownish coloration; Grade 3 (G3) refers to the articulation with scarification of severe intensity, presence of crust, ulceration and/or brownish and/or purplish and/or blackish coloration (Fig. 1).

The classification of contact injuries prove to be widely diversified, both nationally and internationally. Some of them focus on the size of the lesion (Arnould et al., 2009; Pagazaurtundua and Warriss, 2006a; Pagazaurtundua and Warriss,

2006b; Welfare Quality Consortium, 2009) whereas others combine size and depth (Allain et al., 2009; Ekstrand et al., 1998; Kaukonen et al., 2016; Kjaer et al., 2006) in different categories. Therefore, the differences of this varieties of assessment systems may not well elucidated and has been a challenge for both scientific point of view and the quality of surveillance systems (Michel et al., 2012; Riber et al., 2020). Thus, it was suggested herein a model for these gross lesions classifications to further be used in accordance with histological and microbiological analysis.

In the histopathological characterization was described that in the CG articulation was not observed noteworthy lesions. In the G1 and G2 groups, it was observed histological alterations only in the superficial regions of the skin, whereas in G1, there were a moderate proliferation of subcutaneous connective tissue, with no signs of inflammation. Moreover, in G2 was observed a moderate proliferation of connective tissue, with a mild ulceration focus of the skin and formation of a keratin crust and degenerated inflammatory cells, considered as a morphological diagnose of mild focal necrotic dermatitis.

In G3 were observed ulceration in the epidermis and dermis, with the presence of crusts of necrotic tissue, keratin, degenerated inflammatory cells and clumps of bacteria in the form of cocci. In the subcutaneous (hypoderm), it was observed proliferation of connective tissue, with deposition of fibrin lump and, mild presence of hemorrhage and inflammatory infiltrate, with the heterophile predominance. These characteristics are considered a morphological find of focally extensive necrotic dermatitis. The inflammation severity of the lesions increased with the crescent macroscopic score, in which the visually more severe and deeper lesions, also presented more accentuated histological findings, especially considering the



Fig. 1. Macroscopic classification of the contact lesion grades at the tibiotarsal region of broilers. Control group, no visible alteration; Grade 1, mild scarification and pinkish or reddish (erythema) coloration; Grade 2, moderate scarification, crust formation and brownish coloration; Grade 3, severe scarification, presence of crust, ulceration and brownish and/or purplish and/or blackish coloration.

presence of inflammatory cell and remnants of inflammation (proliferation of connective tissue and presence of fibrin; Martins et al., 2016).

The microbiological quantification results, described in Table 1, corroborates with the histopathological characterization, in which no inflammatory reaction of bacterial origin was observed in samples from CG and G1 and G2 and bacterial cocci clumps in G3 samples. In the evaluated samples was not evidenced the presence of sulfite-reducing clostridia, and the counts of *E. coli*, *Staphylococcus* spp. and *Streptococcus* spp. of the CG and grade 1 and 2 were within the acceptance threshold required by Brazil and importing trades such as the USA, China, Eurasian Economic Union, Saudi Arabia, South Africa, and others [Agri-Food and Veterinary Authority of Singapore (AVA), 2000; European Commission (EC), 2005; GB National Standards of People's Republic of China, 2014; GCC Standardization Organization (GSO), 2014; IBNORCA, 2019; Ministry of Agriculture, Livestock and Supply, 2017; Ministry of Agriculture, Livestock and Supply, 2019]. By statistical tests, when comparing the results of the CG with the other grades, it was not observed significant differences ($p>0.05$), except in G3, which differed from the CG and other grades.

Although there is an extensive repertoire of information about the predisposing factors for the occurrence of contact injuries in tibiotarsal joints of broiler, especially those related to bedding quality, nutrition, and management, little is known about the microbiological factors associated with these conditions and that could be considered inadequate (Heidemann Olsen et al., 2018). Chadfield et al. (2004) suggested that *E. coli*, *Staphylococcus* spp. and *Streptococcus* spp. are opportunistic bacteria that can prevent from the bird's own organism, which would explain the similarity ($p>0.05$) in bacterial counts observed for G1 and G2 when compared to the CG.

In the case of mild lesions (G1), in which the skin was preserved intact and in the moderate lesions (G2), where only the superficial skin layer was compromised, there was no microbial invasion into the tissues, which Allain et al. (2009) characterize them as “chemical burns” that are the results of the high amount of humidity of the litter and/or high concentration of ammonia in the fecal matter. In contrast, the accentuated skin deterioration process allow for the opportunistic microorganism to invade the subcutaneous tissue (Nagase et al., 2002). That fact was observed in the present study in lesions of G3, in which there was the presence of ulceration and increased bacterial counts, as described by Martins et al. (2016) who observed the presence of areas of necrosis and the presence of inflammatory infiltrate in severe contact lesions.

Thus, the classification of lesions into three scores of hock dermatitis proposed in this work by means of macroscopic, histopathological and microbiological findings provided a simple, practical and scientific-based way for its use both in the poultry industry, as well as in poultry farms in Brazil. Such grouping allows for easier identification, in addition to enabling better characterization of the evolution of changes in the animals, a fact evidenced both by histopathological findings

Table 1. Microbiological quantification of *Escherichia coli*, *Staphylococcus* spp. and *Streptococcus* spp. in contact lesions in the tibiotarsal joint of broiler chickens

Grade	Microorganism (CFU/mL)		
	<i>E. coli</i>	<i>Staphylococcus</i> spp.	<i>Streptococcus</i> spp.
Control	$4.70 \times 10^{1,a}$	$2.73 \times 10^{2,a}$	$1.80 \times 10^{3,a}$
Grade 1	$9.30 \times 10^{1,a}$	$4.40 \times 10^{2,a}$	$1.16 \times 10^{3,a}$
Grade 2	$1.50 \times 10^{2,a}$	$8.10 \times 10^{2,a}$	$1.11 \times 10^{3,a}$
Grade 3	$4.50 \times 10^{4,b}$	$2.60 \times 10^{5,b}$	$4.80 \times 10^{4,a}$

^{a,b} Means followed by different letters in the columns, differ from each other by the Kruskal-Wallis and the Wilcoxon test at 5% probability ($p<0.05$).

(presence of connective tissue, inflammatory infiltrate, ulceration and the formation of keratin crust) and by microbiological findings (increasing microorganisms quantification).

Conclusion

The visual standard proposed in this work, which was correlated and confirmed by the histopathological and microbiological characterization, allows a faster and precise ascertainment of the degrees of contact lesions in tibiotarsal joints of broilers. Furthermore, it is noteworthy that grades 1 and 2 alterations are not due to an inflammatory process caused by pathogens and, therefore, do not represent a public health concern.

Conflicts of Interest

The authors declare no potential conflicts of interest.

Author Contributions

Conceptualization: Cavani R. Data curation: Cavani R, Rubio MS, Alves KAP, Pizauro LJJ. Formal analysis: Cavani R, Rubio MS, Alves KAP, Pizauro LJJ. Methodology: Cavani R. Investigation: Cavani R, Rubio MS, Alves KAP, Pizauro LJJ. Writing - original draft: Cavani R, Rubio MS, Alves KAP, Pizauro LJJ. Writing - review & editing: Cavani R, Rubio MS, Alves KAP, Pizauro LJJ, Cardozo MV, Silva PL, Silva IJO, Avila FA.

Ethics Approval

The experiment was approved by the Ethics Committee on the Use of Animals (CEUA) under protocol no. 1183/21, located at São Paulo State University (UNESP), School of Agricultural and Veterinary Sciences, Jaboticabal, São Paulo, Brazil.

References

- Agri-Food and Veterinary Authority of Singapore [AVA]. 2000. Circular/Directive no. 2000/1 – Microbiological specifications for imported meat. AVA, Singapore.
- Allain V, Mirabito L, Arnould C, Colas M, Le Bouquin S, Lupo C, Michel V. 2009. Skin lesions in broiler chickens measured at the slaughterhouse: Relationships between lesions and between their prevalence and rearing factors. *Br Poult Sci* 50:407-417.
- ApexBrasil. 2018. China trade and investment opportunities: Food and beverage. Available from: <https://www.cebc.org.br/2019/03/28/opportunidades-de-comercio-e-investimento-na-china-alimentos-e-bebidas/>. Accessed at Aug 11, 2021.
- Arnould C, Butterworth A, Knierim U. 2009. Standardization of clinical scoring in poultry. In *Assessment of animal welfare measures for layers and broilers*. Forkman B, Keeling L (ed). Cardiff University, Cardiff, UK. pp 7-30.
- Avila VS, Oliveira U, Figueiredo EAP, Costa CAF, Abreu VMN, Rosa PS. 2008. Evaluation of alternative materials to replace wood shavings as aviary bedding. *Rev Bras Zootec* 37:273-277.

- Bessei W. 2006. Welfare of broilers: A review. *Worlds Poult Sci J* 62:455-466.
- Chadfield MS, Christensen JP, Christensen H, Bisgaard M. 2004. Characterization of streptococci and enterococci associated with septicaemia in broiler parents with a high prevalence of endocarditis. *Avian Pathol* 33:610-617.
- Chen J, Tellez G, Escobar J. 2016. Identification of biomarkers for footpad dermatitis development and wound healing. *Front Cell Infect Microbiol* 6:26.
- Ekstrand C, Carpenter TE, Andersson I, Algiers B. 1998. Prevalence and control of foot-pad dermatitis in broilers in Sweden. *Br Poult Sci* 39:318-324.
- European Commission [EC]. 2005. Microbiological criteria for foodstuffs. Commission Regulation (EC) No 2073/2005. European Union, Brussels, Belgium.
- GB National Standards of People's Republic of China. 2014. National food standard fresh and frozen poultry products. GB 16869-2005. GB National Standards of People's Republic of China, Beijing, China.
- GCC Standardization Organization [GSO]. 2014. Microbiological criteria for foodstuffs. GSO/FDS 1016/2014. GSO, Riyadh, Saudi Arabia.
- Heidemann Olsen R, Christensen H, Kabell S, Bisgaard M. 2018. Characterization of prevalent bacterial pathogens associated with pododermatitis in table egg layers. *Avian Pathol* 47:281-285.
- Instituto Boliviano de Normalización y Calidad [IBNORCA]. 2019. Carnes y derivados – Poultry meat - Microbiological Requirements. NB:310013. IBNORCA, La Paz, Bolivia.
- Kaukonen E, Norring M, Valros A. 2016. Effect of litter quality on foot pad dermatitis, hock burns and breast blisters in broiler breeders during the production period. *Avian Pathol* 45:667-673.
- Kjaer JB, Su G, Nielsen BL, Sørensen P. 2006. Foot pad dermatitis and hock burn in broiler chickens and degree of inheritance. *Poult Sci* 85:1342-1348.
- Louton H, Bergmann S, Reese S, Erhard M, Bachmeier J, Rösler B, Rauch, E. 2018. Animal- and management-based welfare indicators for a conventional broiler strain in 2 barn types (Louisiana barn and closed barn). *Poult Sci* 97:2754-2767.
- Louton H, Piller A, Bergmann S, Erhard M, Stracke J, Spindler B, Kemper N, Schmidt P, Schade B, Boehm B, Kappe E, Bachmeier J, Schwarzer A. 2020. Histologically validated scoring system for the assessment of hock burn in broilers. *Avian Pathol* 49:230-242.
- Martins BB, Martins MRFB, Mendes AA, Fernandes BCS, Aguiar EF. 2016. Footpad dermatitis in broilers: Differences between strains and gender. *Rev Bras Cienc Avic* 18:461-466.
- Michel V, Prampart E, Mirabito L, Allain V, Arnould C, Huonnic D, Le Bouquin S, Albaric O. 2012. Histologically-validated footpad dermatitis scoring system for use in chicken processing plants. *Br Poult Sci* 53:275-281.
- Ministry of Agriculture, Livestock and Supply. 2017. Memorandum n° 36/2017/DIPOA/MAPA/SDA/MAPA. Ministry of Agriculture, Livestock and Supply, Brasilia, Brazil.
- Ministry of Agriculture, Livestock and Supply. 2019. Official notice n° 33/2019/CGCOA/DIPOA/SDA/MAPA. Ministry of Agriculture, Livestock and Supply, Brasilia, Brazil.
- Munasinghe KA, Schwarz JG, Nyame AK. 2014. Chicken collagen from law market value by-products as an alternate source. *J Food Process* 2014:298295.
- Nagase N, Sasaki A, Yamashita K, Shimizu A, Wakita Y, Kitai S, Kawano J. 2002. Isolation and species distribution of staphylococci from animal and human skin. *J Vet Med Sci* 64:245-250.
- Pagazaurtundua A, Warriss PD. 2006a. Measurements of footpad dermatitis in broiler chickens at processing plants. *Vet Rec*

158:679-682.

- Pagazaurtundua A, Warriss PD. 2006b. Levels of foot pad dermatitis in broiler chickens reared in 5 different systems. *Br Poult Sci* 47:529-532.
- Riber AB, Rangstrup-Christensen L, Hansen MS, Hinrichsen LK, Herskin MS. 2020. Characterisation of footpad lesions in organic and conventional broilers. *Animal* 14:119-128.
- Thøfner ICN, Poulsen LL, Bisgaard M, Christensen H, Olsen RH, Christensen JP. 2019. Correlation between footpad lesions and systemic bacterial infections in broiler breeders. *Vet Res* 50:38.
- Tobergte DR, Curtis S. 2013. *Python for data analysis*. O'Reilly, Sebastopol, CA, USA.
- Welfare Quality Consortium. 2009. *Welfare Quality® assessment protocol for poultry*. Welfare Quality Consortium, Lelystad, The Netherlands. p 114.
- Wilson A, Norden N. 2015. *The R Project for statistical computing*. The R Foundation, Vienna, Austria.