

REVIEW

Analytical Methods and Effects of Bioactive Peptides Derived from Animal Products: A Mini-Review

Jae Won Jeong^{1,†}, Seung Yun Lee², Da Young Lee¹, Jae Hyeon Kim¹,
 Seung Hyeon Yun¹, Juhyun Lee¹, Ermie Jr. Mariano¹, Sung Sil Moon^{3,†},
 and Sun Jin Hur^{1,*}

¹Department of Animal Science and Technology, Chung-Ang University, Anseong 17546, Korea

²Division of Animal Science, Division of Applied Life Science (BK21 Four), Institute of Agriculture & Life Science, Gyeongsang National University, Jinju 52828, Korea

³Sunjin Technology & Research Institute, Icheon 17332, Korea


OPEN ACCESS

Received February 23, 2024

Revised April 4, 2024

Accepted April 9, 2024

***Corresponding author** : Sun Jin Hur
 Department of Animal Science and
 Technology, Chung-Ang University,
 Anseong 17546, Korea
 Tel: +82-31-670-4673
 Fax: +82-31-670-3108
 E-mail: hursj@cau.ac.kr

***ORCID**

Jae Won Jeong
<https://orcid.org/0000-0001-5240-1875>
 Seung Yun Lee
<https://orcid.org/0000-0002-8861-6517>
 Da Young Lee
<https://orcid.org/0000-0002-3172-0815>
 Jae Hyeon Kim
<https://orcid.org/0000-0003-1174-4737>
 Seung Hyeon Yun
<https://orcid.org/0000-0002-9940-2960>
 Juhyun Lee
<https://orcid.org/0000-0001-6777-4447>
 Ermie Jr. Mariano
<https://orcid.org/0000-0003-2630-4603>
 Sung Sil Moon
<https://orcid.org/0000-0003-2734-8931>
 Sun Jin Hur
<https://orcid.org/0000-0001-9386-5852>

[†] These authors contributed equally to this work.

Abstract Peptides with bioactive effects are being researched for various purposes. However, there is a lack of overall research on pork-derived peptides. In this study, we reviewed the process of obtaining bioactive peptides, available analytical methods, and the study of bioactive peptides derived from pork. Pepsin and trypsin, two representative protein digestive enzymes in the body, are hydrolyzed by other cofactors to produce peptides. Bicinchoninic acid assay, sodium dodecyl sulfate-polyacrylamide gel electrophoresis, chromatography, and *in vitro* digestion simulation systems are utilized to analyze bioactive peptides for protein digestibility and molecular weight distribution. Pork-derived peptides mainly exhibit antioxidant and antihypertensive activities. The antioxidant activity of bioactive peptides increases the accessibility of amino acid residues by disrupting the three-dimensional structure of proteins, affecting free radical scavenging, reactive oxygen species inactivation, and metal ion chelating. In addition, the antihypertensive activity decreases angiotensin II production by inhibiting angiotensin converting enzyme and suppresses blood pressure by blocking the AT1 receptor. Pork-derived bioactive peptides, primarily obtained using papain and pepsin, exhibit significant antioxidant and antihypertensive activities, with most having low molecular weights below 1 kDa. This study may aid in the future development of bioactive peptides and serve as a valuable reference for pork-derived peptides.

Keywords pork, bioactive peptide, angiotensin converting enzyme inhibitory peptide, antioxidative peptide

Introduction

In general, peptides have a smaller molecular structure than proteins, consisting of 2–50 amino acids. Certain peptides play a role in regulating the activity of other molecules. Bioactive peptides consist of 2–20 amino acids and have a relatively small molecular weight compared to proteins (Lafarga and Hayes, 2014). The market for

bioactive peptides is expanding with the growth of functional food and beverage products, and they are widely applied in functional foods, natural health products, health foods, and cosmetics (Chalamaiah et al., 2019). This growth can be attributed to the fact that consumers are becoming more health-conscious, and industries are utilizing functional ingredients to develop new products. Bioactive peptides are found in food proteins, especially in milk, meat, fish, and legumes. They are utilized as ingredients in functional foods and pharmaceuticals due to their beneficial effects on the human digestive, endocrine, nervous, and cardiovascular systems, among others, and their role in health (Heres et al., 2021a; Sánchez and Vázquez, 2017). The efficacy of bioactive peptides is often determined by their molecular weight and amino acid sequence because the amino acids that comprise the peptide sequence can have varying properties and effects. Livestock-derived bioactive peptides have been reported to have antioxidant, antihypertensive, antithrombotic, and antimicrobial activities, which have positive effects on disease prevention and blood circulation (Aluko, 2015; Kim et al., 2024; Lafarga and Hayes, 2014; Rubak et al., 2022). Previous studies have confirmed that bioactive peptides that modulate various biological actions can be obtained from pork (Arihara, 2006). In particular, antioxidant and antimicrobial active peptides isolated from pork muscle proteins provide important health benefits to humans and can be utilized as functional ingredients in foods (Di Bernardini et al., 2011). However, the effective utilization of pork-derived bioactive peptides has not been adequately studied. Therefore, in this study, we introduced a method for protein digestion analysis that can be utilized to obtain peptides and categorize various potencies and types of bioactive peptides derived from pork.

Process of Protein Digestion by Enzymes

Proteins consumed by humans must be hydrolyzed by proteolytic enzymes secreted by the stomach, pancreas, and small intestine in order to be digested and absorbed by the body. After proteins digestion, peptides present in the intestinal lumen typically consist of 2–6 amino acids, which account for about 80% of the total amino acids (Bhutia and Ganapathy, 2018). In the intestinal lumen, the amount of amino acids present in peptides is higher than that of free amino acids (Adibi and Mercer, 1973). Consequently, most peptides and free amino acids are transported across the intestinal epithelium into the digestive tract through the brush border membrane transport system. The majority of peptides are then hydrolyzed to free amino acids, which make up about 90% of the total amino acids (Bhutia and Ganapathy, 2018).

Ingested proteins are broken down by a variety of enzymes secreted by the body's digestive system. The initial step in this process is performed by pepsin, a proteolytic enzyme secreted by the stomach. Pepsin is initially secreted as an inactive precursor, called pepsinogen, which is produced by the chief cells of the stomach (Gupta, 2018). This inactive precursor, pepsinogen, is then activated through an autocatalytic reaction in the acidic pH environment of the stomach, resulting in the production of pepsin (Gupta, 2018). Protein hydrolysates processed by pepsin are mostly in the form of polypeptides, and only a small amount of free amino acids is released through hydrolysis (Hinsberger and Sandhu, 2004). After undergoing digestion in the stomach, gastric contents pass through the duodenum and jejunum, where they stimulate cells in the intestinal mucosa to produce cholecystokinin (Liddle, 1997). Cholecystokinin then triggers the secretion of pancreatic juice, which is rich in proteolytic enzymes, and causes the gallbladder to contract and release bile (Liddle, 1997). In addition, when the gastric contents reach the small intestine, the acidic pH environment created by gastric acid prompts S-cells in the duodenum to release secretin, which is produced by these cells (DiGregorio and Sharma, 2019). Secretin increases the secretion of bicarbonate from the pancreas and biliary tract. This neutralizes the acidic pH environment in the duodenum caused by stomach acid to a pH level of around 6–8 and reduces the secretion of stomach acid (Bhutia and Ganapathy, 2018; DiGregorio and Sharma, 2019).

The pancreas is a vital digestive organ that produces and secretes proteolytic enzymes into the small intestine to digest ingested protein. The major pancreatic proteolytic enzymes have been identified as trypsin, chymotrypsin, elastase, and carboxypeptidase (Whitcomb and Lowe, 2007). Similar to pepsin, these enzymes are initially secreted as inactive precursors, including trypsinogen, chymotrypsinogen, proelastase, and procarboxypeptidase (Whitcomb and Lowe, 2007). Among them, trypsinogen is first activated to trypsin by enteropeptidase in the small intestine. Activated trypsin then acts on chymotrypsinogen, proelastase, and procarboxypeptidase to form active chymotrypsin, elastase, and carboxypeptidase (Bhutia and Ganapathy, 2018). Trypsin is highly reactive towards peptides containing the basic amino acids arginine and lysine, while chymotrypsin hydrolyzes peptides containing the aromatic amino acids tyrosine, phenylalanine, and tryptophan (Whitcomb and Lowe, 2007). In addition, elastase acts on peptide binding sites formed by the non-polar amino acids glycine and alanine (Whitcomb and Lowe, 2007). Consequently, the proteolytic enzymes in the pancreas hydrolyze proteins that are mostly present in polypeptide form during small intestinal digestion into oligopeptides and free amino acids consisting of 6–8 amino acids (Bhutia and Ganapathy, 2018). These oligopeptides are then hydrolyzed into smaller forms of peptides, such as tripeptides and dipeptides, by brush border peptidases found in the microvilli composed of small intestinal enterocytes (Hooton et al., 2015). Finally, the myriad of tripeptides and dipeptides are absorbed into the small intestine where they are broken down into amino acids by cytoplasmic peptidases and released into the bloodstream (Boron and Boulpaep, 2016).

Analysis Methods for Protein Digestibility

Methods such as the bicinchoninic acid (BCA) assay, sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), gel permeation chromatography (GPC), and *in vitro* digestion system have been utilized to analyze protein digestibility and molecular weight distribution, which are relevant for peptide acquisition (Li et al., 2017; Rezvankhah et al., 2021; Wen et al., 2015).

The BCA assay is a highly sensitive method for quantifying proteins by comparing their chromogenic reactions. The principle behind this method is that Cu^{2+} ions are reduced to Cu^+ ions by peptide binding of proteins in an alkaline environment. The Cu^+ ions then combine with BCA to form a purple complex. This assay is similar to the Lowry assay but has the advantage of being relatively simple and resistant to compounds that may interfere with the results (Walker, 2009). Once the reaction is complete, the complex can be analyzed using a spectrophotometer to measure the amount of protein at the maximum absorption wavelength of 562 nm (He, 2011). This reaction is mainly influenced by the presence of four amino acid residues (cysteine, cystine, tyrosine, and tryptophan) in the protein molecule (Fischer et al., 1999). Therefore, the BCA assay can be utilized to determine the initial protein content of a sample.

SDS-PAGE is an electrophoresis technique that analyzes the movement of charged protein molecules in an electric field. It is commonly used to separate proteins by size and analyze them qualitatively (Rajput and Sharma, 2011; Roy et al., 2012). SDS, an anionic surfactant with a strong protein denaturing effect, binds to proteins at a constant rate. During this process, the proteins are transformed into a linear chain structure and become negatively charged (Farrell, 2009; Rajput and Sharma, 2011). The proteins then move in the electric field containing the polyacrylamide gel according to their molecular size, with smaller proteins migrating and separating faster than larger proteins (Rajput and Sharma, 2011). Therefore, SDS-PAGE separates polypeptides based on molecular size, making it the best experimental method for analyzing protein digestibility as a function of protein molecular weight (Righetti, 2005). Li et al. (2017) and Wen et al. (2015) used SDS-PAGE to screen for changes in molecular weight and compare the digestibility of pork protein before and after *in vitro* digestion. Specifically, Li

et al. (2017) compared the molecular weight of four types of pork proteins (cooked, emulsion-type sausage, dry-cured, and stewed) before digestion, after pepsin digestion, and after pepsin and trypsin digestion. Additionally, Wen et al. (2015) compared the differences in molecular weight of proteins from four types of cooked meat (pork, beef, chicken, and fish) before digestion, after pepsin digestion, and after pepsin and trypsin digestion. In both studies, compared to undigested samples, samples treated with pepsin alone lost protein bands greater than 150 kDa, and a relatively greater amount of protein bands between 50–100 kDa was identified (Li et al., 2017; Wen et al., 2015). These results indicate that high molecular weight proteins were digested and decomposed into small sizes. In samples treated with pepsin and trypsin together, proteins were decomposed more effectively, and the degraded proteins were confirmed to be clustered in the range between 2–10 kDa (Li et al., 2017; Wen et al., 2015).

Size exclusion chromatography (SEC), a chromatography method developed in 1955, is the most commonly method used to separate polymers such as proteins and peptides according to their molecular size. SEC is utilized for various purposes including adsorption, desalting, and determining molecular weight distribution (Sorci and Belfort, 2014; Wang et al., 2017). As the polymer moves through the column, larger molecules elute faster because they cannot penetrate the pores of the gel, while smaller molecules can penetrate the pores and move freely, increasing the elution time (Boone and Adamec, 2016; Deb et al., 2019). SEC is often used interchangeably with GPC or gel filtration chromatography. GPC is a method of separating molecules by size through elution from a column composed of porous gels such as dextran, agarose, and polyacrylamide. This method can be utilized for extracting specific proteins and analyzing the molecular weight distribution of hydrolysates (Jia et al., 2010; Lee et al., 2022; Ting et al., 2013).

Protein digestibility can be assessed through both *in vivo* and *in vitro* experiments. Kjeldahl assay can be used after feeding experimental animals to determine the crude protein content in feed samples and feces, as well as to calculate feed intake for examining the digestibility of apparent proteins and peptides (AOAC, 2000; Kumar et al., 2019). *In vivo* digestion experiments can provide the most accurate results, but they are time-consuming, costly, and subject to ethical constraints (Boisen and Eggum, 1991; Guerra et al., 2012).

In vitro digestion simulation systems are more efficient compared to *in vivo* digestion experiments and are widely used to evaluate protein digestibility and physiological properties. *In vitro* models have been proposed as an alternative to the financial and ethical challenges of *in vivo* experiments involving humans or animals (Bohn et al., 2018). These systems can be utilized to rapidly screen various food structures and ingredients. In particular, meat (18%) has been identified as the most commonly studied food product using *in vitro* digestion simulation systems after plant foods (45%; Coles et al., 2005; Hur et al., 2011). In a previous study, an *in vitro* digestion simulation system was utilized to identify biochemical indicators of digestibility and nutritional properties of pork muscle protein following different meat processing methods (Bax et al., 2012). Lee et al. (2020) investigated the digestibility and antioxidant properties of beef protein according to aging period and cooking conditions by simulating the digestive conditions of infants. Gallego et al. (2020) also employed an *in vitro* digestion simulation system to evaluate the antioxidant activity of peptides detected after digestion in dry-brined pork hindquarters. However, further testing is needed to confirm the similarity of results obtained from studies using these *in vitro* digestion simulation systems when applied to *in vivo* models.

Utilizing an *in vitro* digestion mimicry system, researchers have identified peptides in fibrillar protein hydrolysates from porcine loin muscle that exhibit partial sequence homology to peptides found through *in vivo* digestion experiments (Escudero et al., 2010b). For example, the peptide AGDDAPR, identified in pork actin, has been found to share partial sequence homology with AGDDAPRAVF and AGFAGDDAPR identified in the duodenum or jejunum of pigs after

consuming beef and trout (Bauchart et al., 2007; Escudero et al., 2010b). However, the digestive enzymes, conditions, and other factors in for each stage of digestion in the *in vitro* simulation system can vary based on age and sex, making it challenging to replicate results from *in vivo* animal experiments. Therefore, comprehensive research is needed to achieve similar outcomes as *in vivo* experiments.

The Applicable Bioavailability Methods for Bioactive Peptides

Antioxidative activities

The antioxidant activity of proteins and peptides is manifested through mechanisms such as free radical scavenging, inactivation of reactive oxygen species, chelation of metal ion, and antioxidant enzyme activity (Elias et al., 2008; Yan et al., 2020).

Free radicals are atoms, molecules, or ions that possess an unpaired electron, making them unstable and highly reactive with other organic compounds (Lobo et al., 2010). These free radicals and other reactive oxygen species derived from oxygen are generated in the body through normal cell metabolism or exposure to external factors such as smoking, ultraviolet light, ozone, and X-rays (Bagchi and Puri, 1998; Carocho and Ferreira, 2013). Reactive oxygen species include the free radicals superoxide ion (O_2^-), hydroxyl radical (HO), hydroperoxyl radical (HO_2), and nitric oxide (NO), as well as other reactive oxygen species such as singlet oxygen (O_2), hydrogen peroxide (H_2O_2), peroxyxynitrite ($ONOO^-$), and hypochlorous acid (HClO; Carocho and Ferreira, 2013; Lobo et al., 2010). These reactive species are neutralized by antioxidant enzymes such as superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase, and various antioxidants (Rock et al., 1996). However, when the balance between reactive species and antioxidants is disrupted, the overabundance of reactive species causes oxidative stress (Rock et al., 1996). Free radicals and other reactive oxygen species exhibit high reactivity with most cellular molecules, including amino acids, sugars, and lipids, causing cellular damage and disruption of homeostasis (Lobo et al., 2010; Young and Woodside, 2001). In addition, excessive oxidative stress can contribute to the development of cancer, liver, kidney, cardiovascular, and neurodegenerative diseases (Carocho and Ferreira, 2013; Soomro, 2019; Tönnies and Trushina, 2017). Chelation is the formation of chelate compounds through the coordination bonding of organic substances with metal ions such as iron and copper (van Lith and Ameer, 2016). Metal ions can trigger redox reactions, leading to oxidative stress and the generation of free radicals that damage biomolecules (van Lith and Ameer, 2016; Yan et al., 2020). Furthermore, an imbalance of metal ions such as iron, copper, zinc, and calcium, along with oxidative stress, can contribute to the development of Alzheimer's disease, a neurodegenerative condition (Wang et al., 2020). Therefore, it is important for antioxidants to effectively inhibit and reduce the interactions of reactive oxygen species with DNA, proteins, lipids, and sugars.

Proteins represent the three-dimensional structure of polypeptides, and most peptides with antioxidant activity are located inside this structure. Therefore, disrupting the three-dimensional structure of proteins through methods such as heat treatment can increase the solvent accessibility of amino acid residues in peptides, thereby enhancing their antioxidant activities (Elias et al., 2008). Furthermore, enzymatic hydrolysis can increase antioxidant activity by breaking peptide bonds to expose amino acid residues. Studies have shown that the antioxidant activity of enzymatically hydrolyzed peptides is higher than that of undigested proteins (Elias et al., 2008; Park and Chin, 2011). In addition, proteins lacking metal ion storage or transport capabilities can chelate metal ions. Proteins with exposed histidine, glutamic acid, and aspartic acid on their surface have been shown to chelate metal ions (Elias et al., 2008). Therefore, there is a need for chelators derived from natural sources with minimal side effects that can bind to these metal ions to form chelate compounds. Peptides derived from pork skeletal

muscle proteins have been found to chelate ferrous ions (Fe^{2+}), and their chelating ability is enhanced through *in vitro* digestion (Zhu et al., 2016). Previous studies have also confirmed that peptides derived from pork proteins and collagen can chelate metal ions (Li et al., 2007; Saiga et al., 2003; Xing et al., 2016).

Experimental methods for measuring the antioxidant activity of proteins and peptides include the 2,2-azinobis-3-ethylbenzothiazoline-6-sulfonic acid (ABTS) radical scavenging assay, 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging assay, iron-chelating assay, and reducing power assay (Acharya, 2017; Bhalodia et al., 2013; Zhong and Shahidi, 2015).

The ABTS assay is considered more responsive and less sensitive to pH than the DPPH assay. Additionally, ABTS assay saves time, money, and sample volume, making it a more efficient option (Moniruzzaman et al., 2012; Shalaby and Shanab, 2013).

The DPPH assay is widely used to evaluate antioxidant activity and is an electron transfer-based assay (Huang et al., 2005; Zhong and Shahidi, 2015). DPPH is a stable nitrogen radical with a dark purple color. Unlike the ABTS assay, the DPPH assay does not require the generation of radicals before conducting the test (Prior et al., 2005).

Iron-chelating assays are used for analyzing antioxidant activity by measuring the capacity of proteins and peptides to chelate Fe^{2+} . This is determined by the level of chromogenicity, as proteins and peptides chelate Fe^{2+} to form chelate compounds, and ferrozine binds to the unchelated Fe^{2+} (Santos et al., 2017). Therefore, low chromogenicity indicates a strong ability of proteins and peptides to bind and chelate Fe^{2+} , indicating high antioxidant activity (Gülçin, 2005).

The reducing power assay is another method that can be utilized to measure the antioxidant activity of proteins and peptides. Since antioxidants also function as reducing agents, reducing power is an important indicator of antioxidant activity (Shahidi and Zhong, 2015). This assay measures reducing power by detecting the reaction of a substance with potassium ferricyanide to form potassium ferrocyanide, which then reacts with ferric chloride to form a ferric-ferrous complex. This reaction results in a yellowish discoloration as the ferric form of potassium ferricyanide is reduced to the ferrous form (Bhalodia et al., 2013; Park et al., 2015).

Experimental methods for measuring antioxidant enzyme activity in proteins and peptides *in vitro* include the SOD assay, CAT assay, and peroxidase (POD) assay (Dasgupta and Klein, 2014; Haida and Hakiman, 2019).

SOD, an antioxidant enzyme, plays an important role in protecting biomolecules from oxidative stress induced by reactive oxygen species (Boguszewska et al., 2010). The SOD assay measures the activity of SOD, which catalyzes the conversion of the free radical superoxide ion (O_2^-) into hydrogen peroxide (H_2O_2) and singlet oxygen (O_2 ; McCord and Fridovich, 1969).

CAT, another antioxidant enzyme, is found in most tissues including the liver and stomach of animals. The CAT assay measures the activity of CAT, which catalyzes the conversion of H_2O_2 to O_2 and H_2O (Liu and Kokare, 2017; Miranda-Bautista et al., 2017). CAT can inhibit cellular damage caused by oxidative stress by reducing the amount of H_2O_2 , a reactive oxygen species produced *in vivo* (Catalán et al., 2018). This is based on the principle that the amount of resorufin, a product that is reduced to O_2 and H_2O by CAT, and the remaining H_2O_2 reacts with horseradish peroxidase (HRP) and a non-fluorescent probe, which is then analyzed by fluorescence or absorbance measurements (Pinto et al., 2011). Similarly, the POD assay determines the activity of POD by measuring the amount of resorufin produced by the reaction of H_2O_2 with HRP and a non-fluorescent probe. Therefore, these antioxidant assays can be used to predict the mechanism of antioxidant activity of bioactive peptides by considering their principles.

Angiotensin converting enzyme inhibitory assay

The antihypertensive activity of proteins and peptides can be measured through the angiotensin converting enzyme (ACE) inhibitory assay. ACE plays a crucial role in the renin-angiotensin system (RAS), which regulates blood pressure (Gurley and

Coffman, 2007; Fig. 1). Gurley and Coffman (2007) have shown that renin in the blood converts angiotensinogen produced by the liver to angiotensin I, which is then converted to angiotensin II by ACE. Within the RAS, ACE converts angiotensin I, an inactive decapeptide, to angiotensin II, an octapeptide with vasoconstrictor activity, and inactivates bradykinin, which exhibits vasodilator activity (Mora et al., 2018; Zhuo et al., 2013). Angiotensin II binds to two G protein coupled-receptors, the AT1 receptor and AT2 receptor, to carry out its biological functions (Wu et al., 2017). The AT1 receptor is associated with a variety of physiological functions, including vasoconstriction, secretion of hormones such as noradrenalin and aldosterone, and sodium reabsorption, while the AT2 receptor promotes vasodilation and sodium excretion (Carey, 2017; Contreras et al., 2003; Kaschina and Unger, 2003; Wu et al., 2017). In this context, antihypertensive functional peptides can reduce angiotensin II production by inhibiting ACE and lower blood pressure by blocking the AT1 receptor (Contreras et al., 2003; Ferrario et al., 2005). In addition, antihypertensive functional peptides play a role in balancing the vasoconstrictor and dilator effects of angiotensin I and bradykinin (Mora et al., 2018).

On the other hand, ACE2, which has been identified as a homologue of ACE, is known to play a physiological role in the regulation of homeostasis (Turner, 2015). In addition, ACE2 cleaves the amino acids at the C-terminus of angiotensin II to form angiotensins 1–7, which bind to the Mas receptor and exert anti-inflammatory, vasodilatory, and antifibrotic effects (Barroso et al., 2015; Shenoy et al., 2015; Simões e Silva et al., 2013). Similarly, ACE2 can hydrolyze angiotensin I to produce angiotensin 1–9, which can be converted to angiotensin 1–7 by ACE action (Donoghue et al., 2000). Previous studies have reported that angiotensin 1–9 can exhibit vasodilatory functions, reducing blood pressure and preventing cardiomyocyte hypertrophy (Gonzalez et al., 2018; Sotomayor-Flores et al., 2020). Therefore, the antihypertensive activity of the peptide may be mainly determined by ACE inhibition.




Fig. 1. The renin-angiotensin system. ACE, angiotensin converting enzyme.

Bioactive Peptides in Pork

It has been confirmed that bioactive peptides exhibit little bioactivity in their normal protein-bound state, and their activity is triggered by protein degradation through processes such as ripening, fermentation, enzymatic hydrolysis, and digestion (Arihara and Ohata, 2008; Xing et al., 2019). Previous studies have shown that plant-derived bioactive peptides are extracted using digestive enzymes such as trypsin, chymotrypsin, and pepsin, or plant-derived proteolytic enzymes papain, bromelain, and ficin (Ryan et al., 2011; Singh et al., 2019). Additionally, alkaline proteases from microbial fermentation processes have been identified to be used to produce highly nutritious protein hydrolysates (Sharma et al., 2019; Sumantha et al., 2006). The use of commercialized proteolytic enzymes, including Protamex[®] and Flavourzyme[®], for the production of ACE inhibitory active peptides has been previously documented (Mirdhayati et al., 2016). Furthermore, it has been reviewed that bioactivities, such as antioxidant and ACE inhibitory activities of protein hydrolysates formed by using alcalase with other proteolytic enzymes are increased (Tacias-Pascacio et al., 2020). In addition, large amounts of bioactive peptides have been produced from pork after *in vitro* digestion, confirming that pork can be a major source of bioactive peptides (Escudero et al., 2010b). The process of bioactive peptide formation from proteins in meat is shown in Fig. 2.




Fig. 2. The analytical process for assessing the digestibility and bioactivities of myofibrillar proteins and peptide extracts from Jeju black pigs and three-way crossbred pigs (Landrace×Yorkshire×Duroc, LYD). ICR, Institute of Cancer Research; BCA, bichinchonic acid; HPLC, high-performance liquid chromatography; SDS-PAGE, sodium dodecyl sulfate-polyacrylamide gel electrophoresis; GPC, gel permeation chromatography; ACE, angiotensin converting enzyme; SOD, superoxide dismutase; CAT, catalase; POD, peroxidase.

Table 1 displays the peptides with antioxidant activity derived from porcine proteins. All of these peptides have a small molecular weight, mostly less than 1 kDa. In addition, bioactive peptides with high DPPH radical scavenging and metal ion chelating activities were extracted from protein hydrolysate obtained from pork source fiber protein (Saiga et al., 2003). Carnosine and anserine, representative peptides with antioxidant activity, were also obtained from porcine loin muscle (Simonetti et al., 2019). Furthermore, bioactive peptides predicted to be generated after hydrolysis from pork myofibrillar proteins were identified through *in silico* analysis as potentially exhibiting a variety of bioactivities, including antioxidant, antihypertensive, antithrombotic, and dipeptidyl peptidase-IV (DPP-IV) inhibition (Keska and Stadnik, 2017). DPP-IV is an enzyme that degrades incretin, a blood sugar-regulating hormone released when food is consumed. Inhibition of DPP-IV increases incretin content, stimulating the release of insulin and inhibiting the release of glucagon, which regulates blood sugar (Drucker, 2007). Peptides extracted from dry-cured pork ham with a molecular weight of less than 1 kDa exhibit the highest antioxidant activity (Xing et al., 2018). Meanwhile, meat-derived bioactive peptides are considered to have higher antioxidant activity as they contain more hydrophobic amino acids (leucine, isoleucine, and valine) and aromatic amino acids (tryptophan, tyrosine, and phenylalanine; Peighambardoust et al., 2021).

Table 1. Antioxidant peptides in pork

Protein source	Peptide sequence	Treatment	MW (Da) ¹⁾	Reference
Porcine myofibrillar protein (actin)	DSGVT	Papain	650.3	Saiga et al. (2003)
Porcine myofibrillar protein (unknown)	IEAEGE	Papain	646.4	Saiga et al. (2003)
Porcine myofibrillar protein (tropomyosin)	DAQEKLE	Papain	832.5	Saiga et al. (2003)
Porcine myofibrillar protein (tropomyosin)	EELDNALN	Papain	916.9	Saiga et al. (2003)
Porcine myofibrillar protein (myosin heavy chain)	VPSIDDQEELM	Papain	1,275.0	Saiga et al. (2003)
Porcine muscle (actomyosin)	DLYA	Papain	480.5	Arihara (2006)
Porcine muscle (actomyosin)	SLYA	Papain	452.5	Arihara (2006)
Porcine muscle (actomyosin)	VW	Papain	303.4	Arihara (2006)
Porcine ham skeletal muscle proteins	GKFNV, HA, LPGGGT, LPGGGHGDL	Dry-cured; Pepsin+trypsin	-	Zhu et al. (2016)
Porcine <i>biceps femoris</i> muscle proteins	GLAGA, SAGNPN	Dry-cured	-	Escudero et al. (2013)
Porcine fresh ground ham	QYP	Fermentation	-	Ohata et al. (2016)
Porcine <i>biceps femoris</i> muscle proteins	DLEE	Dry-cured	504.2	Xing et al. (2016)
Porcine ham muscle proteins	MDPKYR, TKYRVP	Dry-cured	-	Gallego et al. (2019)
Porcine <i>longissimus dorsi</i> muscle	VW, LW	<i>In vitro</i> gastro-intestinal digestion	<3,000	Martini et al. (2019)
Porcine ham muscle proteins (actin)	EAGPSIVHR, ALPHAIMR, AGFAGDDAPR, VAPEEHPTL, DEAGPSIVH, AGPSIVHRK	Dry-cured	908.1–992.1	Wang et al. (2021)
Porcine ham muscle proteins (tropomyosin)	MDAIKKK, DPIIQDR	Dry-cured	833.0–856.0	Wang et al. (2021)

¹⁾ Molecular weight measured in Daltons (Da).

Among the bioactive peptides, the most extensively studied are angiotensin-converting enzyme inhibitory peptides (Arihara and Ohata, 2008).

According to the World Health Organization, approximately 1.13 billion people worldwide have high blood pressure (WHO, 2013; WHO, 2021). High blood pressure can weaken the heart, damage artery walls, alter blood flow, and lead to complications such as stroke, heart disease, kidney failure, vision loss, and hardening of the arteries (Williams et al., 2018). Due to the severe side effects of various synthetic drugs used to treat hypertension, there has been extensive research on bioactive peptides derived from food proteins that can effectively treat hypertension without causing adverse reactions (Toldrá et al., 2018). Table 2 displays the antihypertensive functional peptides derived from porcine proteins, with most originating from fibrillar proteins such as myosin, actin, and troponin. Previous studies have shown that peptides with a molecular weight of less than 10 kDa have superior antioxidant and antihypertensive properties compared to larger peptides with relatively larger molecular weights. Some peptides obtained from pork proteins through *in vitro* digestion have shown ACE inhibitory activity (Escudero et al., 2010a; Escudero et al., 2012). For example, peptides (MYPGIA and VIPEL) derived from pork actin and GAPDH, and peptides (KRQYDI and VKAGF) isolated from pork myosin heavy chain and actin exhibit ACE inhibitory activity (Escudero et al., 2010a; Muguruma et al., 2009). Peptides KAPVA and PTPVP from titin, and

Table 2. Angiotensin I-converting enzyme (ACE)-inhibitory peptides in pork

Protein source	Peptide sequence	Treatment	MW (Da) ¹⁾	Reference
Porcine muscle (myosin)	MNPPK	Thermolysin	585.7 ²⁾	Nakashima et al. (2002)
Porcine muscle (myosin)	ITTNP	Thermolysin	-	Nakashima et al. (2002)
Porcine muscle (myosin light chain)	VKKVLGNP	Pepsin	854.0	Katayama et al. (2007)
Porcine muscle (troponin)	KRQKYDI	Pepsin	950.1 ²⁾	Muguruma et al. (2009)
Porcine muscle (myosin heavy chain)	KRVITY	Pepsin	805.97	Muguruma et al. (2009)
Porcine muscle (actin)	VKAGF	Pepsin	520.62	Muguruma et al. (2009)
Porcine muscle (nebulin)	RPR	Pepsin+pancreatin	-	Escudero et al. (2012)
Porcine muscle (titin)	KAPVA	Pepsin+pancreatin	-	Escudero et al. (2012)
Porcine muscle (titin)	PTPVP	Pepsin+pancreatin	-	Escudero et al. (2012)
Porcine muscle (actin)	MYPGIA	Pepsin+pancreatin	-	Escudero et al. (2010b)
Porcine muscle (GAPDH)	VIPEL	Pepsin+pancreatin	-	Escudero et al. (2010b)
Porcine <i>longissimus dorsi</i> muscle (actin)	VFPS, LKYPI, AVF, MYPGIA	<i>In vitro</i> gastro-intestinal digestion	<3,000	Martini et al. (2019)
Porcine <i>longissimus dorsi</i> muscle	VW, IW, VF, WL, LW, VIP, LGI, LPF, IVP, IL, LLF, WM, FIV, LR, ILP, VLP, PL, LF, IAIP, IR, IF, GLx, AV, AI, DL, NIIPA	<i>In vitro</i> gastro-intestinal digestion	<3,000	Martini et al. (2019)
Porcine ham muscle proteins	GGVPGG, TKYRVP, HCNKKYRSEM	Dry-cured	-	Gallego et al. (2019)
Porcine ham muscle proteins	EL, EV, RL, EEL, ESV	Dry-cured	-	Hao et al. (2020)
Porcine ham muscle proteins	GA, VF	Dry-cured	-	Heres et al. (2021b), Heres et al. (2022)

¹⁾ Molecular weight measured in Daltons (Da).

²⁾ The peptide molecular weight was derived from the PubChem.

peptide RPR from neblin in pork enzymatic hydrolysate, show strong ACE inhibitory activity (Escudero et al., 2012). Furthermore, differences in the amino acid composition of bioactive peptides may affect ACE-inhibitory activity. For example, differences in the composition of amino acids that make up peptides, such as acidic amino acids (aspartic acid and glutamic acid), and the presence of positively charged amino acids in the carboxyl group can affect the increase in ACE-inhibitory activity (Daskaya-Dikmen et al., 2017; Peighamardoust et al., 2021).

Conclusion

In this study, we presented a protein digestion analysis method and a peptide bioactivity analysis method that can be utilized for peptide acquisition. The digestive enzymes present in the intestinal tract include pepsin, trypsin, chymotrypsin, and procarboxypeptidase, with cholecystokinin and secretin playing auxiliary roles in protein digestion. Proteins are hydrolyzed in the body to generate peptides. Methods such as BCA assay, SDS-PAGE, and chromatography have been used to analyze protein digestibility and molecular weight distribution, which are applicable to peptide acquisition. In recent years, *in vitro* digestion simulation systems have been utilized to evaluate protein digestibility and changes in activity. In addition, the ACE inhibitory and antioxidant properties of bioactive peptides derived from pork suggest potential industrial applications. In particular, papain has been primarily used as a hydrolyzing agent for antioxidant peptides in pork. Actomyosin and tropomyosin are found in myofibrillar proteins, and they have molecular weights below 1 kDa. The antihypertensive activity is often attributed to the use of pepsin as a hydrolyzing agent in pork, with most peptides identified having a molecular weight of lower than 1 kDa. Therefore, this study can serve as a basis for the effective utilization in the development of pork-derived bioactive peptides and exploration of their bioactivity in the future. Furthermore, the advancement of pork-derived bioactive peptides may aid in promoting domestic pork consumption.

Conflicts of Interest

The authors declare no potential conflicts of interest.

Acknowledgements

This work was conducted with the support of Chung-Ang University. This work was carried out with the support of “Cooperative Research Program for Agriculture Science and Technology Development (Project No. PJ016201)” Rural Development Administration, Korea.

Author Contributions

Conceptualization: Jeong JW, Hur SJ. Validation: Lee SY, Lee DY, Lee J, Mariano EJ, Moon SS. Investigation: Jeong JW, Lee DY, Kim JH, Yun SH, Mariano EJ. Writing - original draft: Jeong JW, Hur SJ. Writing - review & editing: Jeong JW, Lee SY, Lee DY, Kim JH, Yun SH, Lee J, Mariano EJ, Moon SS, Hur SJ.

Ethics Approval

This article does not require IRB/IACUC approval because there are no human and animal participants.

References

- Acharya K. 2017. Simplified methods for microtiter based analysis of *in vitro* antioxidant activity. *Asian J Pharm* 11:S327-S335.
- Adibi SA, Mercer DW. 1973. Protein digestion in human intestine as reflected in luminal, mucosal, and plasma amino acid concentrations after meals. *J Clin Invest* 52:1586-1594.
- Aluko RE. 2015. Antihypertensive peptides from food proteins. *Annu Rev Food Sci Technol* 6:235-262.
- AOAC. 2000. Official methods of analysis. Association of Official Analytical Chemists, Arlington, VA, USA.
- Arihara K. 2006. Functional properties of bioactive peptides derived from meat proteins. In *Advanced technologies for meat processing*. Nollet NML, Toldrá F (ed). CRC Press, Boca Raton, FL, USA. pp 245-274.
- Arihara K, Ohata M. 2008. Bioactive compounds in meat. In *Meat biotechnology*. Toldrá F (ed). Springer, New York, NY, USA. pp 231-249.
- Bagchi K, Puri S. 1998. Free radicals and antioxidants in health and disease: A review. *East Mediterr Health J* 4:350-360.
- Barroso LC, Silveira KD, Teixeira MM, Simões Silva AC. 2015. Mas and inflammation. In *The protective arm of the renin angiotensin system (RAS): Functional aspects and therapeutic implications*. Unger T, Steckelings UM, dos Santos RAS (ed). Academic Press, Cambridge, MA, USA. pp 213-217.
- Bauchart C, Morzel M, Chambon C, Mirand PP, Reynès C, Buffière C, Rémond D. 2007. Peptides reproducibly released by *in vivo* digestion of beef meat and trout flesh in pigs. *Br J Nutr* 98:1187-1195.
- Bax ML, Aubry L, Ferreira C, Daudin JD, Gatellier P, Rémond D, Santé-Lhoutellier V. 2012. Cooking temperature is a key determinant of *in vitro* meat protein digestion rate: Investigation of underlying mechanisms. *J Agric Food Chem* 60:2569-2576.
- Bhalodia NR, Nariya PB, Acharya RN, Shukla VJ. 2013. *In vitro* antioxidant activity of hydro alcoholic extract from the fruit pulp of *Cassia fistula* Linn. *Ayu* 34:209-214.
- Bhutia YD, Ganapathy V. 2018. Protein digestion and absorption. In *Physiology of the gastrointestinal tract*. 6th ed. Said HM (ed). Academic Press, Cambridge, MA, USA. pp 1063-1086.
- Boguszewska D, Grudkowska M, Zagdańska B. 2010. Drought-responsive antioxidant enzymes in potato (*Solanum tuberosum* L.). *Potato Res* 53:373-382.
- Bohn T, Carriere F, Day L, Deglaire A, Egger L, Freitas D, Golding M, Le Feunteun S, Macierzanka A, Menard O, Miralles B, Moscovici A, Portmann R, Recio I, Rémond D, Santé-Lhoutellier V, Wooster TJ, Lesmes U, Mackie AR, Dupont D. 2018. Correlation between *in vitro* and *in vivo* data on food digestion. What can we predict with static *in vitro* digestion models? *Crit Rev Food Sci Nutr* 58:2239-2261.
- Boisen S, Eggum BO. 1991. Critical evaluation of *in vitro* methods for estimating digestibility in simple-stomach animals. *Nutr Res Rev* 4:141-162.
- Boone C, Adamec J. 2016. Top-down proteomics. In *Proteomic profiling and analytical chemistry: The crossroads*. 2nd ed. Ciborowski P, Silberring J (ed). Elsevier, Amsterdam, The Netherlands. pp 175-191.
- Boron WF, Boulpaep EL. 2016. Nutrient digestion and absorption. In *Medical physiology*. Boron WF, Boulpaep EL (ed). Elsevier, Amsterdam, The Netherlands. pp 914-943.
- Carey RM. 2017. AT2 receptors: Potential therapeutic targets for hypertension. *Am J Hypertens* 30:339-347.
- Carocho M, Ferreira ICFR. 2013. A review on antioxidants, prooxidants and related controversy: Natural and synthetic

- compounds, screening and analysis methodologies and future perspectives. *Food Chem Toxicol* 51:15-25.
- Catalán V, Frühbeck G, Gómez-Ambrosi J. 2018. Inflammatory and oxidative stress markers in skeletal muscle of obese subjects. In *Obesity: Oxidative stress and dietary antioxidants*. Moral AM, Aguilera García CM (ed). Academic Press, Cambridge, MA, USA. pp 163-189.
- Chalamaiah M, Ulug SK, Hong H, Wu J. 2019. Regulatory requirements of bioactive peptides (protein hydrolysates) from food proteins. *J Funct Foods* 58:123-129.
- Coles LT, Moughan PJ, Darragh AJ. 2005. *In vitro* digestion and fermentation methods, including gas production techniques, as applied to nutritive evaluation of foods in the hindgut of humans and other simple-stomached animals. *Anim Feed Sci Technol* 123–124:421-444.
- Contreras F, de la Parte MA, Antonia M, Cabrera J, Ospino N, Israili ZH, Velasco M. 2003. Role of angiotensin II AT1 receptor blockers in the treatment of arterial hypertension. *Am J Ther* 10:401-408.
- Dasgupta A, Klein K. 2014. Methods for measuring oxidative stress in the laboratory. In *Antioxidants in food, vitamins and supplements: Prevention and treatment of disease*. Dasgupta A, Klein K (ed). Elsevier, Amsterdam, The Netherlands. pp 19-40.
- Daskaya-Dikmen C, Yucetepe A, Karbancioglu-Guler F, Daskaya H, Ozcelik B. 2017. Angiotensin-I-converting enzyme (ACE)-inhibitory peptides from plants. *Nutrients* 9:316.
- Deb PK, Kokaz SF, Abed SN, Paradkar A, Tekade RK. 2019. Pharmaceutical and biomedical applications of polymers. In *Basic fundamentals of drug delivery: A volume in advances in pharmaceutical product development and research*. Tekade RK (ed). Academic Press, Cambridge, MA, USA. pp 203-267.
- Di Bernardini R, Harnedy P, Bolton D, Kerry J, O'Neill E, Mullen AM, Hayes M. 2011. Antioxidant and antimicrobial peptidic hydrolysates from muscle protein sources and by-products. *Food Chem* 124:1296-1307.
- DiGregorio N, Sharma S. 2019. Physiology, secretin. Available from: <https://europepmc.org/article/nbk/nbk537116>. Accessed at Sep 13, 2022.
- Donoghue M, Hsieh F, Baronas E, Godbout K, Gosselin M, Stagliano N, Donovan M, Woolf B, Robison K, Jeyaseelan R, Breitbart RE, Acton S. 2000. A novel angiotensin-converting enzyme–related carboxypeptidase (ACE2) converts angiotensin I to angiotensin 1-9. *Circ Res* 87:e1-e9.
- Drucker DJ. 2007. The role of gut hormones in glucose homeostasis. *J Clin Invest* 117:24-32.
- Elias RJ, Kellerby SS, Decker EA. 2008. Antioxidant activity of proteins and peptides. *Crit Rev Food Sci Nutr* 48:430-441.
- Escudero E, Mora L, Fraser PD, Aristoy MC, Toldrá F. 2013. Identification of novel antioxidant peptides generated in Spanish dry-cured ham. *Food Chem* 138:1282-1288.
- Escudero E, Sentandreu MA, Arihara K, Toldrá F. 2010a. Angiotensin I-converting enzyme inhibitory peptides generated from *in vitro* gastrointestinal digestion of pork meat. *J Agric Food Chem* 58:2895-2901.
- Escudero E, Sentandreu MÁ, Toldrá F. 2010b. Characterization of peptides released by *in vitro* digestion of pork meat. *J Agric Food Chem* 58:5160-5165.
- Escudero E, Toldrá F, Sentandreu MA, Nishimura H, Arihara K. 2012. Antihypertensive activity of peptides identified in the *in vitro* gastrointestinal digest of pork meat. *Meat Sci* 91:382-384.
- Farrell RE Jr. 2009. Resilient ribonucleases. In *RNA methodologies: A laboratory guide for isolation and characterization*. 4th ed. Farrell RE Jr (ed). Academic Press, Cambridge, MA, USA. pp 155-172.
- Ferrario CM, Jessup J, Chappell MC, Averill DB, Brosnihan KB, Tallant EA, Diz DI, Gallagher PE. 2005. Effect of

- angiotensin-converting enzyme inhibition and angiotensin II receptor blockers on cardiac angiotensin-converting enzyme. *Circulation* 111:2605-2610.
- Fischer T, Elenko E, McCaffery JM, DeVries L, Farquhar MG. 1999. Clathrin-coated vesicles bearing GAIP possess GTPase-activating protein activity *in vitro*. *Proc Natl Acad Sci USA* 96:6722-6727.
- Gallego M, Mauri L, Aristoy MC, Toldrá F, Mora L. 2020. Antioxidant peptides profile in dry-cured ham as affected by gastrointestinal digestion. *J Funct Foods* 69:103956.
- Gallego M, Mora L, Toldrá F. 2019. Potential cardioprotective peptides generated in Spanish dry-cured ham. *J Food Bioact* 6:110-117.
- Gonzalez L, Novoa U, Moya J, Gabrielli L, Jalil JE, García L, Chiong M, Lavandero S, Ocaranza MP. 2018. Angiotensin-(1-9) reduces cardiovascular and renal inflammation in experimental renin-independent hypertension. *Biochem Pharmacol* 156:357-370.
- Guerra A, Etienne-Mesmin L, Livrelli V, Denis S, Blanquet-Diot S, Alric M. 2012. Relevance and challenges in modeling human gastric and small intestinal digestion. *Trends Biotechnol* 30:591-600.
- Gülçin İ. 2005. The antioxidant and radical scavenging activities of black pepper (*Piper nigrum*) seeds. *Int J Food Sci Nutr* 56:491-499.
- Gupta A. 2018. Digestion and absorption of proteins. In *Comprehensive biochemistry for dentistry: Textbook for dental students*. Gupta A (ed). Springer, Singapore. pp 367-375.
- Gurley SB, Coffman TM. 2007. The renin-angiotensin system and diabetic nephropathy. *Semin Nephrol* 27:144-152.
- Haida Z, Hakimian M. 2019. A comprehensive review on the determination of enzymatic assay and nonenzymatic antioxidant activities. *Food Sci Nutr* 7:1555-1563.
- Hao L, Gao X, Zhou T, Cao J, Sun Y, Dang Y, Pan D. 2020. Angiotensin I-converting enzyme (ACE) inhibitory and antioxidant activity of umami peptides after *in vitro* gastrointestinal digestion. *J Agric Food Chem* 68:8232-8241.
- He F. 2011. BCA (bicinchoninic acid) protein assay. *BI Protoc* 1:e44.
- Heres A, Gallego M, Mora L, Toldrá F. 2022. Identification and quantitation of bioactive and taste-related dipeptides in low-salt dry-cured ham. *Int J Mol Sci* 23:2507.
- Heres A, Mora L, Toldrá F. 2021a. Inhibition of 3-hydroxy-3-methyl-glutaryl-coenzyme A reductase enzyme by dipeptides identified in dry-cured ham. *Food Prod Process Nutr* 3:1-14.
- Heres A, Saldaña C, Toldrá F, Mora L. 2021b. Identification of dipeptides by MALDI-ToF mass spectrometry in long-processing Spanish dry-cured ham. *Food Chem Mol Sci* 3:100048.
- Hinsberger A, Sandhu BK. 2004. Digestion and absorption. *Curr Paediatr* 14:605-611.
- Hooton D, Lentle R, Monro J, Wickham M, Simpson R. 2015. The secretion and action of brush border enzymes in the mammalian small intestine. In *Reviews of physiology, biochemistry and pharmacology*. Nilius B, Gudermann T, Jahn R, Lill R, Petersen OH, de Tombe PP (ed). Springer, Cham, Switzerland. pp 59-118.
- Huang D, Ou B, Prior RL. 2005. The chemistry behind antioxidant capacity assays. *J Agric Food Chem* 53:1841-1856.
- Hur SJ, Lim BO, Decker EA, McClements DJ. 2011. *In vitro* human digestion models for food applications. *Food Chem* 125:1-12.
- Jia J, Zhou Y, Lu J, Chen A, Li Y, Zheng G. 2010. Enzymatic hydrolysis of Alaska pollack (*Theragra chalcogramma*) skin and antioxidant activity of the resulting hydrolysate. *J Sci Food Agric* 90:635-640.
- Kaschina E, Unger T. 2003. Angiotensin AT1/AT2 receptors: Regulation, signalling and function. *Blood Press* 12:70-88.

- Katayama K, Mori T, Kawahara S, Miake K, Kodama Y, Sugiyama M, Kawamura Y, Nakayama T, Maruyama M, Muguruma M. 2007. Angiotensin-I converting enzyme inhibitory peptide derived from porcine skeletal muscle myosin and its antihypertensive activity in spontaneously hypertensive rats. *J Food Sci* 72:S702-S706.
- Keska P, Stadnik J. 2017. Antimicrobial peptides of meat origin: An *in silico* and *in vitro* analysis. *Protein Pept Lett* 24:165-173.
- Kim JH, Lee DY, Lee SY, Mariano E Jr, Jeong JW, Yun SH, Lee J, Park J, Choi Y, Han D, Kim JS, Jo C, Hur SJ. 2024. Study on the digestion-induced changes in the characteristics and bioactivity of Korean native and overseas cattle-derived peptides. *Food Sci Anim Resour* 44:551-569.
- Kumar A, Badgajar PC, Mishra V, Sehrawat R, Babar OA, Upadhyay A. 2019. Effect of microfluidization on cholesterol, thermal properties and *in vitro* and *in vivo* protein digestibility of milk. *LWT-Food Sci Technol* 116:108523.
- Lafarga T, Hayes M. 2014. Bioactive peptides from meat muscle and by-products: Generation, functionality and application as functional ingredients. *Meat Sci* 98:227-239.
- Lee JA, Kim MJ, Shin MR, Roh SS, Lee JB, Seo Y, Choi HG, Park HJ. 2022. Determination of the protein quality of low-molecular weight water-soluble chicken breast powder by a protein digestibility corrected amino acid score (PDCAAS) analysis. *J Korean Soc Food Sci Nutr* 51:439-447.
- Lee S, Jo K, Lee HJ, Jo C, Yong HI, Choi YS, Jung S. 2020. Increased protein digestibility of beef with aging in an infant *in vitro* digestion model. *Meat Sci* 169:108210.
- Li B, Chen F, Wang X, Ji B, Wu Y. 2007. Isolation and identification of antioxidative peptides from porcine collagen hydrolysate by consecutive chromatography and electrospray ionization–mass spectrometry. *Food Chem* 102:1135-1143.
- Li L, Liu Y, Zou X, He J, Xu X, Zhou G, Li C. 2017. *In vitro* protein digestibility of pork products is affected by the method of processing. *Food Res Int* 92:88-94.
- Liddle RA. 1997. Cholecystokinin cells. *Annu Rev Physiol* 59:221-242.
- Liu X, Kokare C. 2017. Microbial enzymes of use in industry. In *Biotechnology of microbial enzymes: Production, biocatalysis and industrial applications*. Brahmachari G (ed). Academic Press, Cambridge, MA, USA. pp 267-298.
- Lobo V, Patil A, Phatak A, Chandra N. 2010. Free radicals, antioxidants and functional foods: Impact on human health. *Pharmacogn Rev* 4:118-126.
- Martini S, Conte A, Tagliacruzchi D. 2019. Comparative peptidomic profile and bioactivities of cooked beef, pork, chicken and turkey meat after *in vitro* gastro-intestinal digestion. *J Proteomics* 208:103500.
- McCord JM, Fridovich I. 1969. Superoxide dismutase: An enzymic function for erythrocyte hemocuprein (hemocuprein). *J Biol Chem* 244:6049-6055.
- Miranda-Bautista J, Bañares R, Vaquero J. 2017. The gastrointestinal system: Anatomy and sources of oxidative stress. In *Gastrointestinal tissue: Oxidative stress and dietary antioxidants*. Gracia-Sancho J, Salvadó J (ed). Academic Press, Cambridge, MA, USA. pp 3-20.
- Mirdhayati I, Hermanianto J, Wijaya CH, Sajuthi D, Arihara K. 2016. Angiotensin converting enzyme (ACE) inhibitory and antihypertensive activities of protein hydrolysate from meat of Kacang goat (*Capra aegagrus hircus*). *J Sci Food Agric* 96:3536-3542.
- Moniruzzaman M, Khalil MI, Sulaiman SA, Gan SH. 2012. Advances in the analytical methods for determining the antioxidant properties of honey: A review. *Afr J Tradit Complement Altern Med* 9:36-42.
- Mora L, Gallego M, Toldrá F. 2018. ACEI-inhibitory peptides naturally generated in meat and meat products and their health

- relevance. *Nutrients* 10:1259.
- Muguruma M, Ahmed AM, Katayama K, Kawahara S, Maruyama M, Nakamura T. 2009. Identification of pro-drug type ACE inhibitory peptide sourced from porcine myosin B: Evaluation of its antihypertensive effects *in vivo*. *Food Chem* 114:516-522.
- Nakashima Y, Arihara K, Sasaki A, Mio H, Ishikawa S, Itoh M. 2002. Antihypertensive activities of peptides derived from porcine skeletal muscle myosin in spontaneously hypertensive rats. *J Food Sci* 67:434-437.
- Ohata M, Uchida S, Zhou L, Arihara K. 2016. Antioxidant activity of fermented meat sauce and isolation of an associated antioxidant peptide. *Food Chem* 194:1034-1039.
- Park JH, Bae NY, Park SH, Kim MJ, Kim KBWR, Choi JS, Ahn DH. 2015. Antioxidant effect of *Sargassum coreanum* root and stem extracts. *KSBB J* 30:155-160.
- Park SY, Chin KB. 2011. Antioxidant activities of pepsin hydrolysates of water- and salt-soluble protein extracted from pork hams. *Int J Food Sci Technol* 46:229-235.
- Peighambaroust SH, Karami Z, Pateiro M, Lorenzo JM. 2021. A review on health-promoting, biological, and functional aspects of bioactive peptides in food applications. *Biomolecules* 11:631.
- Pinto PCAG, Costa ADF, Lima JLFC, Saraiva MLMFS. 2011. Automated evaluation of the effect of ionic liquids on catalase activity. *Chemosphere* 82:1620-1628.
- Prior RL, Wu X, Schaich K. 2005. Standardized methods for the determination of antioxidant capacity and phenolics in foods and dietary supplements. *J Agric Food Chem* 53:4290-4302.
- Rajput YS, Sharma R. 2011. SDS-PAGE: Principle and applications. In *Chemical analysis of value added dairy products and their quality assurance*. Sharma R (ed). National Dairy Research Institute, Karnal, India. p 81.
- Rezvankhah A, Yarmand MS, Ghanbarzadeh B, Mirzaee H. 2021. Generation of bioactive peptides from lentil protein: Degree of hydrolysis, antioxidant activity, phenol content, ACE-inhibitory activity, molecular weight, sensory, and functional properties. *J Food Meas Charact* 15:5021-5035.
- Righetti PG. 2005. Electrophoresis | Polyacrylamide gels. In *Encyclopedia of analytical science*. 2nd ed. Worsfold P, Townshend A, Poole C (ed). Elsevier, Amsterdam, The Netherlands. pp 396-407.
- Rock CL, Jacob RA, Bowen PE. 1996. Update on the biological characteristics of the antioxidant micronutrients: Vitamin C, vitamin E, and the carotenoids. *J Am Diet Assoc* 96:693-702.
- Roy VK, Kumar NS, Gurusubramanian G. 2012. Proteins: Structure, properties and their separation by SDS-polyacrylamide gel electrophoresis. *Sci Vis* 12:170-181.
- Rubak YT, Nuraida L, Iswantini D, Prangdimurti E. 2022. Angiotensin-I-converting enzyme inhibitory peptides in goat milk fermented by lactic acid bacteria isolated from fermented food and breast milk. *Food Sci Anim Resour* 42:46-60.
- Ryan JT, Ross RP, Bolton D, Fitzgerald GF, Stanton C. 2011. Bioactive peptides from muscle sources: Meat and fish. *Nutrients* 3:765-791.
- Saiga A, Tanabe S, Nishimura T. 2003. Antioxidant activity of peptides obtained from porcine myofibrillar proteins by protease treatment. *J Agric Food Chem* 51:3661-3667.
- Sánchez A, Vázquez A. 2017. Bioactive peptides: A review. *Food Qual Saf* 1:29-46.
- Santos JS, Brizola VRA, Granato D. 2017. High-throughput assay comparison and standardization for metal chelating capacity screening: A proposal and application. *Food Chem* 214:515-522.
- Shahidi F, Zhong Y. 2015. Measurement of antioxidant activity. *J Funct Foods* 18:757-781.

- Shalaby EA, Shanab SMM. 2013. Antioxidant compounds, assays of determination and mode of action. *Afr J Pharm Pharmacol* 7:528-539.
- Sharma M, Gat Y, Arya S, Kumar V, Panghal A, Kumar A. 2019. A review on microbial alkaline protease: An essential tool for various industrial approaches. *Ind Biotechnol* 15:69-78.
- Shenoy V, Ferreira AJ, Katovich M, Raizada MK. 2015. Angiotensin-converting enzyme 2/angiotensin-(1-7)/mas receptor axis: Emerging pharmacological target for pulmonary diseases. In *The protective arm of the renin angiotensin system (RAS): Functional aspects and therapeutic implications*. Unger T, Steckelings UM, dos Santos RAS (ed). Academic Press, Cambridge, MA, USA. pp 269-274.
- Simões e Silva AC, Silveira KD, Ferreira AJ, Teixeira MM. 2013. ACE2, angiotensin-(1-7) and Mas receptor axis in inflammation and fibrosis. *Br J Pharmacol* 169:477-492.
- Simonetti A, Perna A, Gambacorta E. 2019. Comparison of antioxidant compounds in pig meat from Italian autochthonous pig Suino Nero Lucano and a modern crossbred pig before and after cooking. *Food Chem* 292:108-112.
- Singh PK, Shrivastava N, Ojha BK. 2019. Enzymes in the meat industry. In *Enzymes in food biotechnology: Production, applications, and future prospects*. Kuddus M (ed). Academic Press, Cambridge, MA, USA. pp 111-128.
- Soomro S. 2019. Oxidative stress and inflammation. *Open J Immunol* 9:1-20.
- Sorci M, Belfort G. 2014. Insulin oligomers: Detection, characterization and quantification using different analytical methods. In *Bio-nanoimaging: Protein misfolding and aggregation*. Uversky VN, Lyubchenko YL (ed). Academic Press, Cambridge, MA, USA. pp 233-245.
- Sotomayor-Flores C, Rivera-Mejías P, Vásquez-Trincado C, López-Crisosto C, Morales PE, Pennanen C, Polakovicova I, Aliaga-Tobar V, García L, Carlos Roa J, Rothermel BA, Maracaja-Coutinho V, Ho-Xuan H, Meister G, Chiong M, Paz Ocaranza M, Corvalán AH, Parra V, Lavandero S. 2020. Angiotensin-(1-9) prevents cardiomyocyte hypertrophy by controlling mitochondrial dynamics via miR-129-3p/PKIA pathway. *Cell Death Differ* 27:2586-2604.
- Sumantha A, Larroche C, Pandey A. 2006. Microbiology and industrial biotechnology of food-grade proteases: A perspective. *Food Technol Biotechnol* 44:211-220.
- Tacias-Pascacio VG, Morellon-Sterling R, Siar EH, Tavano O, Berenguer-Murcia Á, Fernandez-Lafuente R. 2020. Use of alcalase in the production of bioactive peptides: A review. *Int J Biol Macromol* 165:2143-2196.
- Ting BCP, Pouliot Y, Gauthier SF, Mine Y. 2013. Fractionation of egg proteins and peptides for nutraceutical applications. In *Separation, extraction and concentration processes in the food, beverage and nutraceutical industries*. Rizvi SSH (ed). Woodhead, Sawston, UK. pp 595-618.
- Toldrá F, Reig M, Aristoy MC, Mora L. 2018. Generation of bioactive peptides during food processing. *Food Chem* 267:395-404.
- Tönnies E, Trushina E. 2017. Oxidative stress, synaptic dysfunction, and Alzheimer's disease. *J Alzheimer's Dis* 57:1105-1121.
- Turner AJ. 2015. ACE2 cell biology, regulation, and physiological functions. In *The protective arm of the renin angiotensin system (RAS): Functional aspects and therapeutic implications*. Unger T, Steckelings UM, dos Santos RAS (ed). Academic Press, Cambridge, MA, USA. pp 185-189.
- van Lith R, Ameer GA. 2016. Antioxidant polymers as biomaterial. In *Oxidative stress and biomaterials*. Dziubla T, Allan Butterfield D (ed). Academic Press, Cambridge, MA, USA. pp 251-296.
- Walker JM. 2009. The bicinchoninic acid (BCA) assay for protein quantitation. In *The protein protocols handbook*. Walker

- JM (ed). Humana Press, Totowa, NJ, USA. pp 11-15.
- Wang J, Guo M, Wang Q, Dong J, Lu S, Lyu B, Ma X. 2021. Antioxidant activities of peptides derived from mutton ham, Xuanwei ham and Jinhua ham. *Food Res Int* 142:110195.
- Wang L, Yin YL, Liu XZ, Shen P, Zheng YG, Lan XR, Lu CB, Wang JZ. 2020. Current understanding of metal ions in the pathogenesis of Alzheimer's disease. *Transl Neurodegener* 9:1-13.
- Wang X, Yu H, Xing R, Li P. 2017. Characterization, preparation, and purification of marine bioactive peptides. *BioMed Res Int* 2017:9746720.
- Wen S, Zhou G, Song S, Xu X, Voglmeir J, Liu L, Zhao F, Li M, Li L, Yu X, Bai Y, Li C. 2015. Discrimination of *in vitro* and *in vivo* digestion products of meat proteins from pork, beef, chicken, and fish. *Proteomics* 15:3688-3698.
- Whitcomb DC, Lowe ME. 2007. Human pancreatic digestive enzymes. *Dig Dis Sci* 52:1-17.
- Williams B, Mancia G, Spiering W, Agabiti Rosei E, Azizi M, Burnier M, Clement DL, Coca A, de Simone G, Dominiczak A, Kahan T, Mahfoud F, Redon J, Ruilope L, Zanchetti A, Kerins M, Kjeldsen SE, Kreutz R, Laurent S, Lip GYH, McManus R, Narkiewicz K, Ruschitzka F, Schmieder RE, Shlyakhto E, Tsioufis C, Aboyans V, Desormais I. 2018. 2018 ESC/ESH guidelines for the management of arterial hypertension: The task force for the management of arterial hypertension of the European Society of Cardiology (ESC) and the European Society of Hypertension (ESH). *Eur Heart J* 39:3021-3104.
- World Health Organization [WHO]. 2013. A global brief on hypertension: Silent killer, global public health crisis: World Health Day 2013. WHO, Geneva, Switzerland. Report No. WHO/DCO/WHO/2013.2.
- World Health Organization [WHO]. 2021. Hypertension. Available from: <https://www.who.int/news-room/fact-sheets/detail/hypertension>. Accessed at Sep 19, 2022.
- Wu J, Liao W, Udenigwe CC. 2017. Revisiting the mechanisms of ACE inhibitory peptides from food proteins. *Trends Food Sci Technol* 69:214-219.
- Xing L, Hu Y, Hu H, Ge Q, Zhou G, Zhang W. 2016. Purification and identification of antioxidative peptides from dry-cured Xuanwei ham. *Food Chem* 194:951-958.
- Xing L, Liu R, Cao S, Zhang W, Guanghong Z. 2019. Meat protein based bioactive peptides and their potential functional activity: A review. *Int J Food Sci Technol* 54:1956-1966.
- Xing L, Liu R, Gao X, Zheng J, Wang C, Zhou G, Zhang W. 2018. The proteomics homology of antioxidant peptides extracted from dry-cured Xuanwei and Jinhua ham. *Food Chem* 266:420-426.
- Yan W, Lin G, Zhang R, Liang Z, Wu W. 2020. Studies on the bioactivities and molecular mechanism of antioxidant peptides by 3D-QSAR, *in vitro* evaluation and molecular dynamic simulations. *Food Funct* 11:3043-3052.
- Young IS, Woodside JV. 2001. Antioxidants in health and disease. *J Clin Pathol* 54:176-186.
- Zhong Y, Shahidi F. 2015. Methods for the assessment of antioxidant activity in foods. In *Handbook of antioxidants for food preservation*. Shahidi F (ed). Woodhead, Sawston, UK. pp 287-333.
- Zhu CZ, Zhang WG, Zhou GH, Xu XL. 2016. Identification of antioxidant peptides of Jinhua ham generated in the products and through the simulated gastrointestinal digestion system. *J Sci Food Agric* 96:99-108.
- Zhuo JL, Ferrao FM, Zheng Y, Li XC. 2013. New frontiers in the intrarenal renin-angiotensin system: A critical review of classical and new paradigms. *Front Endocrinol* 4:166.